

# Announcements

Wednesday, August 30

- ▶ WeBWork due on **Friday** at 11:59pm.
- ▶ The first quiz is on Friday, during recitation. It covers **through Monday's material**.
  - ▶ Quizzes mostly test your understanding of the homework.
  - ▶ Quizzes last 10 minutes. Books, calculators, etc. are not allowed.
  - ▶ There will generally be a quiz every Friday when there's no midterm.
  - ▶ Check the schedule if you want to know what will be covered.
- ▶ My office is Skiles 244 and my office hours are Monday, 1–3pm and Tuesday, 9–11am.
- ▶ Your TAs have office hours too. You can go to any of them. Details on the website.
- ▶ Many other resources are also contained in the “Help” tab of the master website. This includes Math Lab (not to be confused with MyMathLab), a free one-on-many tutoring service, open for many hours most days, provided by the School of Math.

# Reduced Row Echelon Form

## Theorem

Every matrix is row equivalent to one and only one matrix in reduced row echelon form.

We'll give an algorithm, called **row reduction**, which demonstrates that every matrix is row equivalent to *at least one* matrix in reduced row echelon form.

**Note:** Like echelon forms, the row reduction algorithm does not care if a column is augmented: ignore the vertical line when row reducing.

The uniqueness statement is interesting—it means that, no matter *how* you row reduce, you *always* get the same matrix in reduced row echelon form. (Assuming you only do the three legal row operations.) (And you don't make any arithmetic errors.)

Maybe you can figure out why it's true!

# Row Reduction Algorithm

**Step 1a** Swap the 1st row with a lower one so a leftmost nonzero entry is in 1st row (if necessary).

**Step 1b** Scale 1st row so that its leading entry is equal to 1.

**Step 1c** Use row replacement so all entries below this 1 are 0.

**Step 2a** Swap the 2nd row with a lower one so that the leftmost nonzero entry is in 2nd row.

**Step 2b** Scale 2nd row so that its leading entry is equal to 1.

**Step 2c** Use row replacement so all entries below this 1 are 0.

**Step 3a** Swap the 3rd row with a lower one so that the leftmost nonzero entry is in 3rd row.

etc.

**Last Step** Use row replacement to clear all entries above the pivots, starting with the last pivot (to make life easier).

**Example**

$$\left( \begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

[animated]

# Row Reduction

## Example

$$\left( \begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Step 1a: Row swap to make this nonzero.

$$R_1 \leftrightarrow R_2$$

$$R_1 = R_1 \div 2$$

$$R_3 = R_3 - 3R_1$$

Optional: swap rows 2 and 3 to make Step 2b easier later on.

$$R_2 \leftrightarrow R_3$$

$$\left( \begin{array}{ccc|c} 2 & 4 & 6 & 12 \\ 0 & -7 & -4 & 2 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Step 1b: Scale to make this 1.

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Step 1c: Subtract a multiple of the first row to clear this.

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 0 & -5 & -10 & -20 \end{array} \right)$$

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -5 & -10 & -20 \\ 0 & -7 & -4 & 2 \end{array} \right)$$

# Row Reduction

Example, continued

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -5 & -10 & -20 \\ 0 & -7 & -4 & 2 \end{array} \right)$$

Step 2a: This is already nonzero.

Step 2b: Scale to make this 1.

(There are no fractions because of the optional step before.)

$$\underbrace{R_2 = R_2 \div -5}_{\rightsquigarrow}$$

$$\underbrace{R_3 = R_3 + 7R_2}_{\rightsquigarrow}$$

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & 1 & 2 & 4 \\ 0 & -7 & -4 & 2 \end{array} \right)$$

Step 2c: Add 7 times the second row to clear this.

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 10 & 30 \end{array} \right)$$

**Note:** Step 2 never messes up the first (nonzero) column of the matrix, because it looks like this:

"Active" row  $\rightarrow$  
$$\left( \begin{array}{ccc|c} 1 & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \end{array} \right)$$

# Row Reduction

Example, continued

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 10 & 30 \end{array} \right)$$

Step 3a: This is already nonzero.

Step 3b: Scale to make this 1.

$$R_3 = R_3 \div 10$$

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 1 & 3 \end{array} \right)$$

**Note:** Step 3 never messes up the columns to the left.

**Note:** The matrix is now in row echelon form!

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 1 & 3 \end{array} \right)$$

Last step: Add multiples of the third row to clear these.

$$R_2 = R_2 - 2R_3$$

$$R_1 = R_1 - 3R_3$$

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{array} \right)$$

$$\left( \begin{array}{ccc|c} 1 & 0 & 0 & -3 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{array} \right)$$

Last step: Add  $-2$  times the third row to clear this.

$$R_1 = R_1 - 2R_2$$

$$\left( \begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{array} \right)$$

# Row Reduction

Example, continued

Success! The reduced row echelon form is

$$\left( \begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{array} \right) \implies \begin{cases} x & = & 1 \\ y & = & -2 \\ z & = & 3 \end{cases}$$

# Recap

Get a 1 here

$$\begin{pmatrix} \star & \star & \star & \star \\ \star & \star & \star & \star \\ \star & \star & \star & \star \\ \star & \star & \star & \star \end{pmatrix}$$

Clear down

$$\begin{pmatrix} 1 & \star & \star & \star \\ \star & \star & \star & \star \\ \star & \star & \star & \star \\ \star & \star & \star & \star \end{pmatrix}$$

Get a 1 here

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & \star & \star & \star \\ 0 & \star & \star & \star \\ 0 & \star & \star & \star \end{pmatrix}$$

Clear down

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & 1 & \star & \star \\ 0 & \star & \star & \star \\ 0 & \star & \star & \star \end{pmatrix}$$

(maybe these are already zero)

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & 1 & \star & \star \\ 0 & 0 & 0 & \star \\ 0 & 0 & 0 & \star \end{pmatrix}$$

Get a 1 here

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & 1 & \star & \star \\ 0 & 0 & 0 & \star \\ 0 & 0 & 0 & \star \end{pmatrix}$$

Clear down

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & 1 & \star & \star \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & \star \end{pmatrix}$$

Matrix is in REF

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & 1 & \star & \star \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Clear up

$$\begin{pmatrix} 1 & \star & \star & \star \\ 0 & 1 & \star & \star \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Clear up

$$\begin{pmatrix} 1 & \star & \star & 0 \\ 0 & 1 & \star & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Matrix is in RREF

$$\begin{pmatrix} 1 & 0 & \star & 0 \\ 0 & 1 & \star & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Profit?



# Row Reduction

Another example

The linear system

$$2x + 10y = -1$$

$$3x + 15y = 2$$

gives rise to the matrix

$$\left( \begin{array}{cc|c} 2 & 10 & -1 \\ 3 & 15 & 2 \end{array} \right).$$

Let's row reduce it: [\[interactive row reducer\]](#)

$$\left( \begin{array}{cc|c} 2 & 10 & -1 \\ 3 & 15 & 2 \end{array} \right) \xrightarrow{R_1 = R_1 \div 2} \left( \begin{array}{cc|c} 1 & 5 & -\frac{1}{2} \\ 3 & 15 & 2 \end{array} \right) \quad \text{(Step 1b)}$$

$$\xrightarrow{R_2 = R_2 - 3R_1} \left( \begin{array}{cc|c} 1 & 5 & -\frac{1}{2} \\ 0 & 0 & \frac{7}{2} \end{array} \right) \quad \text{(Step 1c)}$$

$$\xrightarrow{R_2 = R_2 \times \frac{2}{7}} \left( \begin{array}{cc|c} 1 & 5 & -\frac{1}{2} \\ 0 & 0 & 1 \end{array} \right) \quad \text{(Step 2b)}$$

$$\xrightarrow{R_1 = R_1 + \frac{1}{2}R_2} \left( \begin{array}{cc|c} 1 & 5 & 0 \\ 0 & 0 & 1 \end{array} \right) \quad \text{(Step 2c)}$$

The row reduced matrix

$$\left( \begin{array}{cc|c} 1 & 5 & 0 \\ 0 & 0 & 1 \end{array} \right)$$

corresponds to the  
*inconsistent* system

$$\begin{aligned} x + 5y &= 0 \\ 0 &= 1. \end{aligned}$$

# Inconsistent Matrices

## Question

What does an augmented matrix in reduced row echelon form look like, if its system of linear equations is inconsistent?

Answer:

$$\left( \begin{array}{cccc|c} 1 & 0 & * & * & 0 \\ 0 & 1 & * & * & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right)$$

An augmented matrix corresponds to an inconsistent system of equations if and only if *the last* (i.e., the augmented) *column is a pivot column*.

## Another Example

The linear system

$$\begin{aligned} 2x + y + 12z &= 1 \\ x + 2y + 9z &= -1 \end{aligned} \quad \text{gives rise to the matrix} \quad \left( \begin{array}{ccc|c} 2 & 1 & 12 & 1 \\ 1 & 2 & 9 & -1 \end{array} \right).$$

Let's row reduce it: [\[interactive row reducer\]](#)

$$\begin{aligned} \left( \begin{array}{ccc|c} 2 & 1 & 12 & 1 \\ 1 & 2 & 9 & -1 \end{array} \right) & \begin{array}{l} R_1 \leftrightarrow R_2 \\ \text{~~~~~} \end{array} & \left( \begin{array}{ccc|c} 1 & 2 & 9 & -1 \\ 2 & 1 & 12 & 1 \end{array} \right) & \text{(Optional)} \\ & \begin{array}{l} R_2 = R_2 - 2R_1 \\ \text{~~~~~} \end{array} & \left( \begin{array}{ccc|c} 1 & 2 & 9 & -1 \\ 0 & -3 & -6 & 3 \end{array} \right) & \text{(Step 1c)} \\ & \begin{array}{l} R_2 = R_2 \div -3 \\ \text{~~~~~} \end{array} & \left( \begin{array}{ccc|c} 1 & 2 & 9 & -1 \\ 0 & 1 & 2 & -1 \end{array} \right) & \text{(Step 2b)} \\ & \begin{array}{l} R_1 = R_1 - 2R_2 \\ \text{~~~~~} \end{array} & \left( \begin{array}{ccc|c} 1 & 0 & 5 & 1 \\ 0 & 1 & 2 & -1 \end{array} \right) & \text{(Step 2c)} \end{aligned}$$

The row reduced matrix

$$\left( \begin{array}{ccc|c} 1 & 0 & 5 & 1 \\ 0 & 1 & 2 & -1 \end{array} \right) \quad \text{corresponds to the linear system} \quad \begin{cases} x + 5z = 1 \\ y + 2z = -1 \end{cases}$$

## Another Example

Continued

The system

$$x + 5z = 1$$

$$y + 2z = -1$$

comes from a matrix in reduced row echelon form. Are we done? Is the system solved?

Yes! Rewrite:

$$x = 1 - 5z$$

$$y = -1 - 2z$$

For any value of  $z$ , there is exactly one value of  $x$  and  $y$  that makes the equations true. But  $z$  can be *anything we want!*

So we have found the solution set: it is all values  $x, y, z$  where

$$x = 1 - 5z$$

$$y = -1 - 2z \quad \text{for } z \text{ any real number.}$$

$$(z = z)$$

This is called the **parametric form** for the solution. [\[interactive picture\]](#)

For instance,  $(1, -1, 0)$  and  $(-4, -3, 1)$  are solutions.

# Free Variables

## Definition

Consider a *consistent* linear system of equations in the variables  $x_1, \dots, x_n$ . Let  $A$  be a row echelon form of the matrix for this system.

We say that  $x_i$  is a **free variable** if its corresponding column in  $A$  is *not* a pivot column.

### Important

1. You can choose *any value* for the free variables in a (consistent) linear system.
2. Free variables come from *columns without pivots* in a matrix in row echelon form.

In the previous example,  $z$  was free because the reduced row echelon form matrix was

$$\left( \begin{array}{ccc|c} 1 & 0 & 5 & 4 \\ 0 & 1 & 2 & -1 \end{array} \right).$$

In this matrix:

$$\left( \begin{array}{cccc|c} 1 & \star & 0 & \star & \star \\ 0 & 0 & 1 & \star & \star \end{array} \right)$$

the free variables are  $x_2$  and  $x_4$ . (What about the last column?)

## One More Example

The reduced row echelon form of the matrix for a linear system in  $x_1, x_2, x_3, x_4$  is

$$\left( \begin{array}{cccc|c} 1 & 0 & 0 & 3 & 2 \\ 0 & 0 & 1 & 4 & -1 \end{array} \right)$$

The free variables are  $x_2$  and  $x_4$ : they are the ones whose columns are *not* pivot columns.

This translates into the system of equations

$$\begin{cases} x_1 & + 3x_4 = 2 \\ & x_3 + 4x_4 = -1 \end{cases} \implies \boxed{\begin{array}{l} x_1 = 2 - 3x_4 \\ x_3 = -1 - 4x_4 \end{array}}$$

What happened to  $x_2$ ? What is it allowed to be? Anything! The general solution is

$$(x_1, x_2, x_3, x_4) = (2 - 3x_4, x_2, -1 - 4x_4, x_4)$$

for any values of  $x_2$  and  $x_4$ . For instance,  $(2, 0, -1, 0)$  is a solution ( $x_2 = x_4 = 0$ ), and  $(5, 1, 3, -1)$  is a solution ( $x_2 = 1, x_4 = -1$ ).

The boxed equation is called the **parametric form** of the general solution to the system of equations. It is obtained by moving all free variables to the right-hand side of the  $=$ .

## Yet Another Example

The linear system

$$x + y + z = 1 \quad \text{has matrix form} \quad (1 \ 1 \ 1 \mid 1).$$

This is in reduced row echelon form. The free variables are  $y$  and  $z$ . The parametric form of the general solution is

$$x = 1 - y - z.$$

Rearranging:

$$(x, y, z) = (1 - y - z, y, z),$$

where  $y$  and  $z$  are arbitrary real numbers. This was an example in the second lecture!

[interactive]

Poll

Is it possible for a system of linear equations to have exactly two solutions?



## Summary

There are *three possibilities* for the reduced row echelon form of the augmented matrix of a linear system.

1. The last column is a pivot column.

In this case, the system is *inconsistent*. There are zero solutions, i.e. the solution set is *empty*. Picture:

$$\left( \begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right)$$

2. Every column except the last column is a pivot column.

In this case, the system has a *unique solution*. Picture:

$$\left( \begin{array}{ccc|c} 1 & 0 & 0 & * \\ 0 & 1 & 0 & * \\ 0 & 0 & 1 & * \end{array} \right)$$

3. The last column is not a pivot column, and some other column isn't either.

In this case, the system has *infinitely many* solutions, corresponding to the infinitely many possible values of the free variable(s). Picture:

$$\left( \begin{array}{cccc|c} 1 & * & 0 & * & * \\ 0 & 0 & 1 & * & * \end{array} \right)$$