

Announcements

Monday, October 30

- ▶ WeBWork 3.1, 3.2 are due Wednesday at 11:59pm.
- ▶ The quiz on Friday covers §§3.1, 3.2.
- ▶ My office is Skiles 244. Rabinoffice hours are Monday, 1–3pm and Tuesday, 9–11am.

Chapter 5

Eigenvalues and Eigenvectors

Section 5.1

Eigenvectors and Eigenvalues

A Biology Question

Motivation

In a population of rabbits:

1. half of the newborn rabbits survive their first year;
2. of those, half survive their second year;
3. their maximum life span is three years;
4. rabbits have 0, 6, 8 baby rabbits in their three years, respectively.

If you know the population one year, what is the population the next year?

f_n = first-year rabbits in year n

s_n = second-year rabbits in year n

t_n = third-year rabbits in year n

The rules say:

$$\begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \begin{pmatrix} f_n \\ s_n \\ t_n \end{pmatrix} = \begin{pmatrix} f_{n+1} \\ s_{n+1} \\ t_{n+1} \end{pmatrix}.$$

Let $A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix}$ and $v_n = \begin{pmatrix} f_n \\ s_n \\ t_n \end{pmatrix}$. Then $A v_n = v_{n+1}$. ← difference equation

A Biology Question

Continued

If you know v_0 , what is v_{10} ?

$$v_{10} = Av_9 = AA v_8 = \dots = A^{10} v_0.$$

This makes it easy to compute examples by computer:

v_0	v_{10}	v_{11}
$\begin{pmatrix} 3 \\ 7 \\ 9 \end{pmatrix}$	$\begin{pmatrix} 30189 \\ 7761 \\ 1844 \end{pmatrix}$	$\begin{pmatrix} 61316 \\ 15095 \\ 3881 \end{pmatrix}$
$\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$	$\begin{pmatrix} 9459 \\ 2434 \\ 577 \end{pmatrix}$	$\begin{pmatrix} 19222 \\ 4729 \\ 1217 \end{pmatrix}$
$\begin{pmatrix} 4 \\ 7 \\ 8 \end{pmatrix}$	$\begin{pmatrix} 28856 \\ 7405 \\ 1765 \end{pmatrix}$	$\begin{pmatrix} 58550 \\ 14428 \\ 3703 \end{pmatrix}$

What do you notice about these numbers?

1. Eventually, each segment of the population doubles every year: $Av_n = v_{n+1} = 2v_n$.
2. The ratios get close to (16 : 4 : 1):

$$v_n = (\text{scalar}) \cdot \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix}.$$

Translation: 2 is an eigenvalue, and $\begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix}$ is an eigenvector!

Eigenvectors and Eigenvalues

Definition

Let A be an $n \times n$ matrix.

Eigenvalues and eigenvectors are only for square matrices.

1. An **eigenvector** of A is a *nonzero* vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} . In other words, Av is a multiple of v .
2. An **eigenvalue** of A is a number λ in \mathbf{R} such that the equation $Av = \lambda v$ has a *nontrivial* solution.

If $Av = \lambda v$ for $v \neq 0$, we say λ is the **eigenvalue for** v , and v is an **eigenvector for** λ .

Note: Eigenvectors are by definition nonzero. Eigenvalues may be equal to zero.

This is the most important definition in the course.

Verifying Eigenvectors

Example

$$A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \quad v = \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix}$$

Multiply:

$$Av = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix} = \begin{pmatrix} 32 \\ 8 \\ 2 \end{pmatrix} = 2v$$

Hence v is an eigenvector of A , with eigenvalue $\lambda = 2$.

Example

$$A = \begin{pmatrix} 2 & 2 \\ -4 & 8 \end{pmatrix} \quad v = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Multiply:

$$Av = \begin{pmatrix} 2 & 2 \\ -4 & 8 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \end{pmatrix} = 4v$$

Hence v is an eigenvector of A , with eigenvalue $\lambda = 4$.

Poll

Which of the vectors

A. $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ B. $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ C. $\begin{pmatrix} -1 \\ 1 \end{pmatrix}$ D. $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$ E. $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$

are eigenvectors of the matrix $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$?

What are the eigenvalues?

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

eigenvector with eigenvalue 2

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = 0 \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

eigenvector with eigenvalue 0

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} -1 \\ 1 \end{pmatrix} = 0 \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

eigenvector with eigenvalue 0

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \end{pmatrix}$$

not an eigenvector

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

is never an eigenvector

Verifying Eigenvalues

Question: Is $\lambda = 3$ an eigenvalue of $A = \begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix}$?

In other words, does $Av = 3v$ have a nontrivial solution?

... does $Av - 3v = 0$ have a nontrivial solution?

... does $(A - 3I)v = 0$ have a nontrivial solution?

We know how to answer that! Row reduction!

$$A - 3I = \begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix} - 3 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} -1 & -4 \\ -1 & -4 \end{pmatrix}$$

Row reduce:

$$\begin{pmatrix} -1 & -4 \\ -1 & -4 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 1 & 4 \\ 0 & 0 \end{pmatrix}$$

Parametric form: $x = -4y$; parametric vector form: $\begin{pmatrix} x \\ y \end{pmatrix} = y \begin{pmatrix} -4 \\ 1 \end{pmatrix}$.

Does there exist an eigenvector with eigenvalue $\lambda = 3$? Yes! Any nonzero multiple of $\begin{pmatrix} -4 \\ 1 \end{pmatrix}$. Check:

$$\begin{pmatrix} 2 & -4 \\ -1 & -1 \end{pmatrix} \begin{pmatrix} -4 \\ 1 \end{pmatrix} = \begin{pmatrix} -12 \\ 3 \end{pmatrix} = 3 \begin{pmatrix} -4 \\ 1 \end{pmatrix}. \quad \checkmark$$

Eigenspaces

Definition

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . The λ -**eigenspace** of A is the set of all eigenvectors of A with eigenvalue λ , plus the zero vector:

$$\begin{aligned}\lambda\text{-eigenspace} &= \{v \text{ in } \mathbf{R}^n \mid Av = \lambda v\} \\ &= \{v \text{ in } \mathbf{R}^n \mid (A - \lambda I)v = 0\} \\ &= \text{Nul}(A - \lambda I).\end{aligned}$$

Since the λ -eigenspace is a null space, it is a *subspace* of \mathbf{R}^n .

How do you find a basis for the λ -eigenspace? Parametric vector form!

Eigenspaces

Example

Find a basis for the 2-eigenspace of

$$A = \begin{pmatrix} 7/2 & 0 & 3 \\ -3/2 & 2 & -3 \\ -3/2 & 0 & -1 \end{pmatrix}.$$

$$A - 2I = \begin{pmatrix} 3/2 & 0 & 3 \\ -3/2 & 0 & -3 \\ -3/2 & 0 & -3 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\begin{array}{l} \text{parametric} \\ \text{form} \end{array} \xrightarrow{\text{~~~~~}} x = -2z$$

$$\begin{array}{l} \text{parametric vector} \\ \text{form} \end{array} \xrightarrow{\text{~~~~~}} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = y \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + z \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix}$$

$$\begin{array}{l} \text{basis} \end{array} \xrightarrow{\text{~~~~~}} \left\{ \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix} \right\}$$

Eigenspaces

Example

Find a basis for the $\frac{1}{2}$ -eigenspace of

$$A = \begin{pmatrix} 7/2 & 0 & 3 \\ -3/2 & 2 & -3 \\ -3/2 & 0 & -1 \end{pmatrix}.$$

$$A - \frac{1}{2}I = \begin{pmatrix} 3 & 0 & 3 \\ -3/2 & 3/2 & -3 \\ -3/2 & 0 & -3/2 \end{pmatrix} \xrightarrow{\text{row reduce}} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\xrightarrow{\text{parametric form}} \begin{cases} x = -z \\ y = z \end{cases}$$

$$\xrightarrow{\text{parametric vector form}} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = z \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}$$

$$\xrightarrow{\text{basis}} \left\{ \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix} \right\}$$


Eigenspaces

Geometry

Eigenvectors, geometrically

An eigenvector of a matrix A is a nonzero vector v such that:

- ▶ Av is a multiple of v , which means
- ▶ Av is collinear with v , which means
- ▶ Av and v are *on the same line*.



v is an eigenvector


w is not an eigenvector

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors
(vectors that don't move off their line)?

v is an eigenvector with eigenvalue -1 .


[interactive]

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors
(vectors that don't move off their line)?

w is an eigenvector with eigenvalue 1.


[interactive]

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors
(vectors that don't move off their line)?

u is *not* an eigenvector.


[interactive]

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors
(vectors that don't move off their line)?

Neither is z .


[interactive]

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors (vectors that don't move off their line)?

The 1-eigenspace is L
(all the vectors x where $Ax = x$).


[interactive]

Eigenspaces

Geometry; example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be reflection over the line L defined by $y = -x$, and let A be the matrix for T .

Question: What are the eigenvalues and eigenspaces of A ? No computations!



Does anyone see any eigenvectors (vectors that don't move off their line)?

The (-1) -eigenspace is **the line $y = x$** (all the vectors x where $Ax = -x$).

[interactive]

Eigenspaces


Geometry; example

$$A = \begin{pmatrix} 7/2 & 0 & 3 \\ -3/2 & 2 & -3 \\ -3/2 & 0 & -1 \end{pmatrix}.$$

Before we computed bases for the 2-eigenspace and the $1/2$ -eigenspace:

$$\text{2-eigenspace: } \left\{ \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix} \right\} \quad \frac{1}{2}\text{-eigenspace: } \left\{ \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix} \right\}$$

Hence the 2-eigenspace is a plane and the $1/2$ -eigenspace is a line.



[interactive]

Eigenspaces

Summary

Let A be an $n \times n$ matrix and let λ be a number.

1. λ is an eigenvalue of A if and only if $(A - \lambda I)x = 0$ has a nontrivial solution, if and only if $\text{Nul}(A - \lambda I) \neq \{0\}$.
2. In this case, finding a basis for the λ -eigenspace of A means finding a basis for $\text{Nul}(A - \lambda I)$ as usual, i.e. by finding the parametric vector form for the general solution to $(A - \lambda I)x = 0$.
3. The eigenvectors with eigenvalue λ are the nonzero elements of $\text{Nul}(A - \lambda I)$, i.e. the nontrivial solutions to $(A - \lambda I)x = 0$.

The Eigenvalues of a Triangular Matrix are the Diagonal Entries

We've seen that finding eigenvectors for a given eigenvalue is a row reduction problem.

Finding all of the eigenvalues of a matrix *is not a row reduction problem!* We'll see how to do it in general next time. For now:

Fact: The eigenvalues of a triangular matrix are the diagonal entries.

Why? $\text{Nul}(A - \lambda I) \neq \{0\}$ if and only if $A - \lambda I$ is not invertible, if and only if $\det(A - \lambda I) = 0$.

$$\begin{pmatrix} 3 & 4 & 1 & 2 \\ 0 & -1 & -2 & 7 \\ 0 & 0 & 8 & 12 \\ 0 & 0 & 0 & -3 \end{pmatrix} - \lambda I_4 = \begin{pmatrix} 3 - \lambda & 4 & 1 & 2 \\ 0 & -1 - \lambda & -2 & 7 \\ 0 & 0 & 8 - \lambda & 12 \\ 0 & 0 & 0 & -3 - \lambda \end{pmatrix}.$$

The determinant is $(3 - \lambda)(-1 - \lambda)(8 - \lambda)(-3 - \lambda)$, which is zero exactly when $\lambda = 3, -1, 8,$ or -3 .

A Matrix is Invertible if and only if Zero is not an Eigenvalue

Fact: A is invertible if and only if 0 is not an eigenvalue of A .

Why?

0 is an eigenvalue of $A \iff Ax = 0x$ has a nontrivial solution

$\iff Ax = 0$ has a nontrivial solution

$\iff A$ is not invertible.

invertible matrix theorem



Eigenvectors with Distinct Eigenvalues are Linearly Independent

Fact: If v_1, v_2, \dots, v_k are eigenvectors of A with *distinct* eigenvalues $\lambda_1, \dots, \lambda_k$, then $\{v_1, v_2, \dots, v_k\}$ is linearly independent.

Why? If $k = 2$, this says v_2 can't lie on the line through v_1 .

But the line through v_1 is contained in the λ_1 -eigenspace, and v_2 does not have eigenvalue λ_1 .

In general: see Lay, Theorem 2 in §5.1 (or work it out for yourself; it's not too hard).

Consequence: An $n \times n$ matrix has at most n distinct eigenvalues.

Difference Equations

Preview

Let A be an $n \times n$ matrix. Suppose we want to solve $Av_n = v_{n+1}$ for all n . In other words, we want vectors v_0, v_1, v_2, \dots , such that

$$Av_0 = v_1 \quad Av_1 = v_2 \quad Av_2 = v_3 \quad \dots$$

We saw before that $v_n = A^n v_0$. But it is inefficient to multiply by A each time.

If v_0 is an *eigenvector* with eigenvalue λ , then

$$v_1 = Av_0 = \lambda v_0 \quad v_2 = Av_1 = \lambda v_1 = \lambda^2 v_0 \quad v_3 = Av_2 = \lambda v_2 = \lambda^3 v_0.$$

In general, $v_n = \lambda^n v_0$. This is *much easier* to compute.

Example

$$A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \quad v_0 = \begin{pmatrix} 16 \\ 4 \\ 1 \end{pmatrix} \quad Av_0 = 2v_0.$$

So if you start with 16 baby rabbits, 4 first-year rabbits, and 1 second-year rabbit, then the population will exactly double every year. In year n , you will have $2^n \cdot 16$ baby rabbits, $2^n \cdot 4$ first-year rabbits, and 2^n second-year rabbits.