

# Announcements

Wednesday, November 01

- ▶ WeBWork 3.1, 3.2 are due today at 11:59pm.
- ▶ The quiz on Friday covers §§3.1, 3.2.
- ▶ My office is Skiles 244. Rabinoffice hours are Monday, 1–3pm and Tuesday, 9–11am.

## Section 5.2

### The Characteristic Equation

# The Invertible Matrix Theorem

## Addenda

We have a couple of new ways of saying “ $A$  is invertible” now:

### The Invertible Matrix Theorem

Let  $A$  be a square  $n \times n$  matrix, and let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$  be the linear transformation  $T(x) = Ax$ . The following statements are equivalent.

1.  $A$  is invertible.
2.  $T$  is invertible.
3.  $A$  is row equivalent to  $I_n$ .
4.  $A$  has  $n$  pivots.
5.  $Ax = 0$  has only the trivial solution.
6. The columns of  $A$  are linearly independent.
7.  $T$  is one-to-one.
8.  $Ax = b$  is consistent for all  $b$  in  $\mathbf{R}^n$ .
9. The columns of  $A$  span  $\mathbf{R}^n$ .
10.  $T$  is onto.
11.  $A$  has a left inverse (there exists  $B$  such that  $BA = I_n$ ).
12.  $A$  has a right inverse (there exists  $B$  such that  $AB = I_n$ ).
13.  $A^T$  is invertible.
14. The columns of  $A$  form a basis for  $\mathbf{R}^n$ .
15.  $\text{Col } A = \mathbf{R}^n$ .
16.  $\dim \text{Col } A = n$ .
17.  $\text{rank } A = n$ .
18.  $\text{Nul } A = \{0\}$ .
19.  $\dim \text{Nul } A = 0$ .
19. The determinant of  $A$  is *not* equal to zero.
20. The number 0 is *not* an eigenvalue of  $A$ .

# The Characteristic Polynomial

Let  $A$  be a square matrix.

$$\begin{aligned}\lambda \text{ is an eigenvalue of } A &\iff Ax = \lambda x \text{ has a nontrivial solution} \\ &\iff (A - \lambda I)x = 0 \text{ has a nontrivial solution} \\ &\iff A - \lambda I \text{ is not invertible} \\ &\iff \det(A - \lambda I) = 0.\end{aligned}$$

This gives us a way to compute the eigenvalues of  $A$ .

## Definition

Let  $A$  be a square matrix. The **characteristic polynomial** of  $A$  is

$$f(\lambda) = \det(A - \lambda I).$$

The **characteristic equation** of  $A$  is the equation

$$f(\lambda) = \det(A - \lambda I) = 0.$$

### Important

The eigenvalues of  $A$  are the roots of the characteristic polynomial  $f(\lambda) = \det(A - \lambda I)$ .

# The Characteristic Polynomial

## Example

Question: What are the eigenvalues of

$$A = \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}?$$

# The Characteristic Polynomial

## Example

**Question:** What is the characteristic polynomial of

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}?$$

What do you notice about  $f(\lambda)$ ?

- ▶ The constant term is  $\det(A)$ , which is zero if and only if  $\lambda = 0$  is a root.
- ▶ The linear term  $-(a + d)$  is the negative of the sum of the diagonal entries of  $A$ .

## Definition

The **trace** of a square matrix  $A$  is  $\text{Tr}(A) =$  sum of the diagonal entries of  $A$ .

### Shortcut

The characteristic polynomial of a  $2 \times 2$  matrix  $A$  is

$$f(\lambda) = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

# The Characteristic Polynomial

## Example

**Question:** What are the eigenvalues of the rabbit population matrix

$$A = \begin{pmatrix} 0 & 6 & 8 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \end{pmatrix}?$$

# Algebraic Multiplicity

## Definition

The **(algebraic) multiplicity** of an eigenvalue  $\lambda$  is its multiplicity as a root of the characteristic polynomial.

This is not a very interesting notion *yet*. It will become interesting when we also define *geometric* multiplicity later.

## Example

In the rabbit population matrix,  $f(\lambda) = -(\lambda - 2)(\lambda + 1)^2$ , so the algebraic multiplicity of the eigenvalue 2 is 1, and the algebraic multiplicity of the eigenvalue  $-1$  is 2.

## Example

In the matrix  $\begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$ ,  $f(\lambda) = (\lambda - (3 - 2\sqrt{2}))(\lambda - (3 + 2\sqrt{2}))$ , so the algebraic multiplicity of  $3 + 2\sqrt{2}$  is 1, and the algebraic multiplicity of  $3 - 2\sqrt{2}$  is 1.



# The Characteristic Polynomial

Poll

**Fact:** If  $A$  is an  $n \times n$  matrix, the characteristic polynomial

$$f(\lambda) = \det(A - \lambda I)$$

turns out to be a polynomial of degree  $n$ , and its roots are the eigenvalues of  $A$ :

$$f(\lambda) = (-1)^n \lambda^n + a_{n-1} \lambda^{n-1} + a_{n-2} \lambda^{n-2} + \cdots + a_1 \lambda + a_0.$$

# The $\mathcal{B}$ -basis

## Review

**Recall:** If  $\{v_1, v_2, \dots, v_m\}$  is a basis for a subspace  $V$  and  $x$  is in  $V$ , then the  **$\mathcal{B}$ -coordinates** of  $x$  are the (unique) coefficients  $c_1, c_2, \dots, c_m$  such that

$$x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m.$$

In this case, the  **$\mathcal{B}$ -coordinate vector** of  $x$  is

$$[x]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix}.$$

**Example:** The vectors

$$v_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

form a basis for  $\mathbf{R}^2$  because they are not collinear.

[\[interactive\]](#)

## Coordinate Systems on $\mathbf{R}^n$

**Recall:** A set of  $n$  vectors  $\{v_1, v_2, \dots, v_n\}$  form a basis for  $\mathbf{R}^n$  if and only if the matrix  $C$  with columns  $v_1, v_2, \dots, v_n$  is invertible.

**Translation:** Let  $\mathcal{B}$  be the basis of columns of  $C$ . Multiplying by  $C$  changes from the  $\mathcal{B}$ -coordinates to the usual coordinates, and multiplying by  $C^{-1}$  changes from the usual coordinates to the  $\mathcal{B}$ -coordinates:

$$[x]_{\mathcal{B}} = C^{-1}x \quad x = C[x]_{\mathcal{B}}.$$


# Similarity

## Definition

Two  $n \times n$  matrices  $A$  and  $B$  are **similar** if there is an invertible  $n \times n$  matrix  $C$  such that

$$A = CBC^{-1}.$$

**What does this mean?** This gives you a different way of thinking about multiplication by  $A$ . Let  $\mathcal{B}$  be the basis of columns of  $C$ .



To compute  $Ax$ , you:

1. multiply  $x$  by  $C^{-1}$  to change to the  $\mathcal{B}$ -coordinates:  $[x]_{\mathcal{B}} = C^{-1}x$
2. multiply this by  $B$ :  $B[x]_{\mathcal{B}} = BC^{-1}x$
3. multiply this by  $C$  to change to usual coordinates:  $Ax = CBC^{-1}x = CB[x]_{\mathcal{B}}$ .


# Similarity

## Definition

Two  $n \times n$  matrices  $A$  and  $B$  are **similar** if there is an invertible  $n \times n$  matrix  $C$  such that

$$A = CBC^{-1}.$$

**What does this mean?** This gives you a different way of thinking about multiplication by  $A$ . Let  $\mathcal{B}$  be the basis of columns of  $C$ .



If  $A = CBC^{-1}$ , then  $A$  and  $B$  do the same thing, but  $B$  operates on the  $\mathcal{B}$ -coordinates, where  $\mathcal{B}$  is the basis of columns of  $C$ .

# Similarity

## Example


$$A = \begin{pmatrix} 1/2 & 3/2 \\ 3/2 & 1/2 \end{pmatrix} \quad B = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix} \quad C = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \quad A = CBC^{-1}.$$

What does  $B$  do geometrically?

It scales the  $x$ -direction by 2 and the  $y$ -direction by  $-1$ .

To compute  $Ax$ , first change to the  $\mathcal{B}$  coordinates, then multiply by  $B$ , then change back to the usual coordinates, where

$$\mathcal{B} = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\} = \{v_1, v_2\} \quad (\text{the columns of } C).$$



# Similarity

## Example


$$A = \begin{pmatrix} 1/2 & 3/2 \\ 3/2 & 1/2 \end{pmatrix} \quad B = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix} \quad C = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \quad A = CBC^{-1}.$$

What does  $B$  do geometrically?

It scales the  $x$ -direction by 2 and the  $y$ -direction by  $-1$ .

To compute  $Ax$ , first change to the  $\mathcal{B}$  coordinates, then multiply by  $B$ , then change back to the usual coordinates, where

$$\mathcal{B} = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\} = \{v_1, v_2\} \quad (\text{the columns of } C).$$



# Similarity

## Example


$$A = \begin{pmatrix} 1/2 & 3/2 \\ 3/2 & 1/2 \end{pmatrix} \quad B = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix} \quad C = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \quad A = CBC^{-1}.$$

What does  $B$  do geometrically?

It scales the  $x$ -direction by 2 and the  $y$ -direction by  $-1$ .

To compute  $Ax$ , first change to the  $\mathcal{B}$  coordinates, then multiply by  $B$ , then change back to the usual coordinates, where

$$\mathcal{B} = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\} = \{v_1, v_2\} \quad (\text{the columns of } C).$$





# Similarity

## Example


$$A = \begin{pmatrix} 1/2 & 3/2 \\ 3/2 & 1/2 \end{pmatrix} \quad B = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix} \quad C = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \quad A = CBC^{-1}.$$

What does  $B$  do geometrically?

It scales the  $x$ -direction by 2 and the  $y$ -direction by  $-1$ .

To compute  $Ax$ , first change to the  $\mathcal{B}$  coordinates, then multiply by  $B$ , then change back to the usual coordinates, where

$$\mathcal{B} = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\} = \{v_1, v_2\} \quad (\text{the columns of } C).$$




# Similarity

## Example

What does  $A$  do geometrically?

- ▶  $B$  scales the  $e_1$ -direction by 2 and the  $e_2$ -direction by  $-1$ .
- ▶  $A$  scales the  $v_1$ -direction by 2 and the  $v_2$ -direction by  $-1$ .

columns of  $C$



[interactive]

Since  $B$  is simpler than  $A$ , this makes it easier to understand  $A$ .

Note the relationship between the eigenvalues/eigenvectors of  $A$  and  $B$ .

# Similarity

Example ( $3 \times 3$ )

$$A = \begin{pmatrix} -3 & -5 & -3 \\ 2 & 4 & 3 \\ -3 & -5 & -2 \end{pmatrix} \quad B = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad C = \begin{pmatrix} -1 & 1 & 0 \\ 1 & -1 & 1 \\ -1 & 0 & 1 \end{pmatrix}$$
$$\implies A = CBC^{-1}.$$

What do  $A$  and  $B$  do geometrically?

- ▶  $B$  scales the  $e_1$ -direction by 2, the  $e_2$ -direction by  $-1$ , and fixes  $e_3$ .
- ▶  $A$  scales the  $v_1$ -direction by 2, the  $v_2$ -direction by  $-1$ , and fixes  $v_3$ .

Here  $v_1, v_2, v_3$  are the columns of  $C$ .

[interactive]

## Similar Matrices Have the Same Characteristic Polynomial

**Fact:** If  $A$  and  $B$  are similar, then they have the same characteristic polynomial.

**Why?** Suppose  $A = CBC^{-1}$ .

**Consequence:** similar matrices have the same eigenvalues! (But different eigenvectors in general.)

### Warning

1. Matrices with the same eigenvalues need not be similar.  
For instance,

$$\begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$$

both only have the eigenvalue 2, but they are not similar.

2. Similarity has nothing to do with row equivalence. For instance,

$$\begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

are row equivalent, but they have different eigenvalues.

## Summary

We did two different things today.

First we talked about characteristic polynomials:

- ▶ We learned to find the eigenvalues of a matrix by computing the roots of the characteristic polynomial  $p(\lambda) = \det(A - \lambda I)$ .
- ▶ For a  $2 \times 2$  matrix  $A$ , the characteristic polynomial is just

$$p(\lambda) = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

- ▶ The **algebraic multiplicity** of an eigenvalue is its multiplicity as a root of the characteristic polynomial.

Then we talked about similar matrices:

- ▶ Two square matrices  $A, B$  of the same size are **similar** if there is an invertible matrix  $C$  such that  $A = CBC^{-1}$ .
- ▶ Geometrically, similar matrices  $A$  and  $B$  do the same thing, except  $B$  operates on the coordinate system  $\mathcal{B}$  defined by the columns of  $C$ :

$$B[x]_{\mathcal{B}} = [Ax]_{\mathcal{B}}.$$

- ▶ This is useful when we can find a similar matrix  $B$  which is *simpler* than  $A$  (e.g., a diagonal matrix).