

Announcements

Monday, November 13

- ▶ The third midterm is on **this Friday, November 17**.
 - ▶ The exam covers §§3.1, 3.2, 5.1, 5.2, 5.3, and 5.5.
 - ▶ About half the problems will be conceptual, and the other half computational.
- ▶ There is a practice midterm posted on the website. It is identical in format to the real midterm (although there may be $\pm 1-2$ problems).
- ▶ Study tips:
 - ▶ There are lots of problems at the end of each section in the book, and at the end of the chapter, for practice.
 - ▶ Make sure to **learn the theorems** and **learn the definitions**, and understand what they mean. There is a reference sheet on the website.
 - ▶ Sit down to do the practice midterm in 50 minutes, with no notes.
 - ▶ Come to office hours!
- ▶ WeBWork 5.3, 5.5 are due Wednesday at 11:59pm.
- ▶ **Double Rabinoffice hours this week:** Monday, 1–3pm; Tuesday, 9–11am; Thursday, 9–11am; Thursday, 12–2pm.
- ▶ Suggest topics for Wednesday's lecture on Piazza.

Geometric Interpretation of Complex Eigenvectors

2×2 case

Theorem

Let A be a 2×2 matrix with complex (non-real) eigenvalue λ , and let v be an eigenvector. Then

$$A = PCP^{-1}$$

where

$$P = \begin{pmatrix} | & | \\ \text{Re } v & \text{Im } v \\ | & | \end{pmatrix} \quad \text{and} \quad C = \begin{pmatrix} \text{Re } \lambda & \text{Im } \lambda \\ -\text{Im } \lambda & \text{Re } \lambda \end{pmatrix}.$$

The matrix C is a composition of rotation by $-\arg(\lambda)$ and scaling by $|\lambda|$:

$$C = \begin{pmatrix} |\lambda| & 0 \\ 0 & |\lambda| \end{pmatrix} \begin{pmatrix} \cos(-\arg(\lambda)) & -\sin(-\arg(\lambda)) \\ \sin(-\arg(\lambda)) & \cos(-\arg(\lambda)) \end{pmatrix}.$$

A 2×2 matrix with complex eigenvalue λ is similar to (rotation by the argument of $\bar{\lambda}$) composed with (scaling by $|\lambda|$). This is multiplication by $\bar{\lambda}$ in $\mathbf{C} \sim \mathbf{R}^2$.

Geometric Interpretation of Complex Eigenvalues


2×2 example

What does $A = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$ do geometrically?

Geometric Interpretation of Complex Eigenvalues

2 × 2 example, continued

$$A = C = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \quad \lambda = 1 - i$$



Geometric Interpretation of Complex Eigenvalues

Another 2×2 example

What does $A = \begin{pmatrix} \sqrt{3} + 1 & -2 \\ 1 & \sqrt{3} - 1 \end{pmatrix}$ do geometrically?

Geometric Interpretation of Complex Eigenvalues


Another 2×2 example, continued

$$A = \begin{pmatrix} \sqrt{3} + 1 & -2 \\ 1 & \sqrt{3} - 1 \end{pmatrix} \quad C = \begin{pmatrix} \sqrt{3} & -1 \\ 1 & \sqrt{3} \end{pmatrix} \quad \lambda = \sqrt{3} - i$$


Geometric Interpretation of Complex Eigenvalues

Another 2×2 example: picture

What does $A = \begin{pmatrix} \sqrt{3} + 1 & -2 \\ 1 & \sqrt{3} - 1 \end{pmatrix}$ do geometrically?



$A = PCP^{-1}$ does the same thing, but with respect to the basis $\mathcal{B} = \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \end{pmatrix} \right\}$ of columns of P :



Classification of 2×2 Matrices with a Complex Eigenvalue


Triptych

Let A be a real matrix with a complex eigenvalue λ . One way to understand the geometry of A is to consider the difference equation $v_{n+1} = Av_n$, i.e. the sequence of vectors v, Av, A^2v, \dots

$$A = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{3}+1 & -2 \\ 1 & \sqrt{3}-1 \end{pmatrix}$$

$$\lambda = \frac{\sqrt{3}-i}{\sqrt{2}}$$

$$|\lambda| > 1$$




“spirals out”
[interactive]

$$A = \frac{1}{2} \begin{pmatrix} \sqrt{3}+1 & -2 \\ 1 & \sqrt{3}-1 \end{pmatrix}$$

$$\lambda = \frac{\sqrt{3}-i}{2}$$

$$|\lambda| = 1$$




“rotates around an ellipse”
[interactive]

$$A = \frac{1}{2\sqrt{2}} \begin{pmatrix} \sqrt{3}+1 & -2 \\ 1 & \sqrt{3}-1 \end{pmatrix}$$

$$\lambda = \frac{\sqrt{3}-i}{2\sqrt{2}}$$

$$|\lambda| < 1$$



“spirals in”
[interactive]

Complex Versus Two Real Eigenvalues

An analogy

Theorem

Let A be a 2×2 matrix with complex eigenvalue $\lambda = a + bi$ (where $b \neq 0$), and let v be an eigenvector. Then

$$A = PCP^{-1}$$

where

$$P = \begin{pmatrix} | & | \\ \text{Re } v & \text{Im } v \\ | & | \end{pmatrix} \quad \text{and} \quad C = (\text{rotation}) \cdot (\text{scaling}).$$

This is very analogous to diagonalization. In the 2×2 case:

Theorem

Let A be a 2×2 matrix with linearly independent eigenvectors v_1, v_2 and associated eigenvalues λ_1, λ_2 . Then

$$A = PDP^{-1}$$

where

$$P = \begin{pmatrix} | & | \\ v_1 & v_2 \\ | & | \end{pmatrix} \quad \text{and} \quad D = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$$

scale x-axis by λ_1
scale y-axis by λ_2

Picture with 2 Real Eigenvalues

We can draw analogous pictures for a matrix with 2 real eigenvalues.

Example: Let $A = \frac{1}{4} \begin{pmatrix} 5 & 3 \\ 3 & 5 \end{pmatrix}$.


This has eigenvalues $\lambda_1 = 2$ and $\lambda_2 = \frac{1}{2}$, with eigenvectors

$$v_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad \text{and} \quad v_2 = \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

Therefore, $A = PDP^{-1}$ with

$$P = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \quad \text{and} \quad D = \begin{pmatrix} 2 & 0 \\ 0 & \frac{1}{2} \end{pmatrix}.$$

So A scales the v_1 -direction by 2 and the v_2 -direction by $\frac{1}{2}$.




Picture with 2 Real Eigenvalues

We can also draw a picture from the perspective a difference equation: in other words, we draw v, Av, A^2v, \dots

$$A = \frac{1}{4} \begin{pmatrix} 5 & 3 \\ 3 & 5 \end{pmatrix} \quad \lambda_1 = 2 \quad \lambda_2 = \frac{1}{2}$$

$|\lambda_1| > 1 \quad |\lambda_2| < 1$



[interactive]

Exercise: Draw analogous pictures when $|\lambda_1|, |\lambda_2|$ are any combination of $< 1, = 1, > 1$.

The Higher-Dimensional Case

Theorem

Let A be a real $n \times n$ matrix. Suppose that for each (real or complex) eigenvalue, the dimension of the eigenspace equals the algebraic multiplicity. Then $A = PCP^{-1}$, where P and C are as follows:

1. C is **block diagonal**, where the blocks are 1×1 blocks containing the real eigenvalues (with their multiplicities), or 2×2 blocks containing the matrices $\begin{pmatrix} \operatorname{Re} \lambda & \operatorname{Im} \lambda \\ -\operatorname{Im} \lambda & \operatorname{Re} \lambda \end{pmatrix}$ for each non-real eigenvalue λ (with multiplicity).
2. The columns of P form bases for the eigenspaces for the real eigenvectors, or come in pairs $(\operatorname{Re} v \ \operatorname{Im} v)$ for the non-real eigenvectors.


For instance, if A is a 3×3 matrix with one real eigenvalue λ_1 with eigenvector v_1 , and one conjugate pair of complex eigenvalues $\lambda_2, \bar{\lambda}_2$ with eigenvectors v_2, \bar{v}_2 , then

$$P = \begin{pmatrix} | & | & | \\ v_1 & \operatorname{Re} v_2 & \operatorname{Im} v_2 \\ | & | & | \end{pmatrix} \quad C = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \operatorname{Re} \lambda_2 & \operatorname{Im} \lambda_2 \\ 0 & -\operatorname{Im} \lambda_2 & \operatorname{Re} \lambda_2 \end{pmatrix}$$

The Higher-Dimensional Case

Example

Let $A = \begin{pmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$. This acts on the xy -plane by rotation by $\pi/4$ and scaling by $\sqrt{2}$. This acts on the z -axis by scaling by 2. Pictures:



Remember, in general $A = PCP^{-1}$ is only *similar* to such a matrix C : so the x, y, z axes have to be replaced by the columns of P .

Summary

- ▶ There is a procedure analogous to diagonalization for matrices with complex eigenvalues. In the 2×2 case, the result is

$$A = PCP^{-1}$$

where C is a rotation-scaling matrix.

- ▶ Multiplication by a 2×2 matrix with a complex eigenvalue λ spirals out if $|\lambda| > 1$, rotates around an ellipse if $|\lambda| = 1$, and spirals in if $|\lambda| < 1$.
- ▶ There are analogous pictures for 2×2 matrices with real eigenvalues.
- ▶ For larger matrices, you have to combine diagonalization and “complex diagonalization”. You get a block diagonal matrix with scalars and rotation-scaling matrices on the diagonal.