

Midterm 3

Review Slides

Coordinate Systems on \mathbf{R}^n

Recall: A set of n vectors $\{v_1, v_2, \dots, v_n\}$ form a basis for \mathbf{R}^n if and only if the matrix C with columns v_1, v_2, \dots, v_n is invertible.

If $x = c_1 v_1 + c_2 v_2 + \dots + c_n v_n$ then

$$[x]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{pmatrix} \implies x = c_1 v_1 + c_2 v_2 + \dots + c_n v_n = C[x]_{\mathcal{B}}.$$

Since $x = C[x]_{\mathcal{B}}$ we have $[x]_{\mathcal{B}} = C^{-1}x$.

Translation: Let \mathcal{B} be the basis of columns of C . Multiplying by C changes from the \mathcal{B} -coordinates to the usual coordinates, and multiplying by C^{-1} changes from the usual coordinates to the \mathcal{B} -coordinates:

$$[x]_{\mathcal{B}} = C^{-1}x \quad x = C[x]_{\mathcal{B}}.$$


Similarity

Definition

Two $n \times n$ matrices A and B are **similar** if there is an invertible $n \times n$ matrix C such that

$$A = CBC^{-1}.$$

What does this mean? This gives you a different way of thinking about multiplication by A . Let \mathcal{B} be the basis of columns of C .



To compute Ax , you:

1. multiply x by C^{-1} to change to the \mathcal{B} -coordinates: $[x]_{\mathcal{B}} = C^{-1}x$
2. multiply this by B : $B[x]_{\mathcal{B}} = BC^{-1}x$
3. multiply this by C to change to usual coordinates: $Ax = CBC^{-1}x = CB[x]_{\mathcal{B}}$.


Similarity

Definition

Two $n \times n$ matrices A and B are **similar** if there is an invertible $n \times n$ matrix C such that

$$A = CBC^{-1}.$$

What does this mean? This gives you a different way of thinking about multiplication by A . Let \mathcal{B} be the basis of columns of C .



If $A = CBC^{-1}$, then A and B do the same thing, but B operates on the \mathcal{B} -coordinates, where \mathcal{B} is the basis of columns of C .

Similarity

Example


$$A = \begin{pmatrix} 1/2 & 3/2 \\ 3/2 & 1/2 \end{pmatrix} \quad B = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix} \quad C = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \quad A = CBC^{-1}.$$

What does B do geometrically?

It scales the x -direction by 2 and the y -direction by -1 .

To compute Ax , first change to the \mathcal{B} coordinates, then multiply by B , then change back to the usual coordinates, where

$$\mathcal{B} = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\} = \{v_1, v_2\} \quad (\text{the columns of } C).$$



Similarity

Example


$$A = \begin{pmatrix} 1/2 & 3/2 \\ 3/2 & 1/2 \end{pmatrix} \quad B = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix} \quad C = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \quad A = CBC^{-1}.$$

What does B do geometrically?

It scales the x -direction by 2 and the y -direction by -1 .

To compute Ax , first change to the \mathcal{B} coordinates, then multiply by B , then change back to the usual coordinates, where

$$\mathcal{B} = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\} = \{v_1, v_2\} \quad (\text{the columns of } C).$$



Similarity

Example


$$A = \begin{pmatrix} 1/2 & 3/2 \\ 3/2 & 1/2 \end{pmatrix} \quad B = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix} \quad C = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \quad A = CBC^{-1}.$$

What does B do geometrically?

It scales the x -direction by 2 and the y -direction by -1 .

To compute Ax , first change to the \mathcal{B} coordinates, then multiply by B , then change back to the usual coordinates, where

$$\mathcal{B} = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\} = \{v_1, v_2\} \quad (\text{the columns of } C).$$



Similarity

Example


$$A = \begin{pmatrix} 1/2 & 3/2 \\ 3/2 & 1/2 \end{pmatrix} \quad B = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix} \quad C = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \quad A = CBC^{-1}.$$

What does B do geometrically?

It scales the x -direction by 2 and the y -direction by -1 .

To compute Ax , first change to the \mathcal{B} coordinates, then multiply by B , then change back to the usual coordinates, where

$$\mathcal{B} = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\} = \{v_1, v_2\} \quad (\text{the columns of } C).$$




Similarity

Example

What does A do geometrically?

- ▶ B scales the e_1 -direction by 2 and the e_2 -direction by -1 .
- ▶ A scales the v_1 -direction by 2 and the v_2 -direction by -1 .

columns of C



Since B is simpler than A , this makes it easier to understand A .

Note the relationship between the eigenvalues/eigenvectors of A and B .

Similarity

Example (3×3)

$$A = \begin{pmatrix} -3 & -5 & -3 \\ 2 & 4 & 3 \\ -3 & -5 & -2 \end{pmatrix} \quad B = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad C = \begin{pmatrix} -1 & 1 & 0 \\ 1 & -1 & 1 \\ -1 & 0 & 1 \end{pmatrix}$$
$$\implies A = CBC^{-1}.$$

What do A and B do geometrically?

- ▶ B scales the e_1 -direction by 2, the e_2 -direction by -1 , and fixes e_3 .
- ▶ A scales the v_1 -direction by 2, the v_2 -direction by -1 , and fixes v_3 .

Here v_1, v_2, v_3 are the columns of C .

[interactive]

Diagonalizable Matrices

Definition

An $n \times n$ matrix A is **diagonalizable** if it is similar to a diagonal matrix:

$$A = PDP^{-1} \quad \text{for } D \text{ diagonal.}$$

The Diagonalization Theorem

An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors.

In this case, $A = PDP^{-1}$ for

$$P = \left(\begin{array}{c|c|c|c} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{array} \right) \quad D = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix},$$

where v_1, v_2, \dots, v_n are linearly independent eigenvectors, and $\lambda_1, \lambda_2, \dots, \lambda_n$ are the corresponding eigenvalues (in the same order).

Corollary

An $n \times n$ matrix with n distinct eigenvalues is diagonalizable.

Algebraic and Geometric Multiplicity

Definition

Let λ be an eigenvalue of a square matrix A . The **geometric multiplicity** of λ is the dimension of the λ -eigenspace.

Theorem

Let λ be an eigenvalue of a square matrix A . Then

$$1 \leq (\text{the geometric multiplicity of } \lambda) \leq (\text{the algebraic multiplicity of } \lambda).$$

Corollary

Let λ be an eigenvalue of a square matrix A . If the algebraic multiplicity of λ is 1, then the geometric multiplicity is also 1.

The Diagonalization Theorem (Alternate Form)

Let A be an $n \times n$ matrix. The following are equivalent:

1. A is diagonalizable.
2. The sum of the geometric multiplicities of the eigenvalues of A equals n .
3. The sum of the algebraic multiplicities of the eigenvalues of A equals n , and *the geometric multiplicity equals the algebraic multiplicity* of each eigenvalue.

Algebraic and Geometric Multiplicity

Example

$$A = \begin{pmatrix} 7/2 & 0 & 3 \\ -3/2 & 2 & -3 \\ -3/2 & 0 & -1 \end{pmatrix}$$

Characteristic polynomial:

$$f(\lambda) = -(\lambda - 2)^2(\lambda - 1/2)$$

Algebraic multiplicity of 2: 2

Algebraic multiplicity of 1/2: 1.

Know already:

- ▶ The 1/2-eigenspace is a line.
- ▶ The 2-eigenspace is a line or a plane.
- ▶ The matrix is diagonalizable if and only if the 2-eigenspace is a plane.

[interactive]

Algebraic and Geometric Multiplicity

Example

$$A - 2I = \begin{pmatrix} 3/2 & 0 & 3 \\ -3/2 & 0 & -3 \\ -3/2 & 0 & -3 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

So a basis for the 2-eigenspace is

$$\left\{ \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right\}.$$

This is a *plane*, so the geometric multiplicity is 2.

$$A - \frac{1}{2}I = \begin{pmatrix} 3 & 0 & 3 \\ -3/2 & 3/2 & -3 \\ -3/2 & 0 & -3/2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{pmatrix}$$

The 1/2-eigenspace is the *line*

$$\text{Span} \left\{ \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \right\}.$$

Diagonalization

Example

The 2-eigenspace has basis $\left\{ \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right\}$.

The 1/2-eigenspace has basis $\left\{ \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \right\}$.

Therefore, $A = PDP^{-1}$ for

$$P = \begin{pmatrix} -2 & 0 & 1 \\ 0 & 1 & -1 \\ 1 & 0 & 1 \end{pmatrix} \quad C = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1/2 \end{pmatrix}.$$

Question: what does A do geometrically?

Diagonalization

Another example

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}.$$

The characteristic polynomial is $(x - 1)^2(x - 2)$.

Algebraic multiplicity of 1: 2

Algebraic multiplicity of 2: 1.

Know already:

- ▶ The 2-eigenspace is a line.
- ▶ The 1-eigenspace is a line or a plane.
- ▶ The matrix is diagonalizable if and only if the 1-eigenspace is a plane.

Check: a basis for the 1-eigenspace is $\{e_1\}$.

Conclusion: A is not diagonalizable!

[interactive]