


## MATH 1553-C MIDTERM EXAMINATION 3

Name		GT Email	@gatech.edu
------	--	----------	-------------

Please **read all instructions** carefully before beginning.

- Please leave your GT ID card on your desk until your TA scans your exam.
- All graded work for Problem  $n$  must appear on the page containing Problem  $n$  or the page labeled “Scratch page for Problem  $n$ ”.
- Each problem is worth 10 points. The maximum score on this exam is 50 points.
- You have 50 minutes to complete this exam.
- There are no aids of any kind (notes, text, etc.) allowed.
- Please show your work.
- You may cite any theorem proved in class or in the sections we covered in the text.
- Good luck!



## Problem 1.

[2 points each]

In this problem, if the statement is always true, circle **T**; otherwise, circle **F**.

All matrices are assumed to have real entries.

- a) **T**    **F**    A  $5 \times 5$  matrix has a real eigenvector.
- b) **T**    **F**    Every diagonalizable  $n \times n$  matrix has  $n$  distinct eigenvalues.
- c) **T**    **F**    If an  $n \times n$  matrix  $A$  has a zero eigenvalue, then  $\text{Nul}(A) \neq \{0\}$ .
- d) **T**    **F**    If  $A$  is an  $n \times n$  matrix, then  $\det(-A) = -\det(A)$ .
- e) **T**    **F**     $\begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$  is similar to  $\begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}$ .

## Solution.


- a) **True:** its characteristic polynomial has odd degree, thus has a real root.
- b) **False:** for instance,  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  is diagonal(izable) but only has one eigenvalue.
- c) **True:** any eigenvector with eigenvalue zero is a in  $\text{Nul}(A)$ .
- d) **False:** this is true if and only if  $n$  is odd. For instance,  $\det\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} = 1$ .
- e) **True:** both are diagonalizable with distinct eigenvalues 1, 2.

## Problem 2.

[2 points each]

*Short answer problems: you need not explain your work.*


- a) What is the area of the triangle in the picture?



- b) Give an example of a  $2 \times 2$  matrix that is invertible but not diagonalizable.

- c) Give an example of two  $2 \times 2$  matrices that have the same characteristic polynomial but are not similar.

- d) Suppose that  $A = P \begin{pmatrix} 1/2 & 0 \\ 0 & -1 \end{pmatrix} P^{-1}$ , where  $P$  has columns  $v_1$  and  $v_2$ . Given  $x$  and  $y$  in the picture below, draw the vectors  $Ax$  and  $Ay$ .



- e) With respect to the picture in (d), find the  $\mathcal{B}$ -coordinates of an eigenvector of  $A$  with eigenvalue  $-1$ , where  $\mathcal{B} = \{v_1, v_2\}$ .

**Solution.**

a) If we double the triangle, we get a parallelogram spanned by

$$v_1 = \begin{pmatrix} 4 \\ 1 \end{pmatrix} \quad \text{and} \quad v_2 = \begin{pmatrix} 2 \\ 4 \end{pmatrix}.$$

The area of the parallelogram is

$$\det \begin{pmatrix} 4 & 2 \\ 1 & 4 \end{pmatrix} = 14.$$

Hence the triangle has area 7.

b)  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ .

c) The matrix  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  in (c) has characteristic polynomial  $(\lambda - 1)^2$ , as does the identity matrix  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ . But the identity matrix is diagonal, and  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  is not diagonalizable, so they are not similar.

d)  $A$  does the same thing as  $D = \begin{pmatrix} 1/2 & 0 \\ 0 & -1 \end{pmatrix}$ , but in the  $v_1, v_2$ -coordinate system. Since  $D$  scales the first coordinate by  $1/2$  and the second coordinate by  $-1$ , hence  $A$  scales the  $v_1$ -coordinate by  $1/2$  and the  $v_2$ -coordinate by  $-1$ .

e)  $A$  scales the  $v_2$ -direction by  $-1$ , so  $v_2$  is a  $-1$ -eigenvector. The  $\mathcal{B}$ -coordinate vector of  $v_2$  is  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ .

### Problem 3.

[2 points each]

Consider the matrix

$$A = \begin{pmatrix} -2\sqrt{3}-1 & 5 \\ -1 & -2\sqrt{3}+1 \end{pmatrix}$$

- Find both complex eigenvalues of  $A$ .
- Find an eigenvector corresponding to each eigenvalue.
- Find an invertible matrix  $P$  and a rotation-scale matrix  $C$  such that  $A = PCP^{-1}$ .
- By what angle does  $C$  rotate?
- Successive multiplication by  $A$ :

spirals in      rotates around an ellipse      spirals out

(circle the best option).

### Solution.

- a) We compute the characteristic polynomial:

$$f(\lambda) = \lambda^2 - \text{Tr}(A)\lambda + \det(A) = \lambda^2 + 4\sqrt{3}\lambda + 16$$

By the quadratic formula,

$$\lambda = \frac{-4\sqrt{3} \pm \sqrt{16 \cdot 3 - 16 \cdot 4}}{2} = -2\sqrt{3} \pm 2i.$$

- b) Let  $\lambda = -2\sqrt{3} - 2i$ . Then

$$A - \lambda I_2 = \begin{pmatrix} 2i-1 & 5 \\ \star & \star \end{pmatrix} \implies v = \begin{pmatrix} 5 \\ 1-2i \end{pmatrix}$$


is an eigenvector. Hence an eigenvector for  $\lambda = -2\sqrt{3} + 2i$  is

$$v = \begin{pmatrix} 5 \\ 1+2i \end{pmatrix}.$$

- c) Using the eigenvalue  $\lambda = -2\sqrt{3} - 2i$  and eigenvector  $v = \begin{pmatrix} 5 \\ 1-2i \end{pmatrix}$ , we can take

$$P = (\text{Re } v \quad \text{Im } v) = \begin{pmatrix} 5 & 0 \\ 1 & 2 \end{pmatrix} \quad C = \begin{pmatrix} \text{Re } \lambda & \text{Im } \lambda \\ -\text{Im } \lambda & \text{Re } \lambda \end{pmatrix} = \begin{pmatrix} -2\sqrt{3} & -2 \\ 2 & -2\sqrt{3} \end{pmatrix}.$$

- d) We need to find the argument of  $\lambda = -2\sqrt{3} - 2i$ . We draw a picture:



$$\theta = \frac{\pi}{6}$$

$$\text{argument} = \pi + \theta = \frac{7\pi}{6}$$

The matrix  $C$  rotates by  $-7\pi/6 = 5\pi/6$ .

- e) The matrix  $C$  scales by a factor of  $|\lambda| = 4 > 1$ , so successive multiplication by  $A$  spirals outward.

### Problem 4.

[10 points]

For which value(s) of  $a$  is  $\lambda = 2$  an eigenvector of this matrix?

$$A = \begin{pmatrix} 3 & -1 & 0 & a \\ a & 3 & 0 & 4 \\ 2 & 0 & 2 & -2 \\ 13 & a & -2 & -7 \end{pmatrix}$$

### Solution.

We need to know which values of  $a$  make the matrix  $A - 2I_4$  noninvertible. We have

$$A - 2I_4 = \begin{pmatrix} 1 & -1 & 0 & a \\ a & 1 & 0 & 4 \\ 2 & 0 & 0 & -2 \\ 13 & a & -2 & -9 \end{pmatrix}.$$

We expand cofactors along the third column, then the second column:

$$\begin{aligned} \det(A - I_4) &= 2 \det \begin{pmatrix} 1 & -1 & a \\ a & 1 & 4 \\ 2 & 0 & -2 \end{pmatrix} \\ &= (2)(1) \det \begin{pmatrix} a & 4 \\ 2 & -2 \end{pmatrix} + (2)(1) \det \begin{pmatrix} 1 & a \\ 2 & -2 \end{pmatrix} \\ &= 2(-2a - 8) + 2(-2 - 2a) = -8a - 20. \end{aligned}$$


This is zero if and only if  $a = -5/2$ .

## Problem 5.

[5 points each]

Let  $A = \begin{pmatrix} 2 & 0 \\ -3 & 1 \end{pmatrix}$ .

- a) Draw all eigenspaces of  $A$ , and label them with the corresponding eigenvalue:



- b) Compute  $A^n$ , where  $n \geq 1$  is any whole number. Your answer should be a single  $2 \times 2$  matrix whose entries are formulas involving  $n$ .

## Solution.

- a) Since  $A$  is upper-triangular, we see immediately that it has eigenvalues 1 and 2. To find an eigenvector with eigenvalue 2, we compute

$$A - 2I = \begin{pmatrix} 0 & 0 \\ -3 & -1 \end{pmatrix} \rightsquigarrow v_1 = \begin{pmatrix} 1 \\ -3 \end{pmatrix}.$$

We eyeball  $v_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$  as an eigenvector with eigenvalue 1. The eigenspaces are the lines through  $v_1$  and  $v_2$ .

- b) We did the computations to diagonalize  $A$  in (a):

$$A = PDP^{-1} \quad P = \begin{pmatrix} 1 & 0 \\ -3 & 1 \end{pmatrix} \quad D = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}.$$

Hence we have

$$\begin{aligned} A^n &= PD^nP^{-1} = \begin{pmatrix} 1 & 0 \\ -3 & 1 \end{pmatrix} \begin{pmatrix} 2^n & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -3 & 1 \end{pmatrix}^{-1} \\ &= \begin{pmatrix} 2^n & 0 \\ -3 \cdot 2^n & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 2^n & 0 \\ 3 - 3 \cdot 2^n & 1 \end{pmatrix}. \end{aligned}$$