

Announcements

Monday, November 27


- ▶ WeBWorK 6.1, 6.2, 6.3 are due Wednesday at 11:59pm.
- ▶ WeBWorK 6.4, 6.5 are posted and will be covered on the final, but they are not graded.
- ▶ No quiz on Friday! But this is the only recitation on chapter 6.
- ▶ My office is Skiles 244. Rabinoffice hours are Monday, 1–3pm and Tuesday, 9–11am.

Section 6.4

The Gram–Schmidt Process

Motivation: Best Approximation

Suppose you measure a data point x which you know for theoretical reasons must lie on a subspace W .



Due to measurement error, though, the measured x is not actually in W . Best approximation: y is the *closest* point to x on W .

How do you know that y is the closest point? The vector from y to x is orthogonal to W : it is in the *orthogonal complement* W^\perp .

Note $x = y + (x - y)$, where y is in W and $x - y$ is in W^\perp . Last time we called this the *orthogonal decomposition* of x :

$$x = x_W + x_{W^\perp} \quad x_W = y \quad x_{W^\perp} = x - y.$$

Orthogonal Decomposition

Review

Recall: If W is a subspace of \mathbf{R}^n , its **orthogonal complement** is

$$W^\perp = \{v \text{ in } \mathbf{R}^n \mid v \text{ is perpendicular to every vector in } W\}$$

Theorem

Every vector x in \mathbf{R}^n can be written as

$$x = x_W + x_{W^\perp}$$


for unique vectors x_W in W and x_{W^\perp} in W^\perp .

The equation $x = x_W + x_{W^\perp}$ is called the **orthogonal decomposition** of x (with respect to W).

The vector x_W is the closest vector to x on W .

[interactive 1]

[interactive 2]



Orthogonal Projections

Review

How do you compute x_W ? (Note $x_{W^\perp} = x - x_W$.)

Recall: a set of nonzero vectors $\{u_1, u_2, \dots, u_m\}$ is **orthogonal** if $u_i \cdot u_j = 0$ when $i \neq j$: each vector is perpendicular to the others.

Definition

Let W be a subspace of \mathbf{R}^n , and let $\{u_1, u_2, \dots, u_m\}$ be an *orthogonal* basis for W . The **orthogonal projection** of a vector x onto W is

$$\text{proj}_W(x) \stackrel{\text{def}}{=} \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i = \frac{x \cdot u_1}{u_1 \cdot u_1} u_1 + \frac{x \cdot u_2}{u_2 \cdot u_2} u_2 + \cdots + \frac{x \cdot u_n}{u_n \cdot u_n} u_n.$$

[interactive]

Let x be a vector and let $x = x_W + x_{W^\perp}$ be its orthogonal decomposition with respect to a subspace W . The following vectors are the same:

- ▶ x_W
- ▶ $\text{proj}_W(x)$
- ▶ The closest vector to x on W


Orthogonal Projection onto a Line

Review

The formula for orthogonal projections is simple when W is a *line*.


Let $L = \text{Span}\{u\}$ be a line in \mathbf{R}^n , and let x be in \mathbf{R}^n . The orthogonal projection of x onto L is the point

$$\text{proj}_L(x) = \frac{x \cdot u}{u \cdot u} u.$$



[interactive]

Example: Compute the orthogonal projection of $x = \begin{pmatrix} -6 \\ 4 \end{pmatrix}$ onto the line L spanned by $u = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$.



Orthogonal Projections

Properties

We can think of orthogonal projection as a *transformation*:

$$\text{proj}_W: \mathbf{R}^n \longrightarrow \mathbf{R}^n \quad x \mapsto \text{proj}_W(x).$$

Theorem

Let W be a subspace of \mathbf{R}^n .

1. proj_W is a *linear* transformation.
2. For every x in W , we have $\text{proj}_W(x) = x$.
3. For every x in W^\perp , we have $\text{proj}_W(x) = 0$.
4. The range of proj_W is W and the null space of proj_W is W^\perp .

Let W be a subspace with orthogonal basis $\mathcal{B} = \{u_1, u_2, \dots, u_m\}$.

For x in W we have $\text{proj}_W(x) = x$, so

$$x = \text{proj}_W(x) = \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i = \frac{x \cdot u_1}{u_1 \cdot u_1} u_1 + \frac{x \cdot u_2}{u_2 \cdot u_2} u_2 + \cdots + \frac{x \cdot u_n}{u_n \cdot u_n} u_n$$
$$\implies [x]_{\mathcal{B}} = \left(\frac{x \cdot u_1}{u_1 \cdot u_1}, \frac{x \cdot u_2}{u_2 \cdot u_2}, \dots, \frac{x \cdot u_m}{u_m \cdot u_m} \right). \quad [\text{interactive}]$$

A Non-Orthogonal Basis

Important: Orthogonal projections require an *orthogonal* basis!

Non-Example: Consider the basis $\mathcal{B} = \{v_1, v_2\}$ of \mathbf{R}^2 , where

$$v_1 = \begin{pmatrix} 2 \\ -1/2 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}.$$

This is not orthogonal: $\begin{pmatrix} 2 \\ -1/2 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 2 \end{pmatrix} = 1 \neq 0$.


Let's try to compute $x = \text{proj}_{\mathbf{R}^2}(x)$ for $x = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ using the basis $\{v_1, v_2\}$:

$$x = \text{proj}_{\mathbf{R}^2}(x) =$$

This does not work!

[interactive]

(compare [orthogonal basis])



Recap

All of the procedures we learned in §§6.2–6.3 require an *orthogonal* basis $\{u_1, u_2, \dots, u_m\}$.

- ▶ Finding the orthogonal projection of a vector x onto the span W of u_1, u_2, \dots, u_m :

$$\text{proj}_W(x) = \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i.$$

- ▶ Finding the orthogonal decomposition of x :

$$x = \text{proj}_W(x) + x_{W^\perp}.$$

- ▶ Finding the \mathcal{B} -coordinates of x :

$$[x]_{\mathcal{B}} = \left(\frac{x \cdot u_1}{u_1 \cdot u_1}, \frac{x \cdot u_2}{u_2 \cdot u_2}, \dots, \frac{x \cdot u_m}{u_m \cdot u_m} \right).$$

Problem: What if your basis isn't orthogonal?

Solution: The Gram–Schmidt process: take any basis and make it orthogonal.

The Gram–Schmidt Process

Procedure

The Gram–Schmidt Process

Let $\{v_1, v_2, \dots, v_m\}$ be a basis for a subspace W of \mathbf{R}^n . Define:

$$1. \quad u_1 = v_1$$

$$2. \quad u_2 = v_2 - \text{proj}_{\text{Span}\{u_1\}}(v_2) = v_2 - \frac{v_2 \cdot u_1}{u_1 \cdot u_1} u_1$$

$$3. \quad u_3 = v_3 - \text{proj}_{\text{Span}\{u_1, u_2\}}(v_3) = v_3 - \frac{v_3 \cdot u_1}{u_1 \cdot u_1} u_1 - \frac{v_3 \cdot u_2}{u_2 \cdot u_2} u_2$$

\vdots

$$m. \quad u_m = v_m - \text{proj}_{\text{Span}\{u_1, u_2, \dots, u_{m-1}\}}(v_m) = v_m - \sum_{i=1}^{m-1} \frac{v_m \cdot u_i}{u_i \cdot u_i} u_i$$

Then $\{u_1, u_2, \dots, u_m\}$ is an *orthogonal* basis for the same subspace W .

Remark


In fact, for every i between 1 and n , the set $\{u_1, u_2, \dots, u_i\}$ is an orthogonal basis for $\text{Span}\{v_1, v_2, \dots, v_i\}$.

The Gram–Schmidt Process

Two vectors

Find an orthogonal basis $\{u_1, u_2\}$ for $W = \text{Span}\{v_1, v_2\}$, where

$$v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad \text{and} \quad v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}.$$



Important: $\text{Span}\{u_1, u_2\} = \text{Span}\{v_1, v_2\} = W$: this is an *orthogonal* basis for the *same* subspace.

The Gram–Schmidt Process

Three vectors

Find an orthogonal basis $\{u_1, u_2, u_3\}$ for $W = \text{Span}\{v_1, v_2, v_3\} = \mathbb{R}^3$, where


$$v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad v_3 = \begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix}.$$

Important: $\text{Span}\{u_1, u_2, u_3\} = \text{Span}\{v_1, v_2, v_3\} = W$: this is an *orthogonal* basis for the *same* subspace.

The Gram–Schmidt Process

Three vectors, continued

$$v_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \quad v_3 = \begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix} \xrightarrow{\text{G-S}} u_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad u_2 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad u_3 = \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$$



The Gram–Schmidt Process

Three vectors in \mathbb{R}^4

Find an orthogonal basis $\{u_1, u_2, u_3\}$ for $W = \text{Span}\{v_1, v_2, v_3\}$, where

$$v_1 = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \quad v_2 = \begin{pmatrix} -1 \\ 4 \\ 4 \\ -1 \end{pmatrix} \quad v_3 = \begin{pmatrix} 4 \\ -2 \\ -2 \\ 0 \end{pmatrix}.$$

Poll

Summary

- ▶ We like orthogonal bases because they let us compute orthogonal projections.
- ▶ The Gram–Schmidt process turns an arbitrary basis into an orthogonal basis.