

Math 1553 Supplement §6.1, 6.2

Supplemental Problems

1. Match the statements (i)-(v) with the corresponding statements (a)-(e). All matrices are 3×3 . There is a unique correspondence. Justify the correspondences in words.

(i) $Ax = \begin{pmatrix} 5 \\ 1 \\ 2 \end{pmatrix}$ has a unique solution.

(ii) The transformation $T(v) = Av$ fixes a nonzero vector.

(iii) A is obtained from B by subtracting the third row of B from the first row of B .

(iv) The columns of A and B are the same; except that the first, second and third columns of A are respectively the first, third, and second columns of B .

(v) The columns of A , when added, give the zero vector.

(a) 0 is an eigenvalue of A .

(b) A is invertible.

(c) $\det(A) = \det(B)$

(d) $\det(A) = -\det(B)$

(e) 1 is an eigenvalue of A .

Solution.

(i) matches with (b).

(ii) matches with (e).

(iii) matches with (c).

(iv) matches with (d).

(v) matches with (a).

2. Find a basis $\mathcal{B}$ for the (-1) -eigenspace of $Z = \begin{pmatrix} 2 & 3 & 1 \\ 3 & 2 & 4 \\ 0 & 0 & -1 \end{pmatrix}$

Solution.

For $\lambda = -1$, we find $\text{Nul}(Z - \lambda I)$.

$$(Z - \lambda I \mid 0) = (Z + I \mid 0) = \left(\begin{array}{ccc|c} 3 & 3 & 1 & 0 \\ 3 & 3 & 4 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right) \xrightarrow{\text{rref}} \left(\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right).$$

Therefore, $x = -y$, $y = y$, and $z = 0$, so

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -y \\ y \\ 0 \end{pmatrix} = y \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}.$$

A basis is $\mathcal{B} = \left\{ \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} \right\}$. We can check to ensure $\begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}$ is an eigenvector with corresponding eigenvalue -1 :

$$Z \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 & 3 & 1 \\ 3 & 2 & 4 \\ 0 & 0 & -1 \end{pmatrix} \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} -2+3 \\ -3+2 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} = - \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}.$$

3. Suppose A is an $n \times n$ matrix satisfying $A^2 = 0$. Find all eigenvalues of A . Justify your answer.

Solution.

If λ is an eigenvalue of A and $v \neq 0$ is a corresponding eigenvector, then

$$Av = \lambda v \implies A(Av) = A\lambda v \implies A^2v = \lambda(Av) \implies 0 = \lambda(\lambda v) \implies 0 = \lambda^2 v.$$

Since $v \neq 0$ this means $\lambda^2 = 0$, so $\lambda = 0$. This shows that 0 is the only possible eigenvalue of A .

On the other hand, $\det(A) = 0$ since $(\det(A))^2 = \det(A^2) = \det(0) = 0$, so 0 must be an eigenvalue of A . Therefore, the only eigenvalue of A is 0 .

4. Give an example of matrices A and B which satisfy the following:
 (I) A and B have the same eigenvalues, and the same algebraic multiplicities for each eigenvalue.
 (II) For some eigenvalue λ , the λ -eigenspace for A has a different dimension than the λ -eigenspace for B .

Justify your answer.

Solution.

Many examples possible. For example, $A = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$.

Both A and B have characteristic equation $\lambda^2 = 0$, so each has eigenvalue $\lambda = 0$ with algebraic multiplicity two. However, the 0 -eigenspace for A is $\mathbf{R}^2$ and thus has dimension 2, while the 0 -eigenspace for B has dimension 1 (the line $y = 0$ in $\mathbf{R}^2$).

5. Let $A = \begin{pmatrix} 5 & -2 & 3 \\ 0 & 1 & 0 \\ 6 & 7 & -2 \end{pmatrix}$. Find the eigenvalues of A .

Solution.

We find the characteristic polynomial $\det(A - \lambda I)$ any way we like. The computation below uses the cofactor expansion along the second row:

$$\begin{aligned}\det(A - \lambda I) &= \det \begin{pmatrix} 5 - \lambda & -2 & 3 \\ 0 & 1 - \lambda & 0 \\ 6 & 7 & -2 - \lambda \end{pmatrix} = (1 - \lambda) \det \begin{pmatrix} 5 - \lambda & 3 \\ 6 & -2 - \lambda \end{pmatrix} \\ &= (1 - \lambda) \cdot [(5 - \lambda)(-2 - \lambda) - 3 \cdot 6] = (1 - \lambda)(\lambda^2 - 3\lambda - 28) \\ &= -\lambda^3 + 4\lambda^2 + 25\lambda - 28 \quad \text{or} \quad (1 - \lambda)(\lambda - 7)(\lambda + 4)\end{aligned}$$

The characteristic equation is thus $(1 - \lambda)(\lambda - 7)(\lambda + 4) = 0$, so the eigenvalues are $\lambda = -4$, $\lambda = 1$, and $\lambda = 7$.

6. Using facts about determinants, justify the following fact: if A is an $n \times n$ matrix, then A and A^T have the same characteristic polynomial.

Solution.

We will use three facts which apply to all $n \times n$ matrices B , Y , Z :

(1) $\det(B) = \det(B^T)$.

(2) $(Y - Z)^T = Y^T - Z^T$

(3) If λ is any scalar then $(\lambda I)^T = \lambda I$ since the identity matrix is completely symmetric about its diagonal.

Using these three facts in order, we find

$$\det(A - \lambda I) = \det((A - \lambda I)^T) = \det(A^T - (\lambda I)^T) = \det(A^T - \lambda I).$$