

**MATH 218 SECTION 3
MIDTERM EXAMINATION 2**

Name		Duke ID Number	
-------------	--	-----------------------	--

Please **read all instructions** carefully before beginning.

- Do not open this test booklet until you are directed to do so.
- You have 75 minutes to complete this exam.
- If you finish early, go back and check your work.
- This exam is closed book.
- You may use a calculator to do arithmetic. No other technology is allowed.
- For full credit you must show your work so that your reasoning is clear.
- Do not spend too much time on any one problem. Read them all through first and attack them in an order that allows you to make the most progress.
- Good luck!

$$\begin{bmatrix} \cos 90^\circ & \sin 90^\circ \\ -\sin 90^\circ & \cos 90^\circ \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

[Hint: this is a joke.]

[This page intentionally left blank]

Problem 1.

[20 points]

Consider the matrix A and its reduced row echelon form:

$$A = \begin{pmatrix} 1 & 1 & 7 & 5 \\ -2 & -2 & 4 & 8 \end{pmatrix} \xrightarrow{\text{RREF}} \begin{pmatrix} 1 & 1 & 0 & -2 \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

a) Compute a basis for the null space of A :

$$\left\{ \begin{pmatrix} \\ \\ \\ \end{pmatrix} \right\}$$

b) Compute a basis for the row space of A :

$$\left\{ \begin{pmatrix} \\ \\ \\ \end{pmatrix} \right\}$$

c) Find the complete solution of $A\mathbf{x} = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$. (Note that a particular solution is $\mathbf{x} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$.)

d) The dimension of $C(A)$ is

e) Find a vector \mathbf{b} such that $A\mathbf{x} = \mathbf{b}$ has no solution, or explain why no such \mathbf{b} exists.

[Scratch work for problem 1]

Problem 2.

[20 points]

Consider the subspace $V = \left\{ \begin{array}{l} \text{all solutions of } x_1 + x_2 + 7x_3 + 5x_4 = 0 \\ -2x_1 - 2x_2 + 4x_3 + 8x_4 = 0 \end{array} \right\}$.

a) Find an orthonormal basis of V :

$$\left(\begin{array}{c} \\ \\ \\ \end{array} \right)$$

b) Find an orthonormal basis of V^\perp :

$$\left(\begin{array}{c} \\ \\ \\ \end{array} \right)$$

c) Compute the matrix P for projection onto V :

$$P = \left(\begin{array}{cc} & \\ & \end{array} \right)$$

d) Compute the projection \mathbf{p} of $\mathbf{v} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix}$ onto V^\perp :

$$\mathbf{p} = \left(\begin{array}{c} \\ \\ \\ \end{array} \right)$$

[Scratch work for problem 2]

Problem 3.

[10 points]

All of the following statements are false. Provide a counterexample to each. You need not justify your answers.

a) Any two vectors in \mathbf{R}^3 span a plane.

b) If $A\mathbf{x} = \mathbf{0}$ has only the trivial solution, then $A\mathbf{x} = \mathbf{b}$ has a solution for every \mathbf{b} in \mathbf{R}^m .

c) If every vector in a subspace V is orthogonal to every vector in another subspace W , then $V = W^\perp$.

d) Any 2×2 orthogonal matrix has the form

$$Q = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

[Scratch work for problem 3]

Problem 4.

[20 points]

- a) Give an example of a 3×2 matrix A and a vector \mathbf{b} in \mathbf{R}^3 such that $A\mathbf{x} = \mathbf{b}$ has more than one least-squares solution.
- b) Give an example of a 3×3 matrix A such that $C(A) = N(A)$, or explain why no such matrix exists.
- c) Write three different nonzero vectors $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ in \mathbf{R}^3 such that $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is linearly dependent, but \mathbf{v}_3 is *not* in $S\{\mathbf{v}_1, \mathbf{v}_2\}$. Clearly indicate which is \mathbf{v}_3 .
- d) Explain why every projection matrix P can be written as QQ^T for an orthogonal matrix Q .

[Scratch work for problem 4]


Problem 5.

[10 points]

Your roommate Karxon is currently taking Math 105L. Karxon scored 72% on the first exam, 81% on the second exam, and 84% on the third exam. Not having taken linear algebra yet, Karxon does not know what kind of score to expect on the final exam. Luckily, you can help out.

- a) The general equation of a line in \mathbf{R}^2 is $y = Cx + D$. Write down the system of linear equations in C and D that would be satisfied by a line passing through the points $(1, 72)$, $(2, 81)$, and $(3, 84)$, and then write down the corresponding matrix equation.

- b) Solve the corresponding least squares problem for C and D , and use this to *write down* and *draw* the the best fit line below. [Use a calculator]



$$y = \boxed{}x + \boxed{}$$

- c) What score does this line predict for the fourth (final) exam?


[Scratch work for problem 5]

Problem 6.

[10 points]


This problem concerns a certain 2×2 matrix A and a vector $\mathbf{b} \in \mathbf{R}^2$. You do not know what they are numerically.

- a) The solutions of $A\mathbf{x} = \mathbf{b}$ are drawn below. Draw $N(A)$ in the same diagram.



- b) The rank of A is .

- c) Suppose that \mathbf{b} is the vector in the picture. Draw the left null space of A in the same picture. [This is the same \mathbf{b} as before, so in particular, $A\mathbf{x} = \mathbf{b}$ has a solution.]



[Scratch work for problem 6]