

MATH 218D
PRACTICE FINAL EXAMINATION

Name		Duke Email	
-------------	--	-------------------	--

Please **read all instructions** carefully before beginning.

- You have 200 minutes to complete this exam and upload your work. The exam itself is meant to take 100 minutes to complete, so hopefully you will have enough time.
- For full credit you must **show your work** so that your reasoning is clear.
- If you need clarification or think you've found a typo, ask a **private question on Piazza**. We'll be monitoring it.
- If you have time, go back and check your work.
- You may use **your class notes** (not the ones from the website) and the **interactive row reducer** during this exam. You may use a **calculator** for doing arithmetic. All other materials and aids are strictly prohibited.
- You are not allowed to receive **outside help** during this exam. Consulting with someone else is considered cheating; suspected instances will result in immediate referral to the Office of Student Conduct.
- Be sure to **tag your answers** on Gradescope, and **use a scanning app**.
- Good luck!

Complete when starting the exam: I will neither give nor receive aid on this exam.

Signed: _____ Time: _____

Complete after finishing the exam: I have neither given nor received aid on this exam.

Signed: _____ Time: _____

This is a practice exam. It is meant to be similar in format, length, and difficulty to the real exam. It is **not** meant as a comprehensive list of study problems. I recommend completing the practice exam in 75 minutes, without distractions.

Problem 1.

[20 points]

Consider the sequence of numbers $0, 1, 5, 31, 185, \dots$ given by the recursive formula

$$a_0 = 0 \quad a_1 = 1 \quad a_n = 5a_{n-1} + 6a_{n-2}.$$

a) Find a matrix A such that

$$A \begin{pmatrix} a_{n-2} \\ a_{n-1} \end{pmatrix} = \begin{pmatrix} a_{n-1} \\ a_n \end{pmatrix}.$$

b) Find the eigenvalues of A , and find corresponding eigenvectors.

c) Give a non-recursive formula for a_n .

Problem 2.

[20 points]

A certain matrix A has singular value decomposition $A = U\Sigma V^T$, where

$$U = \begin{pmatrix} | & | & | & | \\ u_1 & u_2 & u_3 & u_4 \\ | & | & | & | \end{pmatrix} \quad \Sigma = \begin{pmatrix} 4 & 0 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad V = \begin{pmatrix} | & | & | & | & | \\ v_1 & v_2 & v_3 & v_4 & v_5 \\ | & | & | & | & | \end{pmatrix}.$$

- What is the rank of A ?
- What is the maximum value of $\|Ax\|$ subject to $\|x\| = 1$?
- Find orthonormal bases of the four fundamental subspaces of A .
- What is the singular value decomposition of A^T ?
- What is the pseudoinverse of A ?

Problem 3.

[15 points]

Consider the matrix

$$A = \begin{pmatrix} 1 & 2 & 5 & 0 \\ 1 & 2 & 4 & 2 \\ 0 & -1 & 0 & 8 \\ -1 & -3 & -1 & -1 \end{pmatrix}.$$

a) Find a permutation matrix P , a lower-unitriangular matrix L , and an upper-triangular matrix U such that $PA = LU$.

b) Use a) to solve $Ax = b$, for $b = \begin{pmatrix} 12 \\ 1 \\ -30 \\ 6 \end{pmatrix}$.

c) What is $\det(A)$?

Problem 4.

[20 points]

Consider the subspace

$$W = \text{Span} \left\{ \begin{pmatrix} 1 \\ -1 \\ -2 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 3 \\ -1 \\ -1 \end{pmatrix} \right\}.$$

- a) Compute an orthonormal basis for W . [Hint: W is *not* all of \mathbf{R}^3 .]
- b) What is $\dim(W)$?
- c) Compute the matrix P for orthogonal projection onto W . (You may write P as a product of two matrices, without expanding.)
- d) Write an eigenvector of P .
- e) Find the distance from $\begin{pmatrix} 1 \\ 5 \\ -2 \end{pmatrix}$ to W .
- f) Compute a basis for W^\perp .

Problem 5.

[20 points]

Consider the data points

$$\begin{pmatrix} 3 \\ 2 \\ -2 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \\ -2 \end{pmatrix}, \begin{pmatrix} 2 \\ 4 \\ -4 \end{pmatrix}, \begin{pmatrix} 2 \\ 0 \\ -4 \end{pmatrix}.$$

- a) Form the matrix A_0 with the data points as columns, and form the matrix A by subtracting the row averages from A_0 .
- b) Find the eigenvalues and eigenvectors of $S = \frac{1}{3}AA^T$.
- c) Find the line closest to the columns of A .
- d) Find the plane closest to the columns of A .
- e) Find the plane closest to the original data points.

Problem 6.

[30 points]

Give examples of matrices with each of the following properties. If no such matrix exists, explain why. *All matrices in this problem have real entries.*

- a) A 4×4 matrix A such that $\text{Col}(A) = \text{Nul}(A)$.
- b) A 4×6 matrix of rank 6.
- c) A 2×2 matrix whose column space is the line $3x + y = 0$ and with null space $\{0\}$.
- d) A 2×2 matrix A that is not diagonalizable over \mathbf{C} , such that A^2 is diagonalizable.
- e) A 3×4 matrix with singular values 2 and 1.
- f) A positive-semidefinite symmetric matrix that is not positive-definite.
- g) A matrix of rank 1 that cannot be written as a product of a column vector and a row vector.
- h) A nonzero symmetric matrix with characteristic polynomial $p(\lambda) = \lambda^2$.
- i) A matrix A satisfying
$$\dim(\text{Row}(A)^\perp) = 2 \quad \text{and} \quad \dim(\text{Col}(A)^\perp) = 3.$$
- j) A 3×3 matrix with no real eigenvalues.

Problem 7.

[10 points]

Let A be an $m \times n$ matrix. Which of the following are equivalent to the statement “the columns of A are linearly independent?” Circle all that apply.

- (1) A has full column rank.
- (2) $Ax = b$ has a unique solution for every b in \mathbf{R}^m .
- (3) $Ax = b$ has a unique least-squares solution for every b in \mathbf{R}^m .
- (4) $Ax = 0$ has a unique solution.
- (5) A has n pivots.
- (6) $\text{Nul}(A) = \{0\}$.
- (7) $m \geq n$.
- (8) $A^T A$ is invertible.
- (9) AA^T is invertible.
- (10) $A^+ A$ is the identity matrix.
- (11) $\text{Row}(A) = \mathbf{R}^n$.



Problem 8.

[10 points]

A certain 2×2 matrix A has the singular value decomposition

$$A = \begin{pmatrix} | & | \\ u_1 & u_2 \\ | & | \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} | & | \\ v_1 & v_2 \\ | & | \end{pmatrix}^T,$$

where u_1, u_2, v_1, v_2 are drawn in the diagrams below. Given x and y in the diagram on the left, draw Ax and Ay on the diagram on the right.



Problem 9.

[15 points]

A certain diagonalizable 2×2 matrix A is equal to CDC^{-1} , where C has columns w_1, w_2 pictured below, and $D = \begin{pmatrix} 2 & 0 \\ 0 & 1/4 \end{pmatrix}$.

- Draw $C^{-1}v$ on the left.
- Draw $DC^{-1}v$ on the left.
- Draw $Av = CDC^{-1}v$ on the right.

