

MATH 218D
PRACTICE MIDTERM EXAMINATION 1

Name		Duke Email	
-------------	--	-------------------	--

Please **read all instructions** carefully before beginning.

- You have 180 minutes to complete this exam and upload your work. The exam itself is meant to take 75 minutes to complete, so hopefully you will have enough time.
- If you have time, go back and check your work.
- You may use **your class notes** (not the ones from the website) and the **interactive row reducer** during this exam. All other materials are strictly prohibited.
- You are not allowed to receive **outside help** during this exam. Consulting with someone else is considered cheating; suspected instances will result in immediate referral to the Office of Student Conduct.
- For full credit you must show your work so that your reasoning is clear.
- Be sure to **tag your answers** on Gradescope, and **use a scanning app**.
- Good luck!

Complete when starting the exam: I will neither give nor receive aid on this exam.

Signed: _____ Time: _____

Complete after finishing the exam: I have neither given nor received aid on this exam.

Signed: _____ Time: _____

This is a practice exam. It is meant to be similar in format, length, and difficulty to the real exam. It is **not** meant as a comprehensive list of study problems. I recommend completing the practice exam in 75 minutes, without distractions.

Problem 1.

[25 points]

Consider

$$A = \begin{pmatrix} 1 & 2 & 5 & 0 \\ 1 & 2 & 4 & 2 \\ 0 & -1 & 0 & 8 \\ -1 & -3 & -1 & -1 \end{pmatrix} \quad b = \begin{pmatrix} 12 \\ 1 \\ -30 \\ 6 \end{pmatrix}.$$

The goal of this question is to solve the equation $Ax = b$ like a computer would.

- a) Carry out Gaussian reduction with maximal partial pivoting to find a $PA = LU$ decomposition. You should obtain

$$U = \begin{pmatrix} 1 & 2 & 5 & 0 \\ 0 & -1 & 0 & 8 \\ 0 & 0 & 4 & -9 \\ 0 & 0 & 0 & -\frac{1}{4} \end{pmatrix}.$$

Be sure to specify what L and P are.

- b) Solve the equations $Ly = Pb$ and $Ux = y$ to find a solution of $Ax = b$.

Problem 2.

[5 points]

Consider the matrix

$$B = \begin{pmatrix} 1 & -1 & 0 \\ -1 & 7 & 1 \\ 2 & 4 & 1 \end{pmatrix}.$$

Is B invertible? If so, find its inverse. If not, explain why.

Problem 3.

[20 points]

Consider the matrix

$$D = \begin{pmatrix} 1 & 2 & 3 & 2 & 14 & 9 \\ 0 & 0 & 0 & 2 & 10 & 6 \\ 0 & 0 & 0 & 0 & 0 & 3 \end{pmatrix}.$$

Note that this matrix is in row echelon form.

a) Fill in the blanks:

(1) The column space $\text{Col}(D)$ is a subspace of \mathbf{R}^m , where $m = \square$.

(2) The null space $\text{Nul}(D)$ is a subspace of \mathbf{R}^n , where $n = \square$.

b) Write down a vector b such that $Dx = b$ has no solution. If no such vector exists, explain why not.

c) Compute the reduced row echelon form of D .

d) Find a set of vectors that spans $\text{Nul}(D)$.

e) Compute the solutions of $Dx = (2, 2, 0)$, noting that $D(0, 0, 0, 1, 0, 0) = (2, 2, 0)$.

Problem 4.

[5 points]

Decide if a matrix with the following properties has full row rank, full column rank, both, or neither.

a) $Ax = b$ has 0 or 1 solutions, depending on b .

b) $Ax = b$ has 1 solution for every b .

c) $Ax = b$ has 0 or ∞ solutions, depending on b .

d) $Ax = b$ has ∞ solutions, for every b .

Problem 5.

[10 points]

Consider the subspace V and vectors b_1 and b_2 :

$$V = \text{Span} \left\{ \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ -1 \\ -1 \end{pmatrix}, \begin{pmatrix} 3 \\ -5 \\ -4 \end{pmatrix} \right\} \quad b_1 = \begin{pmatrix} -2 \\ 8 \\ 6 \end{pmatrix} \quad b_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

a) Is b_1 contained in V ? If so, write b_1 as a linear combination of the vectors in the span; if not, explain why.

b) Same question for b_2 .

c) Circle one: V is a

point line plane space

Why?


Problem 6.

[10 points]


Give examples of 2×2 matrices A, B, C with ranks 0, 1, and 2, respectively, and draw pictures of the null space and column space. (Be precise!)

a) Rank 0: $A = \begin{pmatrix} & \\ & \end{pmatrix}$

Nul(A)




Col(A)




b) Rank 1: $B = \begin{pmatrix} & \\ & \end{pmatrix}$

Nul(B)




Col(B)



c) Rank 2: $C = \begin{pmatrix} & \\ & \end{pmatrix}$

Nul(C)



Col(C)

