

MATH 218D
MIDTERM EXAMINATION 2

Name		Duke Email	
-------------	--	-------------------	--

Please **read all instructions** carefully before beginning.

- You have 180 minutes to complete this exam and upload your work. The exam itself is meant to take 75 minutes to complete, so hopefully you will have enough time.
- For full credit you must **show your work** so that your reasoning is clear. (You do not need to show your steps in Gauss–Jordan elimination.)
- If you need clarification or think you’ve found a typo, ask a **private question on Piazza**. We’ll be monitoring it.
- If you have time, go back and check your work.
- You may use **your class notes** (not the ones from the website) and the **interactive row reducer** during this exam. You may use a **calculator** for doing arithmetic. All other materials and aids are strictly prohibited.
- You are not allowed to receive **outside help** during this exam. Consulting with someone else is considered cheating; suspected instances will result in immediate referral to the Office of Student Conduct.
- Be sure to **tag your answers** on Gradescope, and **use a scanning app**.
- Good luck!

Complete when starting the exam: I will neither give nor receive aid on this exam.

Signed: _____ Time: _____

Complete after finishing the exam: I have neither given nor received aid on this exam.

Signed: _____ Time: _____

Problem 1.

[20 points]

Find a basis of the orthogonal complement of each of the following subspaces.

a) $\text{Nul} \begin{pmatrix} 1 & 0 & 2 & 3 \\ 2 & 1 & 4 & 1 \end{pmatrix}$

b) $\text{Col} \begin{pmatrix} 1 & 2 & -4 \\ 0 & -1 & 3 \\ 3 & 0 & 6 \\ 4 & -1 & 11 \end{pmatrix}$

c) The subspace of all vectors in \mathbf{R}^4 whose entries sum to zero.

d) The line $\{(t, 2t, 3t) : t \in \mathbf{R}\}$.

e) \mathbf{R}^3

Problem 2.

[25 points]

In this problem we will consider the best-fit plane $z = Bx + Cy + D$ through the data points

$$\begin{pmatrix} 3 \\ -5 \\ b_1 \end{pmatrix} \quad \begin{pmatrix} 1 \\ 1 \\ b_2 \end{pmatrix} \quad \begin{pmatrix} -1 \\ 5 \\ b_3 \end{pmatrix} \quad \begin{pmatrix} 3 \\ -7 \\ b_4 \end{pmatrix}.$$

- a) Compute the matrix A such that the coefficient vector $\hat{x} = (B, C, D)$ is the least-squares solution of

$$A \begin{pmatrix} B \\ C \\ D \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix}.$$

- b) Compute the QR decomposition of A .
- c) Given a vector $b = (b_1, b_2, b_3, b_4)$, explain how a computer would quickly compute \hat{x} using the QR decomposition you found in **b**.
- d) Compute the best-fit plane $z = Bx + Cy + D$ when $(b_1, b_2, b_3, b_4) = (10, -10, 20, -20)$, using your QR decomposition or otherwise.
- e) What is the minimum value of

$$(z(3, -5) - 10)^2 + (z(1, 1) + 10)^2 + (z(-1, 5) - 20)^2 + (z(3, -7) + 20)^2$$

for all planes $z = Bx + Cy + D$?

Problem 3.

[15 points]

Consider the matrix

$$A = \begin{pmatrix} 1 & 0 & 2 & k \\ -2 & 0 & 1 & 1 \\ k & 2 & 1 & 3 \\ 2 & 1 & -1 & 1 \end{pmatrix}.$$

- a) Compute the determinant of A . For what value(s) of k (a real number), if any, is A *not* invertible?
- b) Compute $\det(A^T A)$.
- c) Compute $\det(BAB^{-1})$ where B is a 4×4 invertible matrix.

Problem 4.

[20 points]

Consider the subspace V in \mathbf{R}^4 defined by $x_1 + 2x_2 - 2x_3 - x_4 = 0$.

- a) Compute the orthogonal projection b_{V^\perp} of the vector $b = (0, -3, 3, -2)$ onto V^\perp .
- b) Compute the orthogonal decomposition $b = b_V + b_{V^\perp}$.
- c) Find the matrix P_V for orthogonal projection onto V .
- d) Find a basis for $\text{Nul}(P_V)$.

Problem 5.

[16 points]

- a) Let A be an $m \times n$ matrix of rank r . Which of the following statements are equivalent to “ A has full row rank”?
- (1) $\text{Nul}(A^T) = \{0\}$
 - (2) $n = r$
 - (3) $\text{Col}(A) = \mathbf{R}^m$
 - (4) A has linearly independent columns
 - (5) A has a pivot in every row
 - (6) A is invertible
 - (7) $Ax = b$ is consistent for every vector b
- b) Explain why the projection matrix P_V onto a subspace V can be written as QQ^T for some matrix Q with orthonormal columns.
- c) Find three nonzero vectors $v_1, v_2, v_3 \in \mathbf{R}^3$ such that $\{v_1, v_2, v_3\}$ is linearly *dependent*, but v_3 is not in $\text{Span}\{v_1, v_2\}$. Be sure to label which is v_3 .
- d) Give an example of a 4×4 matrix A such that $\text{Nul}(A) = \text{Row}(A)$, or explain why no such matrix exists.

Problem 6.

[10 points]

Unit vectors u_1 and u_2 and a vector x are drawn in the picture below. Copy this picture onto your page as best you can, then:

- Draw and label $(x \cdot u_1)u_1$ and $(x \cdot u_2)u_2$.
- Draw and label $(x \cdot u_1)u_1 + (x \cdot u_2)u_2$.
- Note that $\{u_1, u_2\}$ is a basis for $V = \mathbf{R}^2$. Explain why $x = x_V \neq (x \cdot u_1)u_1 + (x \cdot u_2)u_2$ does not contradict the projection formula.

