

Math 218D Problem Session

Week 12

1. Rules of vector SVD

- a) $A = 1(1, 0)(1, 0)^T + 3(0, 1)(0, 1)^T$ is not an SVD since $1 < 3$, but singular values must be in decreasing order.
- b) $A = 4(1, 0)(0, 1)^T + 3(0, 1)(1, 0)^T$ is an SVD.
- c) $A = 3(1, -1)(1, 0)^T + 2(1, 1)(0, 1)^T$ is not an SVD, since $(1, -1)$ and $(1, 1)$ are not unit vectors.
- d) $A = -3(1/\sqrt{2}, -1/\sqrt{2}, 0)(1, 0)^T + 2(0, 0, 1)(0, 1)^T$ is not an SVD since $-3 < 0$, but singular values must be positive.
- e) $A = 3(-1/\sqrt{2}, 1/\sqrt{2}, 0)(1, 0)^T + 2(0, 0, 1)(0, 1)^T$ is an SVD.
- f) $A = 5(1, 0, 0)(0, 1)^T + 3(0, 1, 0)(1, 0)^T + 2(0, 0, 1)(0, 1)^T$ is not an SVD, since the vectors $(0, 1)$, $(1, 0)$, $(0, 1)$ are not orthogonal.

- 2. The matrix SVD** Suppose that A is an $m \times n$ matrix of rank r , with SVD $A = U\Sigma V^T$.
- U is an $m \times m$ matrix, Σ is a $m \times n$ matrix, and V is a $n \times n$ matrix. The matrices U and V are orthogonal matrices. The first r diagonal entries of Σ are > 0 .
 - $A^T A = (U\Sigma V^T)^T U\Sigma V^T = V\Sigma^T U^T U\Sigma V^T = V(\Sigma^T \Sigma)V^T$. Therefore $Q_1 = V$ and $D_1 = \Sigma^T \Sigma$. The columns of V are eigenvectors of $A^T A$, and the eigenvalues are the diagonal entries of the $n \times n$ matrix $\Sigma^T \Sigma$, which are $\sigma_1^2, \dots, \sigma_r^2, 0, \dots, 0$.
 - $AA^T = U\Sigma V^T (U\Sigma V^T)^T = U\Sigma V^T V\Sigma^T U^T = U(\Sigma \Sigma^T)U^T$. Therefore $Q_2 = U$ and $D_2 = \Sigma \Sigma^T$. The columns of U are eigenvectors of AA^T , and the eigenvalues are the diagonal entries of the $m \times m$ matrix $\Sigma \Sigma^T$, which are $\sigma_1^2, \dots, \sigma_r^2, 0, \dots, 0$.
 - $V^T v_i = (v_1 \cdot v_i, \dots, v_i \cdot v_i, \dots, v_n \cdot v_i) = (0, \dots, 1, \dots, 0)$, $\Sigma V^T v_i = \Sigma(0, \dots, 1, \dots, 0) = (0, \dots, \sigma_i, \dots, 0)$, $Av_i = U\Sigma V^T v_i = U(0, \dots, \sigma_i, \dots, 0) = \sigma_i Ue_i = \sigma_i u_i$.

3. Computing the vector SVD

- a) $A = \begin{pmatrix} 0 & -1 \\ 3 & 0 \end{pmatrix}$. The matrix $A^T A = \begin{pmatrix} 9 & 0 \\ 0 & 1 \end{pmatrix}$ has eigenvalues $\lambda_1 = 9, \lambda_2 = 1$, with eigenvectors $v_1 = (1, 0)$ and $v_2 = (0, 1)$. The singular values are $\sigma_1 = \sqrt{\lambda_1} = 3$ and $\sigma_2 = \sqrt{\lambda_2} = 1$. The left singular vectors are $u_1 = \frac{1}{3}Av_1 = \frac{1}{3}(0, 3) = (0, 1)$ and $u_2 = \frac{Av_2}{1} = (-1, 0)$. The vector SVD is

$$A = 3(0, 1)(1, 0)^T + 1(-1, 0)(0, 1)^T.$$

- b) $A = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & -3 & 0 & 0 \end{pmatrix}$. The matrix $A^T A = \begin{pmatrix} 4 & 0 & 0 & 0 \\ 0 & 9 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ has eigenvalues $\lambda_1 =$

$9, \lambda_2 = 4, \lambda_3 = 1, \lambda_4 = 0$. The orthonormal eigenvectors for the non-zero eigenvalues are $v_1 = (0, 1, 0, 0)$, $v_2 = (1, 0, 0, 0)$, and $v_3 = (0, 0, 0, 1)$. The singular values are $\sigma_1 = 3, \sigma_2 = 2, \sigma_3 = 1$. The left singular vectors are $u_1 = \frac{1}{3}Av_1 = (0, 0, -1)$, $u_2 = \frac{1}{2}Av_2 = (1, 0, 0)$, $u_3 = \frac{1}{1}Av_3 = (0, 1, 0)$. The vector SVD is

$$A = 3(0, 0, -1)(0, 1, 0, 0)^T + 2(1, 0, 0)(1, 0, 0, 0)^T + 1(0, 1, 0)(0, 0, 0, 1)^T.$$

- c) $A = \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}$. The matrix $A^T A = \begin{pmatrix} 4 & 2 \\ 2 & 5 \end{pmatrix}$ has characteristic polynomial $\lambda^2 - 9\lambda + 16$, with eigenvalues $\lambda_1 = \frac{9+\sqrt{17}}{2}, \lambda_2 = \frac{9-\sqrt{17}}{2}$. We then have eigenvectors $v_1 = \frac{(-2, 4-\lambda_1)}{\|(-2, 4-\lambda_1)\|} \approx (-0.615, -0.788)$, and $v_2 = \frac{(-2, 4-\lambda_2)}{\|(-2, 4-\lambda_2)\|} \approx (-0.788, 0.615)$.

The singular values are $\sigma_1 = \sqrt{\lambda_1} \approx 2.562$ and $\sigma_2 = \sqrt{\lambda_2} \approx 1.562$.

The left singular vectors are $u_1 = \frac{Av_1}{\sigma_1} \approx (-0.788, -0.615)$ and $u_2 = \frac{Av_2}{\sigma_2} \approx (-0.615, 0.788)$.

The vector SVD is, approximately,

$$A = 2.562(-0.788, -0.615)(-0.615, -0.788)^T + 1.562(-0.615, 0.788)(-0.788, 0.615)^T.$$

(The fact that the u and v vectors look so similar seems to be a coincidence.)

4. Computing the matrix SVD

Compute the matrix SVD of each of the following matrices:

- a) $A = \begin{pmatrix} 0 & -1 \\ 3 & 0 \end{pmatrix}$. Since $m = n = r = 2$, we can just use the singular vectors and values found in problem 3a) (no need to find ONB for $\text{Nul}(A)$ or $\text{Nul}(A^T)$.) We have

$$A = U\Sigma V^T = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}^T.$$

- b) $A = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & -3 & 0 & 0 \end{pmatrix}$. We found the vector SVD of this in 3b). Since $m = 3, n = 4, r = 3$, we need to find the additional vector v_4 , an ONB of $\text{Nul}(A)$. Since the matrix $A^T A$ was rather simple, and $\text{Nul}(A^T A) = \text{Nul}(A)$, we can use that $A^T A$ had unit eigenvector $v_4 = (0, 0, 1, 0)$ for the eigenvalue $\lambda_4 = 0$. Then

$$A = U\Sigma V^T = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 3 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}^T.$$

- c) $A = \begin{pmatrix} 1 & 2 \\ 1 & 2 \\ 2 & 4 \end{pmatrix}$. The matrix $A^T A = \begin{pmatrix} 6 & 12 \\ 12 & 24 \end{pmatrix}$ has characteristic polynomial $\lambda^2 - 30\lambda$, with eigenvalues $\lambda_1 = 30$ and $\lambda_2 = 0$. The λ_1 -eigenvector equals $v_1 = \frac{\begin{pmatrix} 12, 24 \end{pmatrix}}{\| \begin{pmatrix} 12, 24 \end{pmatrix} \|} = \frac{\begin{pmatrix} 1, 2 \end{pmatrix}}{\| \begin{pmatrix} 1, 2 \end{pmatrix} \|} = \frac{1}{\sqrt{5}}(1, 2)$, while the λ_2 -eigenvector equals $v_2 = \frac{\begin{pmatrix} 12, -6 \end{pmatrix}}{\| \begin{pmatrix} 12, -6 \end{pmatrix} \|} = \frac{1}{\sqrt{5}}(2, -1)$.

The only singular value is $\sigma_1 = \sqrt{\lambda_1} = \sqrt{30}$. We find the left singular vector $u_1 = \frac{1}{\sqrt{30}}Av_1 = \frac{1}{5\sqrt{6}}(5, 5, 10) = \frac{1}{\sqrt{6}}(1, 1, 2)$.

We already found the vector v_2 spanning $\text{Nul}(A^T A) = \text{Nul}(A)$. It remains to find an ONB u_2, u_3 of $\text{Nul}(A^T)$. It is not hard to see that $(1, -1, 0)$ and $(2, 0, -1)$ are a basis of $\text{Nul}(A^T)$, but they are not orthonormal.

Doing Gram-Schmidt, we first replace $(1, -1, 0)$ with $(1, -1, 0)$ and replace $(2, 0, -1)$ with $(2, 0, -1) - \frac{(2, 0, -1) \cdot (1, -1, 0)}{(1, -1, 0) \cdot (1, -1, 0)}(1, -1, 0) = (2, 0, -1) - (1, -1, 0) = (1, 1, -1)$.

(I avoided using the usual names for vectors in Gram-Schmidt, since it would be easy to confuse with the u and v vectors of SVD, which have a totally different meaning).

Making these unit vectors, we find that $u_2 = \frac{1}{\sqrt{2}}(1, -1, 0)$ and $u_3 = \frac{1}{\sqrt{3}}(1, 1, -1)$ form an ONB of $\text{Nul}(A^T)$.

We conclude that

$$U = \begin{pmatrix} 1/\sqrt{6} & 1/\sqrt{2} & 1/\sqrt{3} \\ 1/\sqrt{6} & -1/\sqrt{2} & 1/\sqrt{3} \\ 2/\sqrt{6} & 0 & -1/\sqrt{3} \end{pmatrix}, V = \begin{pmatrix} 1/\sqrt{5} & 2/\sqrt{5} \\ 2/\sqrt{5} & -1/\sqrt{5} \end{pmatrix}, \Sigma = \begin{pmatrix} \sqrt{30} & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

Warning: Many other answers are possible for U and V . Your columns of V might be off by a sign, your first column of U might be off by a sign, and the final two columns of U can look quite different.

5. Sums of rank 1 matrices

This final problem is not about SVDs, but just about sums of rank one matrices.

- a) Without computing A , we will explain why

$$A = (1, 2, 1)(1, 1)^T + (1, -1, 1)(-1, 1)^T$$

is a rank 2 matrix.

Since $A(1, 1) = (1, 2, 1)((1, 1) \cdot (1, 1)) + (1, -1, 1)((-1, 1) \cdot (1, 1)) = 2(1, 2, 1) + 0$, the vector $2(1, 2, 1)$ is in the column space of A . Similarly, $A(-1, 1) = (1, 2, 1)((-1, 1) \cdot (1, 1)) + (1, -1, 1)((-1, 1) \cdot (-1, 1)) = 0 + 2(1, -1, 1)$, so $2(1, -1, 1)$ is also in the column space of A since these two column space vectors are linearly independent, the rank of A is at least 2. Since A is a 3×2 matrix, its rank is at most 2. Therefore the rank of A equals 2.

- b) If $A = u_1 v_1^T + \cdots + u_r v_r^T$ for some vectors $u_i \in \mathbf{R}^m$ and $v_j \in \mathbf{R}^n$, we will explain why the rank of A is at most r .

For any vector x , $Ax = (v_1 \cdot x)u_1 + \cdots + (v_r \cdot x)u_r$. Therefore any vector b for which $Ax = b$ is consistent must be a linear combination of u_1, \dots, u_r . In other words, $\text{Col}(A) \subset \text{Span}\{u_1, \dots, u_r\}$. Since $\dim \text{Span}\{u_1, \dots, u_r\} \leq r$, and $\dim \text{Col}(A) \leq \dim \text{Span}\{u_1, \dots, u_r\}$, we conclude that $\text{rank}(A) = \dim \text{Col}(A) \leq r$.

- c) Suppose that the vectors $u_1, \dots, u_r \in \mathbf{R}^m$ are a linearly independent set of vectors, and the vectors $v_1, \dots, v_r \in \mathbf{R}^n$ are also linearly independent. We consider the matrix $A = u_1 v_1^T + \cdots + u_r v_r^T$.

Since the v_i vectors are linearly independent, $\text{Span}\{v_1, \dots, v_r\}$ is r -dimensional, while $\text{Span}\{v_2, \dots, v_r\}$ is $(r - 1)$ -dimensional. Using Gram-Schmidt, we can find a vector $v \in \text{Span}\{v_1, \dots, v_r\}$ which is orthogonal to $\text{Span}\{v_2, \dots, v_r\}$ but not orthogonal to v_1 (i.e. we can project v_1 onto the orthogonal complement of $\text{Span}\{v_2, \dots, v_r\}$).

Using this vector v , $Av = (v_1 \cdot v)u_1 + \cdots + (v_r \cdot v)u_r = (v_1 \cdot v)u_1$, since $v \cdot v_2 = 0$, $v \cdot v_3 = 0, \dots$. In other words, $A\left(\frac{v}{v_1 \cdot v}\right) = u_1$, which verifies that u_1 is in $\text{Col}(A)$.

A similar argument shows that each of the vectors u_i is in $\text{Col}(A)$. Therefore $\text{Span}\{u_1, \dots, u_r\} \subset \text{Col}(A)$. Since the u_i vectors are linearly independent, $r = \dim \text{Span}\{u_1, \dots, u_r\} \leq \dim \text{Col}(A) = \text{rank}(A)$. On the other hand, by **b**, $\text{rank}(A) \leq r$. Therefore $\text{rank}(A) = r$.