

MATH 218D-1
PRACTICE FINAL EXAMINATION

Name		Duke Email	
-------------	--	-------------------	--

Please **read all instructions** carefully before beginning.

- Do not open this test booklet until you are directed to do so.
- You have 180 minutes to complete this exam.
- If you finish early, go back and check your work.
- You may use a **calculator** for doing arithmetic, but you should not need one. You may use a 8.5×11 " **note sheet** as well. All other materials and aids are strictly prohibited.
- For full credit you must **show your work** so that your reasoning is clear, unless otherwise indicated.
- Do not spend too much time on any one problem. Read them all through first and attack them in an order that allows you to make the most progress.
- Good luck!

This is a practice exam. It is meant to be similar in format, length, and difficulty to the real exam. It is **not** meant as a comprehensive list of study problems. I recommend completing the practice exam in 180 minutes, without notes or distractions.

Problem 1.

[20 points]

Consider the matrix

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 1 \\ 0 & 3 & 1 \end{pmatrix}.$$

a) Compute the $A = LU$ decomposition of A .

b) Compute A^{-1} .

c) Express A^{-1} as a product of elementary matrices.

d) Solve $Ax = \begin{pmatrix} 3 \\ 0 \\ -1 \end{pmatrix}$.

Problem 2.

[20 points]

A certain 3×2 matrix A satisfies $AA^T = QDQ^T$ for

$$Q = \begin{pmatrix} 1/\sqrt{3} & 1/\sqrt{2} & 1/\sqrt{6} \\ -1/\sqrt{3} & 0 & 2/\sqrt{6} \\ 1/\sqrt{3} & -1/\sqrt{2} & 1/\sqrt{6} \end{pmatrix} \quad D = \begin{pmatrix} 4 & 0 & 0 \\ 0 & 1/9 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

- What is the rank of A ?
- What are the singular values of A ?
- What are the maximum and minimum values of $\|Ax\|$ subject to $\|x\| = 1$?
- Find orthonormal bases for $\text{Col}(A)$ and $\text{Nul}(A^T)$.

Suppose now that the columns of A are *orthogonal*.

- What are the two possibilities for $A^T A$?
- You have enough information to determine the columns of A up to sign: the longer column is $\pm(?)$, and the shorter column is $\pm(?)$.


Problem 3.

[20 points]

Consider the three data points

$$\begin{pmatrix} 3 \\ 1 \end{pmatrix}, \begin{pmatrix} -4 \\ -4 \end{pmatrix}, \begin{pmatrix} 1 \\ 3 \end{pmatrix}.$$

- Form the matrix A with the data points as columns, and note that the row averages are equal to zero.
- Compute the covariance matrix $S = \frac{1}{3-1}AA^T$. What are its eigenvalues $\lambda_1 > \lambda_2$ and corresponding unit eigenvectors v_1, v_2 ?
- Sketch the line in the direction of largest variance in the grid below.
- What is the variance of the data in the direction perpendicular to the line you drew in c)?



The **data points** in the problem.

Problem 4.


[20 points]

Consider the three data points

$$\begin{pmatrix} 3 \\ 1 \end{pmatrix}, \begin{pmatrix} -4 \\ -4 \end{pmatrix}, \begin{pmatrix} 1 \\ 3 \end{pmatrix}$$

of the previous problem. In this problem, we will approximate these data points with a best-fit line in the least-squares sense.

- The general equation of a line in the plane is $y = Bx + C$. Find a system of linear equations that would be satisfied by a line passing through the above three points, and write this as a matrix equation $Ax = b$.
- Find the least-squares solution \hat{x} of $Ax = b$. What is the equation of the best-fit line?
- Sketch the best-fit line in the grid below.
- The line you drew in this problem is different than the line you drew in the previous problem. How can that be?



The **data points** in the problem.

Problem 5.

[15 points]

Consider the matrix

$$A = \begin{pmatrix} 4 & 2 & -4 \\ 0 & 2 & 0 \\ 2 & 2 & -2 \end{pmatrix}.$$

a) Find an invertible matrix C and a diagonal matrix D such that $A = CDC^{-1}$.

b) Compute $A^{100} \begin{pmatrix} 3 \\ 0 \\ 2 \end{pmatrix}$.

Problem 6.

[25 points]

Consider the subspace

$$W = \text{Span} \left\{ \begin{pmatrix} 1 \\ -1 \\ 0 \\ -2 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 4 \\ 3 \end{pmatrix}, \begin{pmatrix} 3 \\ -1 \\ 4 \\ -1 \end{pmatrix}, \begin{pmatrix} 4 \\ -8 \\ -2 \\ 0 \end{pmatrix} \right\}.$$

- a) Find an orthonormal basis of W .

Now consider the subspace V with orthonormal basis

$$\left\{ \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}, \frac{1}{\sqrt{10}} \begin{pmatrix} 1 \\ -2 \\ -1 \\ 2 \end{pmatrix}, \frac{1}{\sqrt{10}} \begin{pmatrix} 2 \\ 1 \\ -2 \\ -1 \end{pmatrix} \right\}$$

(The reason for changing subspaces is to prevent carry-through error in the rest of the problem.)

- b) Compute the matrix P_V for orthogonal projection onto V . (You can write P_V as a product of two matrices without multiplying it out.)
- c) Find the orthogonal decomposition $x = x_V + x_{V^\perp}$ for $x = (1, 0, 1, 0)$. Your answer should involve fractions and not decimals.
- d) Find a basis for V^\perp .
[Hint: you already did this in c).]
- e) Find an implicit equation for V : that is, $V = \{(x_1, x_2, x_3, x_4) : (?) = 0\}$.
- f) Orthogonally diagonalize P_V : that is, find an orthogonal matrix Q and a diagonal matrix D such that $P_V = QDQ^T$.
[Hint: You have already done all of the necessary calculations.]

Problem 7.

[20 points]

True/false questions: no justification is needed.

- a) **T** **F** If $Ax = b$ has at least one solution for every $b \in \mathbf{R}^m$, then A has full row rank.
- b) **T** **F** Every elementary matrix is invertible.
- c) **T** **F** If x is in a subspace V , then the projection of x onto V is the zero vector.
- d) **T** **F** A triangular matrix A with real entries can have a complex (non-real) eigenvalue.
- e) **T** **F** A diagonalizable $n \times n$ matrix admits n linearly independent eigenvectors.
- f) **T** **F** The maximum value of $\|Ax\|$ subject to $\|x\| = 1$, is the largest eigenvalue of A .
- g) **T** **F** If A is an $m \times n$ matrix with linearly dependent columns, then the columns of A do not span \mathbf{R}^m .
- h) **T** **F** If A and B are $n \times n$ matrices and $\det(A) = 0$, then the columns of AB are linearly dependent.
- i) **T** **F** If A has linearly independent columns, then A^+A is the identity matrix.
- j) **T** **F** If λ is a eigenvalue of $A^T A$ and $\lambda > 0$, then λ is also an eigenvalue of AA^T .



Problem 8.

[10 points]

A certain 2×2 matrix A has the singular value decomposition

$$A = 3u_1v_1^T + 2u_2v_2^T$$

where u_1, u_2, v_1, v_2 are drawn in the diagrams below. The unit circle $\{x: \|x\| = 1\}$ is drawn on the left. Draw $\{Ax: \|x\| = 1\}$ on the right.



Problem 9.

[15 points]

The solution set of $Ax = b$ for a certain 3×2 matrix A is drawn below.

- Draw and label the solution sets of $Ax = \frac{1}{2}b$, $Ax = 0$, and $Ax = -b$ on the same diagram.
- What is the rank of A ?
- Explain why there exists some vector $b' \in \mathbf{R}^3$ such that $Ax = b'$ is inconsistent.

