

**MATH 218D-1**  
**PRACTICE MIDTERM EXAMINATION 2**

<b>Name</b>		<b>Duke Email</b>	
-------------	--	-------------------	--

Please **read all instructions** carefully before beginning.

- Do not open this test booklet until you are directed to do so.
- You have 75 minutes to complete this exam.
- If you finish early, go back and check your work.
- You may use a **calculator** for doing arithmetic, but you should not need one. All other materials and aids are strictly prohibited.
- For full credit you must **show your work** so that your reasoning is clear, unless otherwise indicated.
- Do not spend too much time on any one problem. Read them all through first and attack them in an order that allows you to make the most progress.
- Good luck!

This is a practice exam. It is meant to be similar in format, length, and difficulty to the real exam. It is **not** meant as a comprehensive list of study problems. I recommend completing the practice exam in 75 minutes, without notes or distractions.

## Problem 1.

[15 points]

Consider the vectors

$$v_1 = \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} \quad v_3 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}.$$

a) Is  $\{v_1, v_2, v_3\}$  linearly independent? If not, find a linear relation.

b) Compute the dimension of  $\text{Span}\{v_1, v_2, v_3\}$ .

c) Write  $(2, 6, -2)$  as a linear combination of  $v_1, v_2, v_3$ .

## Problem 2.

[20 points]

Find a basis of the orthogonal complement of each of the following subspaces.

a)  $\text{Nul} \begin{pmatrix} 1 & 0 & 2 & 3 \\ 2 & 1 & 4 & 1 \end{pmatrix}$

b)  $\text{Col} \begin{pmatrix} 1 & 2 & -4 \\ 0 & -1 & 3 \\ 3 & 0 & 6 \\ 4 & -1 & 11 \end{pmatrix}$

c) The subspace of all vectors in  $\mathbf{R}^4$  whose entries sum to zero.

d) The line  $\{(t, 2t, 3t) : t \in \mathbf{R}\}$ .

e)  $\mathbf{R}^3$

### Problem 3.

[25 points]

In this problem we will consider the best-fit plane  $z(x, y) = Bx + Cy + D$  through the data points

$$\begin{pmatrix} 3 \\ -5 \\ b_1 \end{pmatrix} \quad \begin{pmatrix} 1 \\ 1 \\ b_2 \end{pmatrix} \quad \begin{pmatrix} -1 \\ 5 \\ b_3 \end{pmatrix} \quad \begin{pmatrix} 3 \\ -7 \\ b_4 \end{pmatrix}.$$

- a) Find the matrix  $A$  such that the coefficient vector  $\hat{x} = (B, C, D)$  is the least-squares solution of

$$A \begin{pmatrix} B \\ C \\ D \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix}.$$

- b) Compute the QR decomposition of  $A$ .

c) Given a vector  $b = (b_1, b_2, b_3, b_4)$ , explain how a computer would quickly compute  $\hat{x}$  using the  $QR$  decomposition you found in **b**).

d) Compute the best-fit plane  $z = Bx + Cy + D$  when  $(b_1, b_2, b_3, b_4) = (10, -10, 20, -20)$ , using your  $QR$  decomposition or otherwise.

e) What is the minimum value of

$$(z(3, -5) - 10)^2 + (z(1, 1) + 10)^2 + (z(-1, 5) - 20)^2 + (z(3, -7) + 20)^2$$

for all planes  $z = Bx + Cy + D$ ?

### Problem 4.

[20 points]

Consider the subspace  $V$  in  $\mathbf{R}^4$  defined by  $x_1 + 2x_2 - 2x_3 - x_4 = 0$ .

a) Compute the orthogonal projection  $b_{V^\perp}$  of the vector  $b = (0, -3, 3, -2)$  onto  $V^\perp$ .

b) Compute the orthogonal decomposition  $b = b_V + b_{V^\perp}$ .

c) Find the matrix  $P_V$  for orthogonal projection onto  $V$ .

d) Find a basis for  $\text{Nul}(P_V)$ .

## Problem 5.


[16 points]

- a) Let  $A$  be an  $m \times n$  matrix of rank  $r$ . Which of the following statements are equivalent to “ $A$  has full row rank”?
- (1)  $\text{Nul}(A^T) = \{0\}$
  - (2)  $n = r$
  - (3)  $\text{Col}(A) = \mathbf{R}^m$
  - (4)  $A$  has linearly independent columns
  - (5)  $A$  has a pivot in every row
  - (6)  $A$  is invertible
  - (7)  $Ax = b$  is consistent for every vector  $b$
- b) Explain why the projection matrix  $P_V$  onto a subspace  $V$  can be written as  $QQ^T$  for some matrix  $Q$  with orthonormal columns. (What is  $Q$  in terms of  $V$ ?)
- c) Find three nonzero vectors  $v_1, v_2, v_3 \in \mathbf{R}^3$  such that  $\{v_1, v_2, v_3\}$  is linearly dependent, but  $v_3$  is not in  $\text{Span}\{v_1, v_2\}$ . Be sure to label which is  $v_3$ .
- d) Give an example of a  $4 \times 4$  matrix  $A$  such that  $\text{Nul}(A) = \text{Row}(A)$ , or explain why no such matrix exists.

## Problem 6.

[10 points]

Unit vectors  $u_1$  and  $u_2$  and a vector  $b$  are drawn in the picture below. In the same picture, draw and label **a)**  $(b \cdot u_1)u_1$  and  $(b \cdot u_2)u_2$  and **b)**  $(b \cdot u_1)u_1 + (b \cdot u_2)u_2$ .



- c) Note that  $\{u_1, u_2\}$  is a basis for  $V = \mathbf{R}^2$ . Explain why  $b = b_V \neq (b \cdot u_1)u_1 + (b \cdot u_2)u_2$  does not contradict the projection formula.