

MATH 218D-1
PRACTICE MIDTERM EXAMINATION 2

Name		Duke NetID	
-------------	--	-------------------	--

Please **read all instructions** carefully before beginning.

- Do not open this test booklet until you are directed to do so.
- You have 75 minutes to complete this exam.
- If you finish early, go back and check your work.
- The graders will only see the work on the **printed pages** (front and back). You may use other scratch paper, but the graders will not see anything written there.
- You may use a **calculator** for doing arithmetic, but you should not need one. All other materials and aids are strictly prohibited.
- For full credit you must **show your work** so that your reasoning is clear, unless otherwise indicated.
- Do not spend too much time on any one problem. Read them all through first and attack them in an order that allows you to make the most progress.
- Good luck!

This is a practice exam. It is meant to be similar in format, length, and difficulty to the real exam. It is **not** meant as a comprehensive list of study problems. I recommend completing the practice exam in 75 minutes, without notes or distractions.

Problem 1.

[20 points]

Consider the plane

$$V = \{(x, y, z) : x - y + 2z = 0\}.$$

- Find a basis for V .
- Find an orthogonal basis for V .
- Use the projection formula and your answer to part **b)** to compute the orthogonal projection b_V of the vector $b = (1, 1, -3)$ onto V .
- Find a basis for V^\perp .
- Find an orthogonal basis of \mathbf{R}^3 containing the basis vectors you found in **b)**.

Solution.

- a) There are many answers. If you find the solutions of $x - y + 2z = 0$ in parametric vector form, you get

$$\left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

- b) Running Gram–Schmidt on the above vectors gives

$$\left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix} \right\}.$$

c)

$$b_V = \frac{\begin{pmatrix} 1 \\ 1 \\ -3 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}}{\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + \frac{\begin{pmatrix} 1 \\ 1 \\ -3 \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}}{\begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}} \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \\ -1 \end{pmatrix}.$$

- d) Since $V = \text{Nul}\begin{pmatrix} 1 & -1 & 2 \end{pmatrix}$, the orthogonal complement V^\perp is the row space of $\begin{pmatrix} 1 & -1 & 2 \end{pmatrix}$:

$$\left\{ \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} \right\}.$$

- e) We just add the vectors in **b)** and **d)**:

$$\left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} \right\}.$$

(You could also notice that $b - b_V = (-1, 1, -2)$ spans V^\perp .)

Problem 2.

[20 points]

Consider the matrix

$$A = \begin{pmatrix} 1 & 2 & 5 \\ -1 & 1 & -4 \\ -1 & 4 & -3 \\ 1 & -4 & 7 \\ 1 & 2 & 1 \end{pmatrix}.$$

a) Find the QR decomposition of A . You should get $R = \begin{pmatrix} \sqrt{5} & -\sqrt{5} & 4\sqrt{5} \\ 0 & 6 & -2 \\ 0 & 0 & 4 \end{pmatrix}$.

b) Solve $R\hat{x} = Q^T \begin{pmatrix} 2 \\ -2 \\ 4 \\ -3 \\ 3 \end{pmatrix}$ to find the least-squares solution of $Ax = \begin{pmatrix} 2 \\ -2 \\ 4 \\ -3 \\ 3 \end{pmatrix}$.

c) Compute the matrix P_V for projection onto $V = \text{Col}(A)$.

Solution.

a)
$$Q = \begin{pmatrix} 1/\sqrt{5} & 1/2 & 1/2 \\ -1/\sqrt{5} & 0 & 0 \\ -1/\sqrt{5} & 1/2 & 1/2 \\ 1/\sqrt{5} & -1/2 & 1/2 \\ 1/\sqrt{5} & 1/2 & -1/2 \end{pmatrix}$$

b)
$$\hat{x} = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$$

c)
$$P_V = QQ^T = \frac{1}{10} \begin{pmatrix} 7 & -2 & 3 & 2 & 2 \\ -2 & 2 & 2 & -2 & -2 \\ 3 & 2 & 7 & -2 & -2 \\ 2 & -2 & -2 & 7 & -3 \\ 2 & -2 & -2 & -3 & 7 \end{pmatrix}$$

Problem 3.

[15 points]

Consider the data points

$$b_1 = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad b_2 = \begin{pmatrix} 1 \\ 8 \end{pmatrix} \quad b_3 = \begin{pmatrix} 3 \\ 8 \end{pmatrix} \quad b_4 = \begin{pmatrix} 4 \\ 20 \end{pmatrix}$$

drawn below.



- a) Find the matrix A such that the least-squares solution $\hat{x} = (C, D)$ of

$$A \begin{pmatrix} C \\ D \end{pmatrix} = b = \begin{pmatrix} 0 \\ 8 \\ 8 \\ 20 \end{pmatrix}$$

gives the coefficients of the best-fit line $y = Cx + D$.

- b) Find the equation of the best-fit line by computing the least-squares solution of the above equation. Graph this line in the above grid.
- c) Compute the minimized vector b_{V^\perp} . What does b_{V^\perp} represent in the original best-fit problem? (Here $V = \text{Col}(A)$.)
- d) What is the best-fit line among all lines *passing through the origin*?

Solution.

a)
$$A = \begin{pmatrix} 0 & 1 \\ 1 & 1 \\ 3 & 1 \\ 4 & 1 \end{pmatrix}$$

b)
$$\hat{x} = \begin{pmatrix} 4 \\ 1 \end{pmatrix} \rightsquigarrow y = 4x + 1$$

c) The minimized vector is

$$b_{V^\perp} = b - A\hat{x} = \begin{pmatrix} 0 \\ 8 \\ 8 \\ 20 \end{pmatrix} - \begin{pmatrix} 1 \\ 5 \\ 13 \\ 17 \end{pmatrix} = \begin{pmatrix} -1 \\ 3 \\ -5 \\ 3 \end{pmatrix}.$$

This is the vector of vertical distances from the data points to the graph of the best-fit line, drawn in **red** in the picture.

d) Using $y = Cx$ means solving the least-squares problem

$$\begin{pmatrix} 0 \\ 1 \\ 3 \\ 4 \end{pmatrix} C = \begin{pmatrix} 0 \\ 8 \\ 8 \\ 20 \end{pmatrix} \rightsquigarrow C = \frac{56}{13}.$$


The best-fit line is $y = \frac{56}{13}x$.

Problem 4.

[12 points]

A line V and a vector b are drawn below. Draw *and label*:

- a) The orthogonal projection b_V .
- b) The projection onto the orthogonal complement b_{V^\perp} .
- c) The vector $b - 2b_{V^\perp}$.



Problem 5.

[20 points]

Find a basis of the orthogonal complement of each of the following subspaces.

a) $\text{Nul} \begin{pmatrix} 1 & 0 & 2 & 3 \\ 2 & 1 & 4 & 1 \end{pmatrix}$

b) $\text{Col} \begin{pmatrix} 1 & 2 & -4 \\ 0 & -1 & 3 \\ 3 & 0 & 6 \\ 4 & -1 & 11 \end{pmatrix}$

c) The subspace of all vectors in \mathbf{R}^4 whose entries sum to zero.

d) The line $\{(t, 2t, 3t) : t \in \mathbf{R}\}$.

e) \mathbf{R}^3

Solution.

These are the bases you would obtain if you did the problem the same way I did.

a) $\left\{ \begin{pmatrix} 1 \\ 0 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \\ 4 \\ 1 \end{pmatrix} \right\}$ b) $\left\{ \begin{pmatrix} -3 \\ -6 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -4 \\ -9 \\ 0 \\ 1 \end{pmatrix} \right\}$ c) $\left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \right\}$

d) $\left\{ \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -3 \\ 0 \\ 1 \end{pmatrix} \right\}$ e) $\{\}$

Problem 6.

[16 points]

- a) Let A be an $m \times n$ matrix of rank r . Which of the following statements are equivalent to “ A has full row rank”?
- (1) $\text{Nul}(A^T) = \{0\}$
 - (2) $n = r$
 - (3) $\text{Col}(A) = \mathbf{R}^m$
 - (4) A has linearly independent columns
 - (5) A has a pivot in every row
 - (6) A is invertible
 - (7) $Ax = b$ is consistent for every vector b
- b) Explain why the projection matrix P_V onto a subspace V can be written as QQ^T for some matrix Q with orthonormal columns. (What is Q in terms of V ?)
- c) Find three nonzero vectors $v_1, v_2, v_3 \in \mathbf{R}^3$ such that $\{v_1, v_2, v_3\}$ is linearly dependent, but v_3 is not in $\text{Span}\{v_1, v_2\}$. Be sure to label which is v_3 .
- d) Give an example of a 4×4 matrix A such that $\text{Nul}(A) = \text{Row}(A)$, or explain why no such matrix exists.

Solution.

- a) (1),(3),(5),(7)
- b) Choose an orthonormal basis for V , then let Q be the matrix with those columns.
- c) There are many answers. One is

$$v_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad v_2 = \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix} \quad v_3 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}.$$

- d) This is impossible since $\text{Nul}(A) = \text{Row}(A)^\perp$.