

**MATH 218D-1
MIDTERM EXAMINATION 3**

Name		Duke NetID	
-------------	--	-------------------	--

Please **read all instructions** carefully before beginning.

- Do not open this test booklet until you are directed to do so.
- You have 75 minutes to complete this exam.
- If you finish early, go back and check your work.
- The graders will only see the work on the **printed pages** (front and back). You may use other scratch paper, but the graders will not see anything written there.
- You may use a **calculator** for doing arithmetic, but you should not need one. All other materials and aids are strictly prohibited.
- For full credit you must **show your work** so that your reasoning is clear, unless otherwise indicated.
- Do not spend too much time on any one problem. Read them all through first and attack them in an order that allows you to make the most progress.
- Good luck!

**WHO IS THE
MOST AWESOME
PERSON TODAY?**



[This page intentionally left blank]

Problem 1.

[20 points]

Consider the difference equation

$$\begin{aligned}x_{k+1} &= 3x_k && + 8z_k \\y_{k+1} &= -5x_k + 2y_k + 3z_k \\z_{k+1} &= -x_k + y_k.\end{aligned}$$

a) Setting $v_k = (x_k, y_k, z_k)$, find a matrix A such that $v_{k+1} = Av_k$.

$$A = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix}$$

b) Compute the characteristic polynomial of the matrix you found in a). *Do not factor* $p(\lambda)$.

$$p(\lambda) = \boxed{}$$

[Scratch work for Problem 1]

Problem 1, continued.

Now we change matrices to avoid carry-through error. Consider the matrix

$$B = \begin{pmatrix} 3 & 8 & -28 \\ -5 & 0 & 13 \\ -1 & 1 & 0 \end{pmatrix}.$$

- c) Find an invertible matrix C and a diagonal matrix D such that $B = CDC^{-1}$. The eigenvalues of B are 3, 1, and -1 .

[**Hint:** Choose the ± 1 entry as your pivot when doing elimination!]

$$C = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix}$$
$$D = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix}$$

- d) Solve the difference equation

$$v_{k+1} = Bv_k \quad v_0 = \begin{pmatrix} 5 \\ 5 \\ 2 \end{pmatrix}.$$

$$v_k = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix}$$

[Scratch work for Problem 1]

Problem 2.

[15 points]

Consider the symmetric matrix

$$S = \begin{pmatrix} 3 & -2 & 4 \\ -2 & 6 & 2 \\ 4 & 2 & 3 \end{pmatrix}.$$

Its eigenvalues are 7 and -2 .

a) Find an orthogonal matrix Q and a diagonal matrix D such that $S = QDQ^T$.

[Hint: It helps to scale your eigenvectors to have integer entries before normalizing.]

$$Q = \begin{pmatrix} \\ \\ \end{pmatrix}$$
$$D = \begin{pmatrix} \\ \\ \end{pmatrix}$$

b) What are the algebraic and geometric multiplicities of each eigenvalue?

$$\lambda = 7: \quad \text{AM} = \boxed{} \quad \text{GM} = \boxed{}$$

$$\lambda = -2: \quad \text{AM} = \boxed{} \quad \text{GM} = \boxed{}$$

c) What is the *minimum* value of $q(x) = x^T S x$ subject to $\|x\| = 1$, and at which vectors is the minimum achieved?

$$\text{Minimum} = \boxed{} \quad \text{at } x = \pm \begin{pmatrix} \\ \\ \end{pmatrix}$$

d) Find a matrix A such that $S = A^T A$, or explain why no such matrix exists.

[Scratch work for Problem 2]

Problem 3.

[15 points]

Consider the initial value problem

$$u' = Au, \quad u(0) = \begin{pmatrix} 3 \\ 2 \end{pmatrix} \quad \text{for} \quad A = \begin{pmatrix} 2 & -3 \\ 1 & -1 \end{pmatrix}.$$

a) The matrix A has complex eigenvalues $\lambda = \boxed{}$ and $\bar{\lambda} = \boxed{}$.

b) Find eigenvectors w and \bar{w} for λ and $\bar{\lambda}$, respectively.

$$w = \begin{pmatrix} \\ \end{pmatrix} \quad \bar{w} = \begin{pmatrix} \\ \end{pmatrix}$$

c) Solve the initial value problem. Your answer should only contain real numbers.


$$u = \begin{pmatrix} \\ \end{pmatrix}$$

[Scratch work for Problem 3]

Problem 4.

[10 points]

Find the 2×2 matrix A whose eigenspaces are drawn below. The grid is square.



$$A = \begin{pmatrix} & \\ & \end{pmatrix}$$

[Scratch work for Problem 4]

Problem 5.

[20 points]

Give examples of matrices with each of the following properties. If no such matrix exists, explain why. (No justification is needed if an example does exist.)

All matrices in this problem have real entries.

- a) A matrix with characteristic polynomial $p(\lambda) = -(\lambda - 2)(\lambda - 3)^2$ whose 2-eigenspace is a plane.
- b) A 2×2 diagonalizable matrix with only one eigenvalue.
- c) A 3×3 symmetric matrix that is diagonalizable over the complex numbers but not over the real numbers.
- d) A 3×3 matrix with no real eigenvalues.
- e) A diagonalizable 2×2 matrix with characteristic polynomial $p(\lambda) = \lambda^2$.

[Scratch work for Problem 5]

Problem 6.

[20 points]

True/false problems: **circle** the correct answer. No justification is needed. *All matrices in this problem have real entries.*

- a) **T** **F** A matrix with characteristic polynomial $p(\lambda) = -\lambda^3 + 3\lambda^2 - 2\lambda - 2$ is invertible.
- b) **T** **F** If A is a square matrix and x is a nonzero vector in $\text{Nul}(A)$, then x is an eigenvector of A .
- c) **T** **F** The eigenvalues of a square matrix are the diagonal entries.
- d) **T** **F** If S is symmetric, then either S or $-S$ is positive-semidefinite.
- e) **T** **F** Every 2×2 matrix is diagonalizable over the complex numbers.

[Scratch work for Problem 6]