

Determinants & Cofactors

Last time: we defined determinants using row ops.

- (1) If $A \xrightarrow{R_i + cR_j} B$ then $\det(A) = \det(B)$.
- (2) If $A \xrightarrow{R_i \times c} B$ then $\det(A) = \frac{1}{c} \det(B)$.
- (3) If $A \xrightarrow{R_i \leftrightarrow R_j} B$ then $\det(A) = -\det(B)$
- (4) $\det(I_n) = 1$.

This is the fastest algorithm for computing the det of a general matrix with known entries. But what if the matrix has unknown entries? This becomes tedious because you don't know if an entry is a pivot!

Eg: $\det \begin{pmatrix} -\lambda & 1 & 3 \\ 1 & 2-\lambda & 1 \\ 1 & 1 & -\lambda \end{pmatrix} = ?$ Is $-\lambda$ a pivot?

Cofactor expansion is a handy recursive formula for the determinant that is useful in this setting.

Recursive: Compute $\det(n \times n)$ by computing several $\det((n-1) \times (n-1))$.

Def: Let A be an $n \times n$ matrix.

- The (i,j) minor A_{ij} is the $(n-1) \times (n-1)$ matrix obtained by deleting the i^{th} row & j^{th} column.
- The (i,j) cofactor C_{ij} is

$$C_{ij} = (-1)^{i+j} \det(A_{ij})$$

- The cofactor matrix is the matrix C whose (i,j) entry is C_{ij} .

Eg: $A = \begin{pmatrix} 0 & 1 & 3 \\ 1 & 2 & 1 \\ 1 & 0 & 0 \end{pmatrix}$ $A_{21} = \begin{pmatrix} 0 & 1 & 3 \\ \cancel{1} & \cancel{2} & 1 \\ 1 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 3 \\ 1 & 0 \end{pmatrix}$

$$C_{21} = (-1)^{2+1} \det \begin{pmatrix} 1 & 3 \\ 1 & 0 \end{pmatrix} = -(-3) = 3$$

NB: $(-1)^{i+j}$ follows a checkerboard pattern:

$$\begin{pmatrix} + & - & + \\ - & + & - \\ + & - & + \end{pmatrix} \quad + \stackrel{+}{=} (-1)^{i+j} = 1$$
$$- \stackrel{-}{=} (-1)^{i+j} = -1$$

Thm (Cofactor Expansion): A is an $n \times n$ matrix, $a_{ij} = (i,j)$ entry of A , $C_{ij} = (i,j)$ cofactor.

(1) Cofactor expansion along the i^{th} row:

$$\det(A) = \sum_{j=1}^n a_{ij} C_{ij} = a_{i1} C_{i1} + a_{i2} C_{i2} + \dots + a_{in} C_{in}$$

(2) Cofactor expansion along the j^{th} column:

$$\det(A) = \sum_{i=1}^n a_{ij} C_{ij} = a_{1j} C_{1j} + a_{2j} C_{2j} + \dots + a_{nj} C_{nj}$$

Eg: $A = \begin{pmatrix} 0 & 1 & 3 \\ 1 & 2 & 1 \\ 1 & 1 & 0 \end{pmatrix}$

- Expand cofactors along the 3rd row:

$$\det(A) = 1 \cdot \det\begin{pmatrix} 1 & 3 \\ 2 & 1 \end{pmatrix} + 1 \cdot -\det\begin{pmatrix} 0 & 3 \\ 1 & 1 \end{pmatrix} + 0 \cdot \det\begin{pmatrix} 0 & 1 \\ 1 & 2 \end{pmatrix}$$

$$= 1 \cdot (1 - 6) - 1 \cdot (-3) = -2$$

- Expand cofactors along the 2nd column:

$$\det(A) = 1 \cdot -\det\begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} + 2 \cdot \det\begin{pmatrix} 0 & 3 \\ 1 & 0 \end{pmatrix} + 1 \cdot -\det\begin{pmatrix} 0 & 3 \\ 1 & 1 \end{pmatrix}$$

$$= 1 \cdot -(-1) + 2 \cdot (-3) + 1 \cdot -(-3) = 1 - 6 + 3 = -2$$

Remarks:

- (1) This is a recursive formula: $C_{ij} = \det((n-1) \times (n-1))$
- (2) You can compute $C_{ij} = (-1)^{i+j} \det(A_{ij})$ however you like: you'll always get the same number.
- (3) Expanding along any row or column gives you $\det(A)$ — always the same number.
- (4) This is handy when your matrix has unknown entries or a row/col with a lot of zeros — otherwise it's ridiculously slow = $O(n! \cdot n)$.

Eg: $\det \begin{pmatrix} -\lambda & 1 & 3 \\ 1 & 2-\lambda & 1 \\ 1 & 1 & -\lambda \end{pmatrix}$


$$\begin{aligned}
 &\text{expand} \\
 &\text{1st col} \quad (-\lambda) \det \begin{pmatrix} 2-\lambda & 1 \\ 1 & -\lambda \end{pmatrix} + 1 \cdot \det \begin{pmatrix} 1 & 3 \\ 1 & -\lambda \end{pmatrix} + 1 \cdot \det \begin{pmatrix} 1 & 3 \\ 2-\lambda & 1 \end{pmatrix} \\
 &= -\lambda((2-\lambda)(-\lambda) - 1) + 1 \cdot -(-\lambda - 3) + 1 \cdot (1 - 3(2-\lambda)) \\
 &= -\lambda(-2\lambda + \lambda^2 - 1) + (\lambda + 3) + 1 - 3(2-\lambda) \\
 &= -\lambda^3 + 2\lambda^2 + 5\lambda - 2
 \end{aligned}$$

In fact, for 3×3 matrices it's not so hard to compute the determinant when all entries are unknown:

$$\begin{aligned}
 \det \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix} &= a \det \begin{pmatrix} e & f \\ h & i \end{pmatrix} - b \det \begin{pmatrix} d & f \\ g & i \end{pmatrix} + c \det \begin{pmatrix} d & e \\ g & h \end{pmatrix} \\
 &= a(ei - fh) - b(di - fg) + c(dh - ge) \\
 &= aei + bfg + cdh - afh - bdi - ceq
 \end{aligned}$$

How to remember this?

Sarrus' Scheme:



To compute $\det(3 \times 3 \text{ matrix})$:

$$\begin{aligned}
 \det = & aei + bfg + cdh \\
 & - ceq - afh - bdi
 \end{aligned}$$

Sum the products of forward diagonals, subtract products of backwards diagonals.

Eg: $\det \begin{pmatrix} 0 & 1 & 3 \\ 1 & 2 & 1 \\ 1 & 1 & 0 \end{pmatrix} = 0 \cdot 2 \cdot 0 + 1 \cdot 1 \cdot 1 + 3 \cdot 1 \cdot 1 - 1 \cdot 2 \cdot 3 - 1 \cdot 1 \cdot 0 - 0 \cdot 1 \cdot 1$

 $= 4 - 6 = -2$

Warning: This only works for 3×3 matrices!

→ See the big formula at the end for $n \times n$ matrices.

Eg: $\det \begin{pmatrix} 2 & 5 & -3 & -1 \\ -2 & -3 & 2 & -5 \\ 1 & 3 & -2 & 0 \\ -1 & 6 & 4 & 0 \end{pmatrix}$ Column with lots of zeros

$= -1 \cdot -\det \begin{pmatrix} -2 & -3 & 2 \\ 1 & 3 & -2 \\ -1 & 6 & 4 \end{pmatrix} - 5 \cdot \det \begin{pmatrix} 2 & 5 & -3 \\ 1 & 3 & -2 \\ -1 & 6 & 4 \end{pmatrix}$

$+ 0 \cdot \det \begin{pmatrix} \text{don't} \\ \text{care} \end{pmatrix} + 0 \cdot \det \begin{pmatrix} \text{don't} \\ \text{care} \end{pmatrix}$

$= 1(-24) - 5(11) = -24 - 55 = -79$

only computed two 3×3 dets

Better: Do a row operation first!

$\det \begin{pmatrix} 2 & 5 & -3 & -1 \\ -2 & -3 & 2 & -5 \\ 1 & 3 & -2 & 0 \\ -1 & 6 & 4 & 0 \end{pmatrix} \xrightarrow{R_2 \leftarrow 5R_1} \det \begin{pmatrix} 2 & 5 & -3 & -1 \\ -12 & -28 & 17 & 0 \\ 1 & 3 & -2 & 0 \\ -1 & 6 & 4 & 0 \end{pmatrix}$

$= -1 \cdot -\det \begin{pmatrix} -12 & -28 & 17 \\ 1 & 3 & -2 \\ -1 & 6 & 4 \end{pmatrix} = -79$

only computed one 3×3 det

Methods for Computing Determinants

(1) Special formulas ($2 \times 2, 3 \times 3$)

→ best for small matrices, except 3×3 with lots of 0's

(2) Cofactor expansion

→ best if you have unknown entries, or a row/column with lots of zeros.

(3) Row (& column) operations

→ best if you have a big matrix with no unknown entries & no row or column with lots of zeros.

(4) Any combination of the above

→ e.g. do a row op. to create a column with lots of zeros, then expand cofactors, ...

Thm: Let C be the cofactor matrix of A . Then

$$AC^T = \det(A) I_n = C^T A$$

In particular, if $\det(A) \neq 0$, then

$$A^{-1} = \frac{1}{\det(A)} C^T$$

see supplement

→ Ridiculously inefficient computationally.

Eg: $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \rightarrow A^{-1} = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$

→ generalizes the formula for 2×2 inverse

Cross Products

This is an operation you can do to vectors in \mathbb{R}^3 .

Recall: the unit vectors in \mathbb{R}^3 are

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \quad e_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

Def: Let $v = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$ $w = \begin{pmatrix} d \\ e \\ f \end{pmatrix} \in \mathbb{R}^3$.

The cross product is

$$v \times w = \begin{pmatrix} bf - ec \\ cd - af \\ ae - bd \end{pmatrix} \in \mathbb{R}^3$$

So the cross product is (vector) \times (vector) \rightarrow (vector)

Here's how you remember it:

$$\begin{aligned} \begin{pmatrix} a \\ b \\ c \end{pmatrix} \times \begin{pmatrix} d \\ e \\ f \end{pmatrix} &= \text{"det} \begin{pmatrix} e_1 & e_2 & e_3 \\ a & b & c \\ d & e & f \end{pmatrix}" \text{ expand cofactors} \\ &= e_1 \det \begin{pmatrix} b & c \\ e & f \end{pmatrix} - e_2 \det \begin{pmatrix} a & c \\ d & f \end{pmatrix} + e_3 \det \begin{pmatrix} a & b \\ d & e \end{pmatrix} \\ &= (bf - ec)e_1 - (af - cd)e_2 + (ae - bd)e_3 \\ &= \begin{pmatrix} bf - ec \\ cd - af \\ ae - bd \end{pmatrix} \end{aligned}$$

$$\begin{aligned}
 \text{Eg: } \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \times \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} &= \det \begin{pmatrix} e_1 & e_2 & e_3 \\ 1 & 2 & 1 \\ 0 & 1 & 0 \end{pmatrix} \\
 &= e_1 \det \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix} - e_2 \det \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + e_3 \det \begin{pmatrix} 1 & 2 \\ 1 & 0 \end{pmatrix} \\
 &= -e_1 + e_2 - e_3 = \begin{pmatrix} -1 \\ 1 \\ -1 \end{pmatrix}
 \end{aligned}$$

Def: Let $u, v, w \in \mathbb{R}^3$. The triple product is

$$u \cdot (v \times w) = \det \begin{pmatrix} u^T & - \\ v^T & - \\ w^T & - \end{pmatrix}$$

Check: If $r = (a, b, c)$ $\omega = (d, e, f)$ $u = (g, h, i)$ then

$$u \cdot (v \times \omega)$$

$$= \begin{pmatrix} g & h & i \\ d & e & f \\ 1 & 1 & 1 \end{pmatrix} \cdot (e_1 \det \begin{pmatrix} b & c \\ e & f \end{pmatrix} - e_2 \det \begin{pmatrix} a & c \\ d & f \end{pmatrix} + e_3 \det \begin{pmatrix} a & b \\ d & e \end{pmatrix})$$

$$= g \det \begin{pmatrix} b & c \\ e & f \end{pmatrix} - h \det \begin{pmatrix} a & c \\ d & f \end{pmatrix} + i \det \begin{pmatrix} a & b \\ d & e \end{pmatrix}$$

$$= \det \begin{pmatrix} g & h & i \\ a & b & c \\ d & e & f \end{pmatrix} \quad \checkmark$$

$$\begin{aligned}
 \text{Eg: } u &= \begin{pmatrix} 0 \\ 1 \\ 3 \end{pmatrix} & r &= \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} & \omega &= \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
 v \times \omega &= \begin{pmatrix} -1 \\ 1 \\ -1 \end{pmatrix} & u \cdot (v \times \omega) &= \begin{pmatrix} 0 \\ 1 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 1 \\ -1 \end{pmatrix} &= 1 - 3 = -2 \\
 \det \begin{pmatrix} 0 & 1 & 3 \\ 1 & 2 & 1 \\ 1 & 0 & 0 \end{pmatrix} &= -2
 \end{aligned}$$

Properties:

(1) $\mathbf{v} \times \omega \perp \mathbf{v}$ and $\mathbf{v} \times \omega \perp \omega$


→ because $\mathbf{v} \cdot (\mathbf{v} \times \omega) = \det \begin{pmatrix} \mathbf{v}^T \\ \mathbf{v}^T \\ \omega^T \end{pmatrix} = 0$

(2) $\omega \times \mathbf{v} = -\mathbf{v} \times \omega$

→ because $\det \begin{pmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \\ -\omega^T & - \\ -\mathbf{v}^T & - \end{pmatrix} \stackrel{\text{no sweep}}{=} -\det \begin{pmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \\ \mathbf{v}^T & - \\ \omega^T & - \end{pmatrix}$

(3) $\|\mathbf{v} \times \omega\| = \|\mathbf{v}\| \cdot \|\omega\| \cdot \sin(\theta)$


→ compare $\mathbf{v} \cdot \omega = \|\mathbf{v}\| \cdot \|\omega\| \cdot \cos(\theta)$



(4) $\mathbf{v} \times \omega = 0 \iff \mathbf{v}, \omega$ are **collinear**

(then $\theta = 0$ or $180^\circ \iff \sin(\theta) = 0$)

(5) $\mathbf{v} \times \omega$ points in the direction determined by the **right hand rule**.



NB: (1), (3), & (5) characterize $\mathbf{v} \times \omega$.

The Big Formula

This is an explicit formula for $\det(A)$.

It's useful for some things but not practical — it has $n!$ terms!

Def: A **permutation** of $\{1, \dots, n\}$ is a **re-ordering**

$$\sigma: \{1, \dots, n\} \rightarrow \{1, \dots, n\}$$

$\sigma(i)$ = new number in i^{th} position

Eg:

$\begin{matrix} 1 & 2 & 3 & 4 \\ \downarrow & \downarrow & \downarrow & \downarrow \\ 3 & 1 & 2 & 4 \end{matrix}$	$\sigma(1) = 3$	$\sigma(3) = 2$
	$\sigma(2) = 1$	$\sigma(4) = 4$

Q: how many permutations of $\{1, \dots, n\}$ are there?

- n choices for 1st spot

- $(n-1)$ choices for 2nd spot

⋮

- 1 choice for last spot

So $n \cdot (n-1) \cdot \dots \cdot (1) = n!$

Eg: $n=3: 123 \rightarrow$

$$\underbrace{123 \quad 132 \quad 213 \quad 231 \quad 312 \quad 321}_{6 = 3 \cdot 2 \cdot 1 = 3!}$$

Def: A **transposition** is a permutation that just swaps two numbers.

Eg: $123 \rightarrow 132 \quad 213 \quad 321$

Fact: Any permutation can be obtained by doing some number of transpositions.

(Compare HW3 #5)

Def: The **sign** of a permutation σ is $\text{sign}(\sigma) =$

- $+1$ if it can be obtained by doing an **even** number of transpositions.
- -1 if it can be obtained by doing an **odd** number of transpositions.

Eg: $1234 \rightarrow 3124$:

$$1234 \rightarrow 3214 \rightarrow 3124$$

2 transpositions \Rightarrow sign is $+1$.

Thm (Big Formula): Let A be an $n \times n$ matrix with (ij) entry a_{ij} .

$$\det(A) = \sum_{\text{permutations}} \text{sign}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)}$$

Eg: $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$

Six permutations: $123 \rightarrow$

123	Sign = 1	(0 transpositions)
132	Sign = -1	(transposition)
213	Sign = -1	(transposition)
231	Sign = 1	(2 transpositions)
321	Sign = -1	(transposition)
312	Sign = 1	(2 transpositions)

$\det(A)_{11}$

$$\begin{aligned} & a_{11} a_{22} a_{33} \\ & - a_{11} a_{23} a_{32} \\ & - a_{12} a_{21} a_{33} \\ & + a_{12} a_{23} a_{31} \\ & - a_{13} a_{22} a_{31} \\ & + a_{13} a_{21} a_{32} \end{aligned}$$

