

Math 218D-1: Homework #9

due Wednesday, November 2, at 11:59pm

1. Compute

$$\det \left[\begin{pmatrix} -3 & 3 & 2 \\ 3 & 0 & 0 \\ -9 & 18 & 7 \end{pmatrix} - \lambda I_3 \right]$$

where λ is an unknown real number. Your answer will be a function of λ .

2. a) Compute the determinants of the matrices in HW8#14 in two more ways: by expanding cofactors along a row, and by expanding cofactors along a column. You should get the same answer using all three methods!
- b) Compute the determinants of the matrices in HW8#14(b) and (d) *again* using Sarrus' scheme.
- c) For the matrix of HW8#14(c), sum the products of the forward diagonals and subtract the products of the backward diagonals, as in Sarrus' scheme. Did you get the determinant?

3. Consider the matrix

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 3 & 1 \\ 0 & 1 & 2 \end{pmatrix}.$$

- a) Compute the cofactor matrix C of A .
- b) Compute AC^T . What is the relationship between C^T and A^{-1} ?
4. Consider the $n \times n$ matrix F_n with 1's on the diagonal, 1's in the entries immediately below the diagonal, and -1 's in the entries immediately above the diagonal:

$$F_2 = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \quad F_3 = \begin{pmatrix} 1 & -1 & 0 \\ 1 & 1 & -1 \\ 0 & 1 & 1 \end{pmatrix} \quad F_4 = \begin{pmatrix} 1 & -1 & 0 & 0 \\ 1 & 1 & -1 & 0 \\ 0 & 1 & 1 & -1 \\ 0 & 0 & 1 & 1 \end{pmatrix} \quad \dots$$

- a) Show that $\det(F_2) = 2$ and $\det(F_3) = 3$.
- b) Expand in cofactors to show that $\det(F_n) = \det(F_{n-1}) + \det(F_{n-2})$.
- c) Compute $\det(F_4)$, $\det(F_5)$, $\det(F_6)$, $\det(F_7)$ using **b**).

This shows that $\det(F_n)$ is the n th *Fibonacci number*. (The sequence usually starts with 1, 1, 2, 3, ..., so our $\det(F_n)$ is the usual $n + 1$ st Fibonacci number.)


5. Let A be an $n \times n$ invertible matrix with integer (whole number) entries.
- Explain why $\det(A)$ is an integer.
 - If $\det(A) = \pm 1$, show that A^{-1} has integer entries.
 - If A^{-1} has integer entries, show that $\det(A) = \pm 1$.

6. Let V be a subspace of \mathbf{R}^n . The matrix for reflection over V is

$$R_V = I_n - 2P_{V^\perp},$$

where $P_{V^\perp} = I_n - P_V$ is the projection matrix onto V^\perp .

- a) Suppose that V is the line in the picture. Draw the vectors $R_V x_1, R_V x_2, R_V x_3,$ and $R_V x_4$ as points in the plane.



- b) Show that any reflection matrix R_V is orthogonal.
 [Hint: Recall that $P_{V^\perp}^2 = P_{V^\perp} = P_{V^\perp}^T$.]
- c) Let V be the plane $x + y + z = 0$. Compute R_V and $\det(R_V)$.
- d) Let V be any plane in \mathbf{R}^3 . Prove that $\det(R_V) = -1$, as follows: choose an orthonormal basis $\{u_1, u_2\}$ for V , and let $u_3 = u_1 \times u_2$. Show that the matrix A with columns u_1, u_2, u_3 has determinant 1, and that $R_V A$ has determinant -1 .

Summary: a reflection over a plane in \mathbf{R}^3 has determinant -1 .

- e) Now compute $\det(R_L)$, where L is the x -axis in \mathbf{R}^3 .

7. Use a cross product to find an implicit equation for the plane

$$V = \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix} \right\}.$$

Compare HW6#6(a).

8. a) Let $v = (a, b)$ and $w = (c, d)$ be vectors in the plane, and let $A = \begin{pmatrix} a & c \\ b & d \end{pmatrix}$. By taking the cross product of $(a, b, 0)$ and $(c, d, 0)$, explain how the right-hand rule determines the sign of $\det(A)$.

b) Using the identity

$$\left[\begin{pmatrix} a \\ b \\ c \end{pmatrix} \times \begin{pmatrix} d \\ e \\ f \end{pmatrix} \right] \cdot \begin{pmatrix} g \\ h \\ i \end{pmatrix} = \det \begin{pmatrix} a & d & g \\ b & e & h \\ c & f & i \end{pmatrix},$$

explain how the right-hand rule determines the sign of a 3×3 determinant.

9. Decide if each statement is true or false, and explain why.

a) The determinant of the cofactor matrix of A equals the determinant of A .

b) $u \times v = v \times u$.

c) If $u \times v = 0$ then $u \perp v$.

10. For each matrix A and each vector v , decide if v is an eigenvector of A , and if so, find the eigenvalue λ .

a) $\begin{pmatrix} -20 & 42 & 58 \\ 1 & -1 & -3 \\ -1 & 18 & 26 \end{pmatrix}, \begin{pmatrix} 1 \\ 5 \\ -2 \end{pmatrix}$ b) $\begin{pmatrix} 2 & 3 & 0 \\ -5 & 4 & 2 \\ 3 & 3 & 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 3 \end{pmatrix}$

c) $\begin{pmatrix} -7 & 32 & -76 \\ 7 & -22 & 59 \\ 3 & -11 & 28 \end{pmatrix}, \begin{pmatrix} 3 \\ -2 \\ -1 \end{pmatrix}$ d) $\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}, \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}$

e) $\begin{pmatrix} -3 & 2 & -3 \\ 3 & -3 & -2 \\ -4 & 2 & -3 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$

11. For each matrix A and each number λ , decide if λ is an eigenvalue of A ; if so, find a basis for the λ -eigenspace of A .

a) $\begin{pmatrix} -5 & -14 \\ 3 & 8 \end{pmatrix}, \lambda = 1$ b) $\begin{pmatrix} -5 & -14 \\ 3 & 8 \end{pmatrix}, \lambda = -1$

c) $\begin{pmatrix} 2 & 3 & -15 \\ 5 & -7 & 31 \\ 2 & -3 & 13 \end{pmatrix}, \lambda = 3$ d) $\begin{pmatrix} 2 & 3 & -15 \\ 5 & -7 & 31 \\ 2 & -3 & 13 \end{pmatrix}, \lambda = 2$

e) $\begin{pmatrix} 3 & 1 & -2 \\ -2 & 0 & 4 \\ -1 & -1 & 4 \end{pmatrix}, \lambda = 2$ f) $\begin{pmatrix} 1 & 1 & -2 \\ -2 & -2 & 4 \\ -1 & -1 & 2 \end{pmatrix}, \lambda = 0$

g) $\begin{pmatrix} 7 & 0 & 0 \\ 0 & 7 & 0 \\ 0 & 0 & 7 \end{pmatrix}, \lambda = 7$ h) $\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \lambda = 0$

12. Suppose that A is an $n \times n$ matrix such that $Av = 2v$ for some $v \neq 0$. Let C be any invertible matrix. Consider the matrices

a) A^{-1} b) $A + 2I_n$ c) A^3 d) CAC^{-1} .

Show that v is an eigenvector of a)–c) and that Cv is an eigenvector of d), and find the eigenvalues.

13. Here is a handy trick for computing eigenvectors of a 2×2 matrix.

Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ be a 2×2 matrix with eigenvalue λ . Explain why $\begin{pmatrix} -b \\ a-\lambda \end{pmatrix}$ and $\begin{pmatrix} d-\lambda \\ -c \end{pmatrix}$ are λ -eigenvectors of A if they are nonzero.

For which matrices A does this trick fail?

14. a) Show that A and A^T have the same eigenvalues.
b) Give an example of a 2×2 matrix A such that A and A^T do not share any eigenvectors.
c) A *stochastic matrix* is a matrix with nonnegative entries such that the entries in each column sum to 1. Explain why 1 is an eigenvalue of a stochastic matrix.
[Hint: show that $(1, 1, \dots, 1)$ is an eigenvector of A^T .]

15. a) Find all eigenvalues of the matrix

$$\begin{pmatrix} 1 & -1 & 2 & 3 & 4 \\ 0 & 3 & -1 & -2 & -5 \\ 0 & 0 & 1 & 2 & 4 \\ 0 & 0 & 0 & 2 & 3 \\ 0 & 0 & 0 & 0 & -1 \end{pmatrix}.$$

- b) Explain how to find the eigenvalues of any triangular matrix.
16. Recall that an *orthogonal matrix* is a square matrix with orthonormal columns. Prove that any (real) eigenvalue of an orthogonal matrix Q is ± 1 .
17. Suppose that A is a square matrix such that A^k is the zero matrix for some $k > 0$. Show that 0 is the only eigenvalue of A .

- 18.** Decide if each statement is true or false, and explain why.
- a) If v, w are eigenvectors of a matrix A , then so is $v + w$.
 - b) An eigenvalue of $A + B$ is the sum of an eigenvalue of A and an eigenvalue of B .
 - c) An eigenvalue of AB is the product of an eigenvalue of A and an eigenvalue of B .
 - d) If $Ax = \lambda x$ for some vector x , then λ is an eigenvalue of A .
 - e) A matrix with eigenvalue 0 is not invertible.
 - f) The eigenvalues of A are equal to the eigenvalues of a row echelon form of A .