

# Solving Systems of Equations using Elimination

Here's a system of 3 equations in 3 variables:

$$\begin{cases} x_1 + 2x_2 + 3x_3 = 6 \\ 2x_1 - 3x_2 + 2x_3 = 14 \\ 3x_1 + x_2 - x_3 = -2 \end{cases}$$

How to solve it?

- **Substitution:** solve 1<sup>st</sup> equation for  $x_1$ , substitute into 2<sup>nd</sup> & 3<sup>rd</sup>, continue.
- **Elimination:** "combine" the equations to eliminate variables.

Elimination turns out to scale much better (to more equations & variables), so we'll focus on that.

"replace the 2<sup>nd</sup> equation with the 2<sup>nd</sup> minus 2x the 1<sup>st</sup>"

Eg:

$x_1 + 2x_2 + 3x_3 = 6$	$R_2 \leftarrow 2R_1$	$x_1 + 2x_2 + 3x_3 = 6$
$2x_1 - 3x_2 + 2x_3 = 14$	$\rightsquigarrow$	$-7x_2 - 4x_3 = 2$
$3x_1 + x_2 - x_3 = -2$		$3x_1 + x_2 - x_3 = -2$
	$R_3 \leftarrow 3R_1$	$x_1 + 2x_2 + 3x_3 = 6$
	$\rightsquigarrow$	$-7x_2 - 4x_3 = 2$
		$-5x_2 - 10x_3 = -20$

Now we have **eliminated**  $x_1$  from the 2<sup>nd</sup> & 3<sup>rd</sup> eq.s

These now form 2 equations in 2 variables: simpler!

$$\begin{array}{lcl} x_1 + 2x_2 + 3x_3 = 6 & R_3 \leftarrow \frac{5}{7}R_2 & x_1 + 2x_2 + 3x_3 = 6 \\ -7x_2 - 4x_3 = 2 & \searrow & -7x_2 - 4x_3 = 2 \\ -5x_2 - 10x_3 = -20 & & -\frac{50}{7}x_3 = -\frac{150}{7} \end{array}$$

We eliminated  $x_2$  from the last equation: now it's one equation in one variable. Easy!

We can now solve via back-substitution:

$$-\frac{50}{7}x_3 = -\frac{150}{7} \Rightarrow x_3 = 3.$$

Substitute into 2<sup>nd</sup> equation:

$$-7x_2 - 4x_3 = 2 \leadsto -7x_2 - 4 \cdot 3 = 2$$

Now solve for  $x_2$ :

$$-7x_2 - 12 = 2 \Rightarrow -7x_2 = 14 \Rightarrow x_2 = -2$$

Substitute both into 1<sup>st</sup> equation:

$$x_1 + 2x_2 + 3x_3 = 6 \leadsto x_1 + 2 \cdot (-2) + 3 \cdot 3 = 6$$

Now solve for  $x_1$ :

$$x_1 - 4 + 9 = 6 \Rightarrow x_1 = 1$$

Check:

$$\begin{array}{l} 1 + 2(-2) + 3(3) = 6 \\ 2 \cdot 1 - 3(-2) + 2(3) = 14 \\ 3 \cdot 1 + (-2) - 3 = -2 \end{array}$$



NB: In this case there was one solution - since we could

isolate each variable, all values were determined.

Does this always work?

Eg:  $\bullet$   $4x_2 + 3x_3 = 2$   $\swarrow$  swap  
 $x_1 + x_2 - x_3 = 3$   
 $2x_1 - 3x_2 - 6x_3 = -3$

$x_1$  is already eliminated from  $R_1$ . Fix: swap the 1<sup>st</sup> 2 eqns.

$R_1 \leftrightarrow R_2$

$$\begin{aligned} x_1 + x_2 - x_3 &= 3 \\ 4x_2 + 3x_3 &= 2 \\ 2x_1 - 3x_2 - 6x_3 &= -3 \end{aligned}$$

Now eliminate as before:

$R_3 \leftarrow 2R_1$

$$\begin{aligned} x_1 + x_2 - x_3 &= 3 \\ 4x_2 + 3x_3 &= 2 \\ \bullet -5x_2 - 4x_3 &= -9 \end{aligned}$$

$R_3 \leftarrow \frac{5}{4}R_2$

$$\begin{aligned} x_1 + x_2 - x_3 &= 3 \\ 4x_2 + 3x_3 &= 2 \\ \bullet -\frac{1}{4}x_3 &= -\frac{13}{2} \end{aligned}$$

Solve using back-substitution:

$$-\frac{1}{4}x_3 = -\frac{13}{2} \Rightarrow x_3 = 26$$

Substitute into 2<sup>nd</sup> equation:

$$4x_2 + 3(26) = 2 \Rightarrow x_2 = -19$$

Substitute both into 1<sup>st</sup> equation:

$$x_1 - 19 - 26 = 3 \Rightarrow x_1 = 48$$

NB again there is one solution.

Check:  $4x_2 + 3x_3 = 2$   $4(-19) + 3(26) = 2$   
 $x_1 + x_2 - x_3 = 3 \rightarrow 48 - 19 - 26 = 3$   
 $2x_1 - 3x_2 - 6x_3 = -3$   $2(48) - 3(-19) - 6(26) = -3$  ✓

Eg:  $x_1 + 2x_2 + 3x_3 = 1$   $R_2 \leftarrow 4R_1$   $x_1 + 2x_2 + 3x_3 = 1$   
 $4x_1 + 5x_2 + 6x_3 = 0$   $R_3 \leftarrow 7R_1$   $-3x_2 - 6x_3 = -4$   
 $7x_1 + 8x_2 + 9x_3 = -1$   $-6x_2 - 12x_3 = -8$   
 $R_3 \leftarrow 2R_2$   $x_1 + 2x_2 + 3x_3 = 1$   
 $-3x_2 - 6x_3 = -4$   
 $0 = 0$

Are we done? Yes: choose any value for  $x_3$ , then back-substitute to find  $x_1, x_2$ :

$$-3x_2 = -4 + 6x_3 \Rightarrow x_2 = \frac{4}{3} - 2x_3$$

$$x_1 = 1 - 2x_2 - 3x_3 = 1 - \frac{8}{3} + 4x_3 - 3x_3$$

$$x_1 = -\frac{5}{3} + x_3$$

Eg:  $x_3 = 1 \rightarrow x_1 = -\frac{2}{3}, x_2 = -\frac{2}{3}$

Check:  $-\frac{2}{3} - \frac{4}{3} + 3 = 1$   
 $-\frac{8}{3} - \frac{10}{3} + 6 = 0$   
 $-\frac{14}{3} - \frac{16}{3} + 9 = -1$  ✓

In this case there are infinitely many solutions. We'll deal with this in Week 3.

Eg:  $x_1 + 2x_2 + 3x_3 = 1$   
 $4x_1 + 5x_2 + 6x_3 = 0$   
 $7x_1 + 8x_2 + 9x_3 = 0$

$R_2 \leftarrow -4R_1$   
 $R_3 \leftarrow -7R_1$

$x_1 + 2x_2 + 3x_3 = 1$   
 $-3x_2 - 6x_3 = -4$   
 $-6x_2 - 12x_3 = -7$

tweak previous example

$R_3 \leftarrow -2R_2$

$x_1 + 2x_2 + 3x_3 = 1$   
 $-3x_2 - 6x_3 = -4$   
 $0 = 1$

If our original equations were true, then  $0 = 1$ .

Thus our system has no solutions.  
 (last 2 eqns are parallel planes)

Row Operations are the allowed manipulations we can perform on our equations.

(1)  $x_1 + 2x_2 + 3x_3 = 6$   
 $2x_1 - 3x_2 + 2x_3 = 14$   
 $3x_1 + x_2 - x_3 = -2$

$R_2 \leftarrow -2R_1$

$x_1 + 2x_2 + 3x_3 = 6$   
 $-7x_2 - 4x_3 = 2$   
 $3x_1 + x_2 - x_3 = -2$

row replacement  
 replace  $R_2$  by  $R_2 - 2R_1$

(2)  $x_1 + 2x_2 + 3x_3 = 6$   
 $2x_1 - 3x_2 + 2x_3 = 14$   
 $3x_1 + x_2 - x_3 = -2$

$R_1 \leftrightarrow R_2$

$2x_1 - 3x_2 + 2x_3 = 14$   
 $x_1 + 2x_2 + 3x_3 = 6$   
 $3x_1 + x_2 - x_3 = -2$

row swap  
 (change order)


$$\begin{array}{ll}
 (3) \quad x_1 + 2x_2 + 3x_3 = 6 & R_1 \times 2 \quad 2x_1 + 4x_2 + 6x_3 = 12 \\
 2x_1 - 3x_2 + 2x_3 = 14 & \rightsquigarrow 2x_1 - 3x_2 + 2x_3 = 14 \\
 3x_1 + x_2 - x_3 = -2 & 3x_1 + x_2 - x_3 = -2
 \end{array}$$

scalar multiplication  
(by nonzero scalar)

Obviously if  $(x_1, x_2, x_3)$  is a solution **before** doing a row operation, then it is true **after**. Eg row replacement:

$$\begin{array}{ll}
 x_1 + 2x_2 + 3x_3 \rightarrow 6 = 6 & R_2 \leftarrow 2R_1 \quad x_1 + 2x_2 + 3x_3 \rightarrow 6 = 6 \\
 2x_1 - 3x_2 + 2x_3 \rightarrow 14 = 14 & \rightsquigarrow \quad -7x_2 - 4x_3 \rightarrow 2 = 2
 \end{array}$$

**Fact:** All these operations are **reversible**: if you have a solution  $(x_1, x_2, x_3)$  **after** doing a row operation, then it's also a solution **before**.



This was the whole point: we wanted to **solve** our (original) system of equations!

**Questions:** How do you undo (reverse):

- $R_1 \leftrightarrow R_2$  ?     $R_1 \leftrightarrow R_2$
- $R_1 \times 2$  ?     $R_1 \div 2$
- $R_1 \leftarrow R_2$  ?     $R_1 \leftarrow R_2$

The variables  $x_1, x_2, \dots$  are just placeholders; only their **coefficients** matter. Let's extract them into a **matrix**.

## Three Ways to Write System of Linear Equations

(1) As a **system of equations**:

$$x_1 + 2x_2 + 3x_3 = 6$$

$$2x_1 - 3x_2 + 2x_3 = 14$$

(2) As a **matrix equation**  $Ax = b$

$$\underbrace{\begin{bmatrix} 1 & 2 & 3 \\ 2 & -3 & 2 \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_x = \underbrace{\begin{bmatrix} 6 \\ 14 \end{bmatrix}}_b$$

If you expand out the product you get

$$\begin{bmatrix} x_1 + 2x_2 + 3x_3 \\ 2x_1 - 3x_2 + 2x_3 \end{bmatrix} = \begin{bmatrix} 6 \\ 14 \end{bmatrix}$$

which is what we had before.

The **coefficient matrix**  $A$  comes from the **coefficients** of the variables:

$$\begin{bmatrix} 1 & 2 & 3 \\ 2 & -3 & 2 \end{bmatrix} \leftrightarrow \begin{matrix} 1x_1 + 2x_2 + 3x_3 \\ 2x_1 - 3x_2 + 2x_3 \end{matrix}$$

The vector  $x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$  contains the unknowns or variables.

NB:  $A$  is an  $m \times n$  matrix where

$m$  = # equations

$n$  = # variables

$b \in \mathbb{R}^m \leftarrow$  size  $m$

$x \in \mathbb{R}^n \leftarrow$  size  $n$

(3) As an augmented matrix.

This is a notational convenience: just squash  $A$  &  $b$  together and separate with a line.

$$\begin{bmatrix} 1 & 2 & 3 & | & 6 \\ 2 & -3 & 2 & | & 14 \end{bmatrix} \leftrightarrow \begin{cases} 1x_1 + 2x_2 + 3x_3 = 6 \\ 2x_1 - 3x_2 + 2x_3 = 14 \end{cases}$$

$$[A \mid b]$$

Augmented matrices are good for row operations, which only affect the coefficients (not the variables):

$$\begin{array}{lcl} x_1 + 2x_2 + 3x_3 = 6 & R_2 \leftarrow 2R_1 & x_1 + 2x_2 + 3x_3 = 6 \\ 2x_1 - 3x_2 + 2x_3 = 14 & & -7x_2 - 4x_3 = 2 \end{array}$$

III

$$\begin{bmatrix} 1 & 2 & 3 & | & 6 \\ 2 & -3 & 2 & | & 14 \end{bmatrix} \xrightarrow{R_2 \leftarrow 2R_1} \begin{bmatrix} 1 & 2 & 3 & | & 6 \\ 0 & -7 & -4 & | & 2 \end{bmatrix}$$



Eg: Let's solve the system from before using augmented matrices:

$$\begin{cases} x_1 + 2x_2 + 3x_3 = 6 \\ 2x_1 - 3x_2 + 2x_3 = 14 \\ 3x_1 + x_2 - x_3 = -2 \end{cases} \rightsquigarrow$$

$$\left[ \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right]$$

$$\left[ \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right] \xrightarrow{R_2 - 2R_1}$$

$$\left[ \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 3 & 1 & -1 & -2 \end{array} \right]$$

$$\xrightarrow{R_3 - 3R_1}$$

$$\left[ \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 0 & -5 & -10 & -20 \end{array} \right]$$

$$\xrightarrow{R_3 - \frac{5}{7}R_2}$$

$$\left[ \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 0 & 0 & -\frac{50}{7} & -\frac{150}{7} \end{array} \right]$$


$$\rightsquigarrow \begin{cases} x_1 + 2x_2 + 3x_3 = 6 \\ -7x_2 - 4x_3 = 2 \\ -\frac{50}{7}x_3 = -\frac{150}{7} \end{cases}$$

Now use back-substitution like before.

What does it mean to be "done"?  
(in terms of augmented matrices)

Def: A matrix is in **row echelon form (REF)** if

- (1) The first nonzero entry of each row is to the right of the row above it
- (2) All zero rows are at the bottom




REF: 
$$\begin{bmatrix} 1 & 2 & -1 & 4 \\ 0 & 0 & 3 & 12 \end{bmatrix} \quad \begin{bmatrix} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 0 & 0 & -\frac{50}{7} & -\frac{150}{7} \end{bmatrix}$$

Not REF: 
$$\begin{bmatrix} 1 & 2 & -1 & 4 \\ 2 & 0 & 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 1 & 2 & 3 & 6 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 5 \end{bmatrix}$$

Important: When checking if an **augmented matrix** is in REF, **ignore the augmentation line**.

$$\left[ \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & 0 & 3 & 12 \end{array} \right] \text{ REF? } \left[ \begin{array}{ccc|c} 1 & 2 & -1 & 4 \\ 0 & 0 & 3 & 12 \end{array} \right] \checkmark$$
  
Note: The vertical line in the first matrix is labeled "delete".

**Upshot:** The elimination procedure **terminates** when your (augmented) matrix is in **REF**.



**Def:** The **pivot positions (pivots)** of a matrix are the positions of the  $1^{\text{st}}$  nonzero entries of each row **after** you put it into REF.

$$\begin{bmatrix} \textcolor{red}{1} & 1 & -1 & 4 \\ 0 & 0 & \textcolor{red}{3} & 12 \end{bmatrix} \quad \begin{bmatrix} \textcolor{red}{1} & 2 & 3 & 6 \\ 0 & \textcolor{red}{-7} & -4 & 2 \\ 0 & 0 & \textcolor{red}{-\frac{50}{7}} & -\frac{150}{7} \end{bmatrix}$$
$$\begin{bmatrix} \textcolor{red}{1} & 2 & 3 & 1 \\ 0 & \textcolor{red}{-3} & -6 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$\textcolor{red}{\bullet} = \text{pivots}$

Remarkably, this is well-defined!

Def: The **rank** of a matrix is the number of pivots it has (in REF).

Eg: 
$$\begin{bmatrix} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{bmatrix} \xrightarrow[\text{(p.9)}]{\text{REF}} \begin{bmatrix} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 0 & 0 & -\frac{50}{7} & -\frac{150}{7} \end{bmatrix}$$

rank = 3

$$\begin{bmatrix} 1 & 2 & 3 & 1 \\ 4 & 5 & 6 & 0 \\ 7 & 8 & 9 & -1 \end{bmatrix} \xrightarrow[\text{(p.4)}]{\text{REF}} \begin{bmatrix} 1 & 2 & 3 & 1 \\ 0 & -3 & -6 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

rank = 2

## Number of Solutions (in terms of pivots)

The most basic question you can ask about a system of equations is: **how many** solutions does it have? This is entirely determined by the **pivot positions** / **pivot columns** (columns with a pivot).

(1) The system

$$\left[ \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 0 & 0 & -\frac{50}{7} & -\frac{150}{7} \end{array} \right]$$

(p.2)

had **one** solution. It has a pivot in every column except the augmented column.

This means every variable will be isolated when doing back-substitution.

(2) The system

$$\left[ \begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & -3 & -6 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right]$$

(p.5)

had **no** solutions. It has a pivot in the **augmented column**, which leads to the equation  $0=1$ .

$(\infty)$  The system

$$\left[ \begin{array}{ccc|c} 1 & 2 & 3 & 1 \\ 0 & -3 & -6 & -4 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

(p.4)

had infinitely many solutions. It has no pivot in the augmented column and no pivot in the column for the variable  $x_3$ . You can't isolate  $x_3$ , so you can choose any value.

**NB:** You have to put the system in REF to find its pivots, so you have to do work to know how many solutions there are.

**Def:** A system is consistent if it has at least 1 solution (so 1 or  $\infty$ ). It is inconsistent otherwise.