

Complex Eigenvalues

Some matrices have no (real) eigenvalues. But every matrix has a **complex** eigenvalue: any polynomial $p(\lambda)$ has a **complex** zero.

Eg: $A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ (ccw rotation by 90°)
 $p(\lambda) = \lambda^2 + 1 = (\lambda + i)(\lambda - i)$

Diagonalization still works great even if the eigenvalues are not real.

→ Still can solve difference equations & ODEs

→ Still get real-number answers

So we can apply diagonalization techniques to **more matrices** if we allow complex eigenvalues.

Fact: The **complex** eigenvalues & eigenvectors of a **real** matrix come in **complex conjugate pairs**:

$$Av = \lambda v \iff A\bar{v} = \bar{\lambda}\bar{v}$$

here $v = \begin{pmatrix} z_1 \\ \vdots \\ z_n \end{pmatrix} \rightsquigarrow \bar{v} = \begin{pmatrix} \bar{z}_1 \\ \vdots \\ \bar{z}_n \end{pmatrix}$

Eg: Solve the difference equation

$$v_{k+1} = A v_k \quad A = \begin{pmatrix} 0 & -1 \\ 3 & -3 \end{pmatrix} \quad v_0 = \begin{pmatrix} 2 \\ 3 \end{pmatrix} \quad \parallel \quad \text{No complex \#s in the statement!}$$

(1) Diagonalize:

$$p(\lambda) = \lambda^2 + 3\lambda + 3 \Rightarrow \lambda = \frac{1}{2}(-3 \pm \sqrt{9-12})$$

$$\rightarrow \lambda = \frac{1}{2}(-3 + i\sqrt{3}), \quad \bar{\lambda} = \frac{1}{2}(-3 - i\sqrt{3})$$

Find eigenvectors using the 2×2 trick

$$w = \begin{pmatrix} -b \\ a-\lambda \end{pmatrix} = \begin{pmatrix} 1 \\ -\lambda \end{pmatrix}$$

eigenvector for λ

$$\bar{w} = \begin{pmatrix} 1 \\ -\bar{\lambda} \end{pmatrix}$$

eigenvector for $\bar{\lambda}$

Check: $Aw = \begin{pmatrix} 0 & -1 \\ 3 & -3 \end{pmatrix} \begin{pmatrix} 1 \\ -\lambda \end{pmatrix} = \begin{pmatrix} \lambda \\ 3+3\lambda \end{pmatrix}$

Wait, is this equal to λw ?

$$\lambda w = \lambda \begin{pmatrix} 1 \\ -\lambda \end{pmatrix} = \begin{pmatrix} \lambda \\ -\lambda^2 \end{pmatrix} \stackrel{?}{=} \begin{pmatrix} \lambda \\ 3+3\lambda \end{pmatrix}$$

Yes: $-\lambda^2 = 3+3\lambda$ because

$$\lambda^2 + 3\lambda + 3 = p(\lambda) = 0 \quad \checkmark$$

So $\{w, \bar{w}\}$ is an **eigenbasis**

(different eigenvalues \Rightarrow LI)

(2) Expand the initial state in our eigenbasis:

We need to solve $\begin{pmatrix} 2 \\ 3 \end{pmatrix} = v_0 = x_1 \omega + x_2 \bar{\omega}$.

$$\leadsto \left(\begin{array}{cc|c} 1 & 1 & 2 \\ -\lambda & -\bar{\lambda} & 3 \end{array} \right) \xrightarrow{R_2 + \lambda R_1} \left(\begin{array}{cc|c} 1 & 1 & 2 \\ 0 & \lambda - \bar{\lambda} & 3 + 2\lambda \end{array} \right) \xrightarrow{\quad} = i\sqrt{3}$$

$$\leadsto \begin{cases} x_1 + x_2 = 2 \\ (i\sqrt{3})x_2 = i\sqrt{3} \end{cases} \xrightarrow{\quad} x_2 = 1 \rightarrow x_1 = 1$$

(back-substitution)

$$\text{So } v_0 = \omega_1 + \omega_2$$

$$\text{Solution: } A^k v_0 = \lambda^k \omega + \bar{\lambda}^k \bar{\omega}$$

So far it's exactly the same as for real eigenvalues!

... but we wanted a solution involving only real #s.

Thankfully, $\lambda^k \omega$ and $\bar{\lambda}^k \bar{\omega}$ are complex conjugates,
so


$$\begin{aligned} A^k v_0 &= \lambda^k \omega + \bar{\lambda}^k \bar{\omega} = 2 \operatorname{Re}[\lambda^k \omega] \\ &= 2 \operatorname{Re}\left[\lambda^k \begin{pmatrix} 1 \\ -\lambda \end{pmatrix}\right] = 2 \operatorname{Re}\begin{pmatrix} \lambda^k \\ -\lambda^{k+1} \end{pmatrix} \end{aligned}$$

Recall: Multiplication of complex numbers is much easier
in polar form.

$$\lambda = \frac{1}{2}(-3 + i\sqrt{3}) = r e^{i\theta}$$

$$r = \frac{1}{2}\sqrt{9+3} = \frac{1}{2}\sqrt{12} = \sqrt{3}$$

$$\theta = 150^\circ = 5\pi/6$$



Euler's Formula

$$\text{So } \lambda^k = r^k e^{ik \cdot \frac{5\pi}{6}} = (\sqrt{3})^k \left(\cos \frac{5k\pi}{6} + i \sin \frac{5k\pi}{6} \right)$$

$$\Rightarrow \operatorname{Re}(\lambda^k) = (\sqrt{3})^k \cos \frac{5k\pi}{6}$$

$$\Rightarrow v_k = 2 \begin{pmatrix} \sqrt{3}^k \cos(5k\pi/6) \\ -\sqrt{3}^{k+1} \cos(5(k+1)\pi/6) \end{pmatrix} \quad [\text{demo}]$$

The answer involves **only real numbers** (and cosines-weird!) but we needed complex numbers to get it!

Difference Equations with Complex Eigenvalues:

To solve $v_{k+1} = A v_k$:

(1-2) Diagonalize A and expand v_0 in an eigenbasis, as before. Complex numbers are OK.

→ Remember $Av = \lambda v \Leftrightarrow A\bar{v} = \bar{\lambda}\bar{v}$

(3) Group complex conjugate terms:

$$\lambda^k x \omega + \bar{\lambda}^k \bar{x} \bar{\omega} = 2 \operatorname{Re}(\lambda^k x \omega)$$


(4) Write λ in polar form:

$$\lambda = r e^{i\theta} \Rightarrow \lambda^k = r^k e^{ik\theta} = r^k (\cos k\theta + i \sin k\theta)$$

Multiply this by x and the coordinates of w and take the real part

we get an answer with sines & cosines (but no i 's).

Eg (for 4): $\lambda = 1+i$ $x = 3-2i$ $w = \begin{pmatrix} 1 \\ 2i \end{pmatrix}$



$$\lambda = \sqrt{2} e^{i\pi/4} \Rightarrow \lambda^k = (\sqrt{2})^k e^{ik\pi/4} = 2^{k/2} \left(\cos \frac{k\pi}{4} + i \sin \frac{k\pi}{4} \right)$$

$$\Rightarrow \lambda^k x w = 2^{k/2} \left(\cos \frac{k\pi}{4} + i \sin \frac{k\pi}{4} \right) (3-2i) \begin{pmatrix} 1 \\ 2i \end{pmatrix}$$

$$= 2^{k/2} \left[3 \cos \frac{k\pi}{4} + 2 \sin \frac{k\pi}{4} + i \left(3 \sin \frac{k\pi}{4} - 2 \cos \frac{k\pi}{4} \right) \right] \begin{pmatrix} 1 \\ 2i \end{pmatrix}$$

$$= 2^{k/2} \begin{pmatrix} 3 \cos \frac{k\pi}{4} + 2 \sin \frac{k\pi}{4} + i \left(3 \sin \frac{k\pi}{4} - 2 \cos \frac{k\pi}{4} \right) \\ -6 \sin \frac{k\pi}{4} + 4 \cos \frac{k\pi}{4} + i \left(6 \cos \frac{k\pi}{4} + 4 \sin \frac{k\pi}{4} \right) \end{pmatrix}$$

$$\Rightarrow 2 \operatorname{Re} [\lambda^k x w] = 2 \cdot 2^{k/2} \begin{pmatrix} 3 \cos \frac{k\pi}{4} + 2 \sin \frac{k\pi}{4} \\ -6 \sin \frac{k\pi}{4} + 4 \cos \frac{k\pi}{4} \end{pmatrix}$$

Algebraic & Geometric Multiplicity

Last we will discuss a **criterion for diagonalizability**.

(We like diagonalizable matrices because we can solve difference equations.)

Recall: If λ is a root of a polynomial $p(x)$, its **multiplicity** m is the largest power of $(x-\lambda)$ dividing p :
$$p(x) = (x-\lambda)^m \cdot (\text{other factors})$$

Eg: $p(\lambda) = -\lambda^3 + 3\lambda^2 - 4 = -(\lambda-2)^2(\lambda+1)^1$

$\lambda=2$ has multiplicity **2**; $\lambda=-1$ has multiplicity **1**

Def: Let A be an $n \times n$ matrix with eigenvalue λ .

(1) The **algebraic multiplicity (AM)** of λ is its multiplicity as a root of the characteristic polynomial $p(\lambda)$.

(2) The **geometric multiplicity (GM)** of λ is the dimension of the λ -eigenspace:

$$GM(\lambda) = \dim \text{Nul}(A - \lambda I_n)$$

$$= \# \text{ free variables in } A - \lambda I_n.$$

$$= \# \text{ linearly independent } \lambda\text{-eigenvectors}$$

Eg: $A = \begin{pmatrix} -7 & 3 & 5 \\ -10 & 5 & 6 \\ -9 & 3 & 7 \end{pmatrix}$ $p(\lambda) = -(\lambda-2)^2(\lambda-1)$

So the eigenvalues are 1 & 2.

- $\lambda = 1: AM = 1$.

$$\text{Nul}(A - 1I_3) = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}$$

$$AM \geq GM \checkmark$$

→ this is a line: $GM = 1$

- $\lambda = 2: AM = 2$

$$\text{Nul}(A - 2I_3) = \text{Span} \left\{ \begin{pmatrix} 3 \\ 4 \\ 3 \end{pmatrix} \right\}$$

$$AM \geq GM \checkmark$$

→ this is a line: $GM = 1$

This matrix is not diagonalizable:

only two linearly independent eigenvectors.

[demo]

Eg: $B = \begin{pmatrix} -4 & 3 & 2 \\ -6 & 5 & 2 \\ -6 & 3 & 4 \end{pmatrix}$ $p(\lambda) = -(\lambda-2)^2(\lambda-1)$

So the eigenvalues are 1 & 2.

- $\lambda = 1$: $AM = 1$

$$\text{Nul}(B - 1I_3) = \text{Span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}$$

$$AM \geq GM \checkmark$$

→ this is a line: $GM = 1$

- $\lambda = 2$: $AM = 2$

$$\text{Nul}(B - 2I_3) = \text{Span} \left\{ \begin{pmatrix} 3 \\ 4 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} \right\}$$

$$AM \geq GM \checkmark$$

→ this is a plane: $GM = 2$

This matrix is diagonalizable: an eigenbasis is

$$\left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 3 \\ 4 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} \right\}$$

[demo]

Both matrices have only 2 eigenvalues.

The difference is that B had $AM = GM = 2$ LI 2 eigenvectors and A had one.

Thm ($AM \geq GM$): For any eigenvalue λ of A ,
(algebraic multiplicity of λ)
 \geq (geometric multiplicity of λ) ≥ 1

For a proof, see the supplement.

NB: $GM \geq 1$ just says every eigenvalue has an eigenvector — the eigenspace can't be $\{0\}$, so its dimension is ≥ 1 .

Upshot: if $p(\lambda) = -(\lambda - 2)^2(\lambda - 1)^1$ then

- the 1-eigenspace is necessarily a line:

$$AM = 1 \geq GM \geq 1$$

- the 2-eigenspace is a line or a plane:

$$AM = 2 \geq GM \geq 1$$

- the matrix is diagonalizable $\iff GM(2) = 2$:
then you have $1 + 2 = 3$ LI eigenvectors.

Thm (AM/GM Criterion for Diagonalizability):

Let A be an $n \times n$ matrix.


- A is diagonalizable over the complex numbers
 $\iff AM(\lambda) = GM(\lambda)$ for every eigenvalue λ
- A is diagonalizable over the real numbers
 $\iff AM(\lambda) = GM(\lambda)$ for every eigenvalue λ
and A has no complex eigenvalues.

Eg: $A = \begin{pmatrix} -7 & 3 & 5 \\ -10 & 5 & 6 \\ -9 & 3 & 7 \end{pmatrix}$ is not diagonalizable because
 $AM(2) = 2 \neq 1 = GM(2)$

Corollary: If A has n different eigenvalues then
 A is diagonalizable.

Proof: If A has n different eigenvalues then

$$n = AM(\lambda_1) + \dots + AM(\lambda_n) \implies AM(\lambda_i) = 1$$

$$1 = AM(\lambda_i) \geq GM(\lambda_i) \geq 1 \implies AM(\lambda_i) = GM(\lambda_i) = 1$$


Eg: A 2×2 real matrix with a complex eigenvalue
 λ is diagonalizable (over \mathbb{C}): it has 2
eigenvalues λ and $\bar{\lambda}$.

Proof of the Theorem:

First note that

$$p(\lambda) = (-1)^n (\lambda - \lambda_1)^{m_1} \dots (\lambda - \lambda_r)^{m_r}$$

factors into linear factors (over \mathbb{C}), where $m_i = \text{AM}(\lambda_i)$. Hence

$$\text{AM}(\lambda_1) + \dots + \text{AM}(\lambda_r) = n \quad (\text{sum of the AM's is } n)$$

If A is diagonalizable then it has n LI eigenvectors,

$$\begin{aligned} \text{so } n &= \underbrace{\text{GM}(\lambda_1)}_{\text{AM}(\lambda_1)} + \dots + \underbrace{\text{GM}(\lambda_n)}_{\text{AM}(\lambda_n)} \\ &= \text{AM}(\lambda_1) + \dots + \text{AM}(\lambda_n) = n \end{aligned}$$

This forces $\text{AM}(\lambda_i) = \text{GM}(\lambda_i)$. Conversely, if each $\text{AM}(\lambda_i) = \text{GM}(\lambda_i)$ then

$$n = \text{GM}(\lambda_1) + \dots + \text{GM}(\lambda_n),$$

so when you combine eigenspace bases you get n LI eigenvectors.

