

## DYNAMICS IN A KINETIC MODEL OF ORIENTED PARTICLES WITH PHASE TRANSITION\*

AMIC FROUVELLE<sup>†</sup> AND JIAN-GUO LIU<sup>‡</sup>

**Abstract.** Motivated by a phenomenon of phase transition in a model of alignment of self-propelled particles, we obtain a kinetic mean-field equation which is nothing more than the Smoluchowski equation on the sphere with dipolar potential. In this self-contained article, using only basic tools, we analyze the dynamics of this equation in any dimension. We first prove global well-posedness of this equation, starting with an initial condition in any Sobolev space. We then compute all possible steady states. There is a threshold for the noise parameter: over this threshold, the only equilibrium is the uniform distribution, and under this threshold, the other equilibria are the Fisher–von Mises distributions with arbitrary direction and a concentration parameter determined by the intensity of the noise. For any initial condition, we give a rigorous proof of convergence of the solution to a steady state as time goes to infinity. In particular, when the noise is under the threshold and with nonzero initial mean velocity, the solution converges exponentially fast to a unique Fisher–von Mises distribution. We also found a new conservation relation, which can be viewed as a convex quadratic entropy when the noise is above the threshold. This provides a uniform exponential rate of convergence to the uniform distribution. At the threshold, we show algebraic decay to the uniform distribution.

**Key words.** Smoluchowski equation, nonlinear Fokker–Planck equation, dipolar potential, phase transition, LaSalle invariance principle, steady states, spontaneous symmetry breaking, Onsager’s phase transition

**AMS subject classifications.** 35K55, 35Q84, 35R01, 82B26, 82C26

**DOI.** 10.1137/110823912

**1. Introduction.** Phase transition and large time behavior of large interacting oriented/rod-like particle systems and their mean field limits have shown to be interesting in many physical and biological complex systems. Examples are paramagnetism to ferromagnetism phase transition near Curie temperature, nematic phase transition in liquid crystal or rod-shaped polymers, emerging of flocking dynamics near critical mass of self-propelled particles, etc.

The dynamics on orientation for self-propelled particles proposed by Vicsek et al. [22] to describe, for instance, fish schooling or bird flocking present such a behavior in numerical simulations. As the density increases (or as the noise decreases) and reaches a threshold, one can observe strong correlations between the orientations of particles. The model is discrete in time, and particles move at a constant speed following their orientation. At each time step, the orientation of each particle is updated, replaced by the mean orientation of its neighbors, plus a noise term.

A way to provide a time-continuous version of this dynamical system, which allows one to take a mean-field limit (and even a macroscopic limit), has been proposed by Degond and Motsch [7]. Instead of replacing the orientation at the next time step, they introduce a parameter playing the role of a rate of relaxation towards this mean

---

\*Received by the editors February 10, 2011; accepted for publication (in revised form) January 11, 2012; published electronically March 22, 2012.

<http://www.siam.org/journals/sima/44-2/82391.html>

<sup>†</sup>Institut de Mathématiques de Toulouse, CNRS – UMR 5219, Université de Toulouse, F-31062 Toulouse, France (amic.frouvelle@math.univ-toulouse.fr).

<sup>‡</sup>Department of Physics and Department of Mathematics, Duke University, Durham, NC 27707 (Jian-Guo.Liu@duke.edu). The research of this author was partially supported by NSF grant DMS 10-11738.

orientation. Unfortunately the mean-field limit of this model does not present phase transition. In [12], the first author of the present paper proved the robustness of the behavior of this model when this rate of relaxation depends on a local density. In particular, phase transition is still absent. However, when this parameter is set to be proportional to the local momentum of the neighboring particles, we will see that the model presents a phenomenon of phase transition as the intensity of the noise crosses a threshold. This phenomenon occurs on the orientation dynamics, so here we will consider only the spatial homogeneous dynamics.

In a joint work [6] with Pierre Degond, we have formally derived macroscopic limits for the inhomogeneous case. Using the results of the present paper, we have obtained, in the hydrodynamic limit, that the phase transition now appears as the local density crosses a threshold. Under this threshold, the local equilibria are uniform in orientation, and the corresponding macroscopic model is a nonlinear diffusion for the density. Above this threshold, the system is locally ordered, and the evolution of the local density and orientation is given by a nonconservative first-order system, which appears to be nonhyperbolic.

The particular model is described as follows: we have  $N$  oriented particles, described by vectors  $\omega_1, \dots, \omega_N$  belonging to  $\mathbb{S}$ , the unit sphere of  $\mathbb{R}^n$ , and satisfying the following system of coupled stochastic differential equations (which must be understood in the Stratonovich sense), for  $k \in \llbracket 1, N \rrbracket$ :

$$(1) \quad d\omega_k = (\text{Id} - \omega_k \otimes \omega_k) J_k dt + \sqrt{2\tau} (\text{Id} - \omega_k \otimes \omega_k) \circ dB_t^k,$$

$$(2) \quad J_k = \frac{1}{N} \sum_{j=1}^N \omega_j.$$

The term  $(\text{Id} - \omega_k \otimes \omega_k)$  denotes the projection on the hyperplane orthogonal to  $\omega_k$  and constrains the norm of  $\omega_k$  to be constant. The terms  $B_t^k$  stand for  $N$  independent standard Brownian motions on  $\mathbb{R}^n$ , and then the stochastic term  $(\text{Id} - \omega_k \otimes \omega_k) \circ dB_t^k$  represents the contribution of a Brownian motion on the sphere  $\mathbb{S}$  to the model. For more details on how to define Brownian motion on a Riemannian manifold, see [13].

Without this stochastic term, (1) can be written as

$$\dot{\omega}_k = \nabla_{\omega}(\omega \cdot J_k)|_{\omega=\omega_k},$$

where  $\nabla_{\omega}$  is the tangential gradient on the sphere (see the beginning of section 2.1 for some useful formulas on the unit sphere). So the model can be understood as a relaxation towards a unit vector in the direction of  $J_k$  subjected to a Brownian motion on the sphere with intensity  $\sqrt{2\tau}$ . The only difference with the model proposed in [7] (in the spatial homogeneous case) is that there  $J_k$  is replaced by  $\nu\Omega_k$ , where  $\Omega_k$  is the unit vector in the direction of  $J_k$  and the frequency of relaxation  $\nu$  is constant (or dependent on the local density in [12]). One point to emphasize is that, in that model, the interaction cannot be seen as a sum of binary interactions, contrary to the model presented here. Here the mean momentum  $J_k$  does not depend on the index  $k$  (but this is not true in the inhomogeneous case, where the mean is taken among the neighboring particles).

To simplify notation, we work with the uniform measure of total mass 1 on the sphere  $\mathbb{S}$ . We denote by  $f^N : \mathbb{R}_+ \times \mathbb{S} \rightarrow \mathbb{R}_+$  the probability density function (depending on time) associated with the orientation of one particle. Then, as the number  $N$  of particles tends to infinity,  $f^N$  tends to a probability density function  $f$  satisfying

$$(3) \quad \partial_t f = Q(f),$$

with

$$(4) \quad Q(f) = -\nabla_{\omega} \cdot ((\text{Id} - \omega \otimes \omega)J[f]f) + \tau \Delta_{\omega} f,$$

$$(5) \quad J[f] = \int_{\mathbb{S}} \omega f(\cdot, \omega) d\omega.$$

In the model of [7],  $J[f]$  is replaced just in (4) by  $\nu \Omega[f]$ , where  $\Omega[f]$  is the unit vector in the direction of  $J[f]$ .

The first term of  $Q(f)$  can be formally derived using a direct computation with the empirical distribution of particles. The diffusion part comes from Itô's formula. In a recent work [2], a rigorous derivation of this mean-field limit has been provided, even in the inhomogeneous case. This derivation is linked with the so-called propagation of chaos property. We refer the reader to [21] for an introduction to this notion. The laboratory example given in this reference is the original model of McKean [18], which is a more general version of our system in  $\mathbb{R}^n$  instead of  $\mathbb{S}$  (in that case, (3) is called the McKean–Vlasov equation). The main point is to adapt the theory in the framework of stochastic analysis on Riemannian manifolds.

Notice that (3) can be written in the form

$$\partial_t f = \nabla \cdot (f \nabla \Psi) + \tau \Delta f,$$

with

$$\Psi(\omega, t) = -\omega \cdot J(t) = \int_{\mathbb{S}} K(\omega, \bar{\omega}) f(t, \bar{\omega}) d\bar{\omega}.$$

This equation is known as the Smoluchowski equation (or the nonlinear Fokker–Planck equation) and was introduced by Doi [8] as the gradient flow equation for the Onsager free energy functional

$$(6) \quad \mathcal{F}(f) = \tau \int_{\mathbb{S}} f(\cdot, \omega) \ln f(\cdot, \omega) d\omega + \frac{1}{2} \int_{\mathbb{S} \times \mathbb{S}} K(\omega, \bar{\omega}) f(\cdot, \omega) f(\cdot, \bar{\omega}) d\omega d\bar{\omega}.$$

This functional was proposed by Onsager [19] to describe the equilibrium states of suspensions of rod-like polymers. They are given by the critical points of this functional.

Defining the chemical potential  $\mu$  as the first-order variation of  $\mathcal{F}(f)$  under the constraint  $\int_{\mathbb{S}} f = 1$ , we get  $\mu = \tau \ln f + \Psi$ , and the Smoluchowski equation becomes

$$\partial_t f = \nabla \cdot (f \nabla \mu).$$

In the original work of Onsager [19], the kernel has the form  $K(\omega, \bar{\omega}) = |\omega \times \bar{\omega}|$ , but there is another form, introduced later by Maier and Saupe [17], which leads to similar quantitative results:  $K(\omega, \bar{\omega}) = -(\omega \cdot \bar{\omega})^2$ . In our case, the potential given by  $K(\omega, \bar{\omega}) = -\omega \cdot \bar{\omega}$  is called the dipolar potential. This is a case where the arrow of the orientational direction has to be taken in account.

One of the interesting behaviors of the Smoluchowski equation is the phase transition bifurcation. This is indeed easy to see (here with the dipolar potential) from the following linearization around the uniform distribution: if  $f$  is a probability density function, the solution of (3), we write  $f = 1 + g$ , so  $\int_{\mathbb{S}} g d\omega = 0$  and we can get the equation for  $g$ . We multiply the equation by  $\omega$  and integrate using the formula  $\int_{\mathbb{S}} \omega \otimes \omega d\omega = \frac{1}{n} \text{Id}$  (this is a matrix with trace one and commuting with any

rotation) and the tools in the beginning of section 2.1. We get the linearized equation for  $g$  and  $J[g]$

$$\begin{aligned}\partial_t g &= \tau \Delta_\omega g + (n-1) \omega \cdot J[g] + O(g^2), \\ \frac{d}{dt} J[g] &= (n-1) \left( \frac{1}{n} - \tau \right) J[g] + O(g^2).\end{aligned}$$

Therefore if we take the linear part of this system, we can solve the second equation directly, and the first one becomes the heat equation with a known source term. Finally, around the constant state, the linearized Smoluchowski equation is stable if  $\tau \geq \frac{1}{n}$  and unstable if  $\tau < \frac{1}{n}$ . We expect to find another kind of equilibrium in this regime. The work has initially been done in [10] for the dimension  $n = 3$ ; the distribution obtained is called the Fisher–von Mises distribution [23].

A lot of work has been done to study the equilibrium states for the Maier–Saupe potential and in particular to show the axial symmetry of these steady states. A complete classification has been achieved for the two- and three-dimensional cases in [16] (see also [24], including the analysis of stability under a weak external shear flow). The interesting behavior, besides the phase transition, is the hysteresis phenomenon: before a first threshold, only one family of anisotropic equilibria is stable; then, in addition, the uniform equilibrium becomes stable, and after a second threshold, the only equilibrium is the uniform distribution. In the case of a coupling between the Maier–Saupe and the dipolar potentials, it is shown in [14], [15], [26] that the only stable equilibrium states are axially symmetric. To our knowledge, less work has been done to study the dynamics of the Smoluchowski equation, in particular the rate at which the solution converges to a steady state.

The purpose of this paper is to give a rigorous proof of the phase transition in any dimension for the dipolar potential and to study the large time dynamics and the convergence rates towards equilibrium states.

In section 2, we give some general results concerning (3). We provide a self-contained proof for existence and uniqueness of a solution with an initial nonnegative condition in any Sobolev space. We show that the solution is instantaneously positive and in any Sobolev space (and actually analytic in the space variable), and we obtain uniform bounds in time for each Sobolev norm.

In section 3, we use the Onsager free energy (decreasing in time) to analyze the general behavior of the solution as time goes to infinity. We prove a kind of LaSalle principle, implying that the solution converges, in the  $\omega$ -limit sense, to a given set of equilibria. We determine all the steady states and see that the value  $\frac{1}{n}$  is indeed a threshold for the noise parameter  $\tau$ . Over this threshold, the only equilibrium is the uniform distribution. When  $\tau < \frac{1}{n}$ , two kinds of equilibria exist: the uniform distribution, and a family of nonisotropic distributions (called Fischer–von Mises distributions), with a concentration parameter  $\kappa$  depending on  $\tau$ .

Finally, in section 4, we show that the solution converges strongly to a given equilibrium. We first obtain a new conservation relation, which plays the role of an entropy when  $\tau \geq \frac{1}{n}$  and shows a global convergence to the uniform distribution with rate proportional to  $\tau - \frac{1}{n}$ . Then we prove that, in the supercritical case  $\tau < \frac{1}{n}$ , the solution converges to a nonisotropic equilibrium if and only if the initial drift velocity  $|J[f_0]|$  is nonzero (if it is zero, the equation reduces to the heat equation, and the solution converges exponentially fast to the uniform distribution). We prove in that case that the convergence to this steady state is exponential in time, and we give the asymptotic rate of convergence. Finally, in the critical case  $\tau = \frac{1}{n}$ , we show that

the speed of convergence to the uniform distribution is algebraic (more precisely, the decay in any Sobolev norm is at least  $\frac{C}{\sqrt{t}}$ ).

**2. General results.**

**2.1. Preliminaries: Some results on the unit sphere.** This subsection consists essentially of a main lemma, allowing us to perform some estimates on the norm of integrals of the form  $\int_{\mathbb{S}} g \nabla_{\omega} h$ , where  $h$  and  $g$  are real functions with mean zero.

But let us start with some useful formulas. For  $V$  a constant vector in  $\mathbb{R}^n$ , we have

$$\begin{aligned} \nabla_{\omega}(\omega \cdot V) &= (\text{Id} - \omega \otimes \omega)V, \\ \nabla_{\omega} \cdot ((\text{Id} - \omega \otimes \omega)V) &= -(n - 1)\omega \cdot V, \end{aligned}$$

where  $\nabla_{\omega}$  (resp.,  $\nabla_{\omega} \cdot$ ) stands for the tangential gradient (resp., the divergence) on the unit sphere. When no confusion is possible, we will use just the notation  $\nabla$ . Then, taking the dot product with a given tangent vector field  $A$  or multiplying by a regular function  $f$  and integrating by parts, we get

$$\begin{aligned} \int_{\mathbb{S}} \omega \nabla_{\omega} \cdot A(\omega) d\omega &= - \int_{\mathbb{S}} A(\omega) d\omega, \\ \int_{\mathbb{S}} \nabla_{\omega} f d\omega &= (n - 1) \int_{\mathbb{S}} \omega f d\omega. \end{aligned}$$

We then introduce some notation. We denote by  $\dot{H}^s(\mathbb{S})$  the subspace composed of mean zero functions of the Sobolev space  $H^s(\mathbb{S})$ . This is a Hilbert space, associated with the inner product  $\langle g, h \rangle_{\dot{H}^s}^2 = \langle (-\Delta)^s g, h \rangle$ , where  $\Delta$  is the Laplace–Beltrami operator on the sphere. This also has a sense for any  $s \in \mathbb{R}$  by spectral decomposition of this operator. We will denote by  $\|\cdot\|_{\dot{H}^s}$  the norm on this Hilbert space.

We then define the so-called conformal Laplacian  $\tilde{\Delta}_{n-1}$  on the sphere (see [1]) which plays a role in some Sobolev inequalities. This is a positive definite operator (pseudodifferential operator of degree  $n - 1$ , mapping  $\dot{H}^s(\mathbb{S})$  continuously into  $\dot{H}^{s-n+1}(\mathbb{S})$ , which is a differential operator when  $n$  is odd) given by

$$(7) \quad \tilde{\Delta}_{n-1} = \begin{cases} \prod_{0 \leq j \leq \frac{n-3}{2}} (-\Delta + j(n-j-2)) & \text{for } n \text{ odd,} \\ (-\Delta + (\frac{n}{2} - 1)^2)^{\frac{1}{2}} \prod_{0 \leq j \leq \frac{n}{2}-2} (-\Delta + j(n-j-2)) & \text{for } n \text{ even.} \end{cases}$$

Equivalently, it can also be defined by

$$(8) \quad \tilde{\Delta}_{n-1} Y_{\ell} = \ell(\ell + 1) \dots (\ell + n - 2) Y_{\ell} \text{ for any spherical harmonic } Y_{\ell} \text{ of degree } \ell.$$

Here is the main lemma.

LEMMA 2.1 (estimates on the sphere, valid for any  $s \in \mathbb{R}$ ).

1. If  $h$  is in  $\dot{H}^{-s+1}(\mathbb{S})$  and  $g$  is in  $\dot{H}^s(\mathbb{S})$ , the following integral is well defined and we have

$$(9) \quad \left| \int_{\mathbb{S}} g \nabla h \right| \leq C \|g\|_{\dot{H}^s} \|h\|_{\dot{H}^{-s+1}},$$

where the constant  $C$  depends only on  $s$  and  $n$ .

2. We have the following estimation for any  $g \in \dot{H}^{s+1}(\mathbb{S})$ :

$$(10) \quad \left| \int_{\mathbb{S}} g \nabla (-\Delta)^s g \right| \leq C \|g\|_{\dot{H}^s}^2,$$

where the constant  $C$  depends only on  $s$  and  $n$ .

3. We have the following identity for any  $g \in \dot{H}^{-\frac{n-3}{2}}$ :

$$(11) \quad \int_{\mathbb{S}} g \nabla \tilde{\Delta}_{n-1}^{-1} g = 0.$$

Let us make some remarks on these statements. The first one expresses just the fact that the gradient operator (or, more precisely, any of its component  $e \cdot \nabla$  for a given unit vector  $e$ ) is well defined as an operator sending  $\dot{H}^{-s+1}(\mathbb{S})$  continuously into  $\dot{H}^{-s}(\mathbb{S})$  for any  $s$ .

The second one is actually a commutator estimate. It is equivalent to the fact that for any given unit vector  $e$  and for any  $g, h \in \dot{H}^{s+1}$ , we have

$$\left| \int_{\mathbb{S}} g e \cdot \nabla (-\Delta)^s h + h e \cdot \nabla (-\Delta)^s g \right| \leq \tilde{C} \|g\|_{\dot{H}^s} \|h\|_{\dot{H}^s}.$$

Defining the operator  $F$  by

$$Fg = e \cdot \nabla (-\Delta)^s g - (-\Delta)^s \nabla \cdot ((\text{Id} - \omega \otimes \omega)eg)$$

and integrating by parts, this inequality becomes  $|\int_{\mathbb{S}} h Fg| \leq \tilde{C} \|g\|_{\dot{H}^s} \|h\|_{\dot{H}^s}$ . In other words,  $F$  sends  $\dot{H}^s(\mathbb{S})$  continuously into  $\dot{H}^{-s}(\mathbb{S})$  for any  $s$ .

So since  $F = [e \cdot \nabla, (-\Delta)^s] + (n-1)(-\Delta)^s e \cdot \omega$ , this second statement (10) expresses that the commutator  $[\nabla, (-\Delta)^s]$  is an operator of degree  $2s$ .

With the same point of view, (11) gives an exact computation of the commutator of the gradient and the inverse of conformal Laplacian.

This says just that  $[\nabla, \tilde{\Delta}_{n-1}^{-1}] = -(n-1)\tilde{\Delta}_{n-1}^{-1}\omega$  or, multiplying left and right by  $\tilde{\Delta}_{n-1}$ , that  $[\nabla, \tilde{\Delta}_{n-1}] = (n-1)\omega\tilde{\Delta}_{n-1}$ .

The proof of this lemma relies on some computations on spherical harmonics and is given in section A.1.

**2.2. Existence, uniqueness, positivity, and regularity.** We present here a self-contained proof of well-posedness of the problem (3), working in any Sobolev space for the initial condition. Some analogous claims are given in [5], without proof, starting for a continuous nonnegative function. They are based on arguments of [3], stating that the Galerkin method based on spherical harmonics converges (exponentially fast) to the unique solution. They are weaker with respect to the initial conditions and the positivity but stronger for the regularity of the solution (analytic in space). As a remark we will give the same regularity result and prove it in section A.2.

**DEFINITION 2.2** (weak solution for some  $s \in \mathbb{R}$ ). For  $T > 0$ , the function  $f \in L^2((0, T), H^{s+1}(\mathbb{S})) \cap H^1((0, T), H^{s-1}(\mathbb{S}))$  is said to be a weak solution of (3) if, for almost all  $t \in [0, T]$ , we have for all  $h \in H^{-s+1}(\mathbb{S})$ ,

$$(12) \quad \langle \partial_t f, h \rangle = -\tau \langle \nabla_\omega f, \nabla_\omega h \rangle + \langle f, J[f] \cdot \nabla_\omega h \rangle,$$

where  $\langle \cdot, \cdot \rangle$  is the usual duality product for distributions on the sphere  $\mathbb{S}$ .

Since it is sometimes more convenient to work with mean zero functions (in order to use the main lemma of the previous subsection), we reformulate this problem in

another framework. We set  $f = 1 + g$  so that  $f$  is a weak solution if and only if  $g \in L^2((0, T), \dot{H}^{s+1}(\mathbb{S})) \cap H^1((0, T), \dot{H}^{s-1}(\mathbb{S}))$  with, for almost all  $t \in [0, T]$  and for all  $h \in \dot{H}^{-s+1}(\mathbb{S})$ ,

$$(13) \quad \langle \partial_t g, h \rangle = -\tau \langle \nabla_\omega g, \nabla_\omega h \rangle + (n - 1)J[g] \cdot J[h] + \langle g, J[g] \cdot \nabla_\omega h \rangle.$$

It makes sense to look for a weak solution with prescribed initial condition in  $H^s$ , since it always belongs to  $C([0, T], H^s(\mathbb{S}))$ , as stated by the following proposition.

**PROPOSITION 2.3.** *If  $g \in L^2((0, T), \dot{H}^{s+1}(\mathbb{S})) \cap H^1((0, T), \dot{H}^{s-1}(\mathbb{S}))$ , then, up to redefining it on a set of measure zero, it belongs to  $C([0, T], H^s(\mathbb{S}))$ , and we have*

$$\max_{[0, T]} \|u(t)\|_{H^s}^2 \leq C \int_0^T \|u\|_{\dot{H}^{s+1}}^2 + \|\partial_t u\|_{\dot{H}^{s-1}}^2,$$

where the constant  $C$  depends only on  $T$ .

The proof in the case  $s = 0$  is the same as in [9, section 5.9.2, Thm. 3]. To do the general case, we apply the result to  $(-\Delta)^{\frac{s}{2}}g$ .

**THEOREM 2.4.** *Given an initial probability measure  $f_0$  in  $H^s(\mathbb{S})$ , there exists a unique weak solution  $f$  of (3) such that  $f(0) = f_0$ . This solution is global in time. Moreover,  $f \in C^\infty((0, +\infty) \times \mathbb{S})$ , with  $f(t, \omega) > 0$  for all positive  $t$ .*

We also have the following instantaneous regularity and uniform boundedness estimates (for  $m \in \mathbb{N}$ , the constant  $C$  depends only on  $\tau, m$ , and  $s$ ) for all  $t > 0$ :

$$\|f(t)\|_{H^{s+m}}^2 \leq C \left(1 + \frac{1}{t^m}\right) \|f_0\|_{H^s}^2.$$

The proof consists of several steps, which we will treat as propositions. We first use a Galerkin method to prove existence on a small interval. We then show the continuity with respect to initial conditions on this interval (and hence the uniqueness). Next, we prove the positivity of  $1 + g$  for regular solutions. This gives us a better estimate of  $J[g]$ . Repeating the procedure on the following small interval, and so on, we can show that this extends to any  $t > 0$ . Regularizing the initial condition then gives global existence in any case.

We finally obtain the instantaneous regularity and boundary estimates by decomposing the solution between low and high modes.

For the proof of all propositions, we will denote by  $C_0, C_1, \dots$  some positive constants which depend only on  $s$  and  $\tau$ . We will also fix one parameter  $K > 0$  (which will be a bound on the norm of initial condition) and denote by  $M_0, M_1, \dots$  some positive constants which depend only on  $s, \tau$ , and  $K$ .

**PROPOSITION 2.5** (existence: Galerkin method). *We set*

$$(14) \quad T = \frac{1}{C_1} \ln \left(1 + \frac{1}{1 + 2C_2K}\right),$$

where the constants  $C_1$  and  $C_2$  will be defined later.

*If  $\|g_0\|_{\dot{H}^s} \leq K$ , then we have existence of a weak solution on  $[0, T]$  satisfying (13), uniformly bounded in  $L^2((0, T), \dot{H}^{s+1}(\mathbb{S})) \cap H^1((0, T), \dot{H}^{s-1}(\mathbb{S}))$  by a constant  $M_1$ .*

*Proof.* We denote by  $P_N$  the space spanned by the first  $N$  (nonconstant) eigenvectors of the Laplace–Beltrami operator. This is a finite-dimensional vector space, included in  $\dot{H}^p(\mathbb{S})$  for all  $p$  and containing the functions of the form  $\omega \mapsto V \cdot \omega$  (see section A.1 for more details).

Let  $g^N \in C^1(I, P_N)$  be the unique solution of the following Cauchy problem, defined on a maximal interval  $I \subset \mathbb{R}_+$  (“nonlinear” ODE on a finite-dimensional space):

$$\begin{cases} \frac{d}{dt}g^N = \Pi_N [\tau \Delta_\omega g^N - \nabla_\omega \cdot ((\text{Id} - \omega \otimes \omega)J[g^N](1 + g^N))], \\ g^N(0) = \Pi_N(g_0), \end{cases}$$

where  $\Pi_N$  is the orthogonal projection on  $P_N$ . The first equation is equivalent to the fact that for any  $h \in P_N$ , we have

$$(15) \quad \frac{d}{dt} \langle g^N, h \rangle = -\tau \langle \nabla_\omega g^N, \nabla_\omega h \rangle + (n-1)J[g^N] \cdot J[h] + \langle g^N, J[g^N] \cdot \nabla_\omega h \rangle.$$

The goal is to prove that  $[0, T] \subset I$  and that there exists an extracted sequence  $N_k$  such that, as  $k \rightarrow \infty$ ,

- $g^{N_k}$  converges weakly in  $L^2((0, T), \dot{H}^{s+1}(\mathbb{S}))$  to a function  $g$ ,
- $\partial_t g^{N_k}$  converges weakly to  $\partial_t g$  in  $L^2((0, T), \dot{H}^s(\mathbb{S}))$ , and
- $J[g^{N_k}] \rightarrow J[g]$  uniformly.

We have that  $(-\Delta)^s g^N \in P_N$ , so we can take it for  $h$ , put it in (15), and use the second part of Lemma 2.1 to get

$$(16) \quad \frac{1}{2} \frac{d}{dt} \|g^N\|_{\dot{H}^s}^2 + \tau \|g^N\|_{\dot{H}^{s+1}}^2 \leq C_0 |J[g^N]| \|g^N\|_{\dot{H}^s}^2 + (n-1)^s |J[g^N]|^2$$

$$(17) \quad \leq C_1 \|g^N\|_{\dot{H}^s}^2 (1 + C_2 \|g^N\|_{\dot{H}^s}).$$

Indeed, any component of  $\omega$  belongs to any  $\dot{H}^{-s}$ ; then  $J[g^N] = \langle \omega, g^N \rangle$  is controlled by any  $\dot{H}^s$  norm of  $g^N$ .

Solving this inequality, we obtain for  $0 \leq t < C_1^{-1} \ln(1 + (C_2 \|\Pi^N(g_0)\|_{\dot{H}^s})^{-1})$ ,

$$(18) \quad \|g^N\|_{\dot{H}^s} \leq \frac{\|\Pi^N(g_0)\|_{\dot{H}^s}}{e^{-C_1 t} - C_2 \|\Pi^N(g_0)\|_{\dot{H}^s} (1 - e^{-C_1 t})}.$$

Then we have  $\|g^N(t)\|_{\dot{H}^s} \leq 2\|g_0\|_{\dot{H}^s}$  for all  $t$  in  $[0, T]$ . There is no finite-time blow up in  $[0, T]$ ; then the ODE (15) has a solution on  $[0, T]$  for any  $N \in \mathbb{N}$ .

Now we denote by  $M_0$  a bound for  $|J[g^N]|$  on  $[0, T]$ . The inequality (16) gives

$$\frac{d}{dt} \|g^N\|_{\dot{H}^s}^2 + 2\tau \|g^N\|_{\dot{H}^{s+1}}^2 \leq (1 + M_0)C_3 \|g^N\|_{\dot{H}^s}^2.$$

Solving this inequality, we get for  $t \in [0, T]$ ,

$$(19) \quad \|g^N\|_{\dot{H}^s}^2 + 2\tau \int_0^t \|g^N\|_{\dot{H}^{s+1}}^2 \leq \|g_0\|_{\dot{H}^s}^2 e^{(1+M_0)C_3 t}.$$

We then use the ODE (15) to control the derivative of  $g$ . Taking  $h \in \dot{H}^{-s+1}(\mathbb{S})$ , we write  $h^N = \Pi_N(h)$ , and we get

$$\begin{aligned} \langle \partial_t g^N, h \rangle &= \langle \partial_t g^N, h^N \rangle \\ &\leq \tau \|g^N\|_{\dot{H}^{s+1}} \|h^N\|_{\dot{H}^{-s+1}} + C_4 \|g^N\|_{\dot{H}^s} \|h^N\|_{\dot{H}^{-s+1}} + M_0 \|g^N\|_{\dot{H}^s} \|h^N\|_{\dot{H}^{-s+1}} \\ &\leq (\tau \|g^N\|_{\dot{H}^{s+1}} + C_4 \|g^N\|_{\dot{H}^s} + M_0 \|g^N\|_{\dot{H}^s}) \|h\|_{\dot{H}^{-s+1}}, \end{aligned}$$



and thus we obtain

$$\|\partial_t g^N\|_{\dot{H}^{s-1}}^2 \leq 2\tau^2 \|g^N\|_{\dot{H}^{s+1}}^2 + 2(C_4 + M_0)^2 \|g^N\|_{\dot{H}^s}^2.$$

Integrating in time, we get, together with the estimate (19),

$$\int_0^T \|\partial_t g^N\|_{\dot{H}^{s-1}}^2 \leq \left[ \tau + \frac{2(C_4 + M_0)^2}{(1 + M_0)C_3} \right] \|g_0\|_{\dot{H}^s}^2 e^{(1 + M_0)C_3 T}.$$

Then we can take  $M_1^2 = K^2 e^{(1 + M_0)C_3 T} \max(\tau^{-1}, \tau + \frac{2(C_4 + M_0)^2}{(1 + M_0)C_3})$ , and we get that  $g^N$  is bounded by  $M_1$  in  $L^2((0, T), \dot{H}^{s+1}(\mathbb{S})) \cap H^1((0, T), \dot{H}^{s-1}(\mathbb{S}))$ .

Now, we need just estimates for  $\frac{d}{dt} J[g^N]$ . We can take  $h = \omega \cdot V$  for any constant vector  $V$  in the ODE (15) and use the tools given in the beginning of this section. We finally get

$$\begin{aligned} \left| \frac{d}{dt} J[g^N] \right| &= \left| \frac{n-1}{n} (1 - \tau n) J[g^N] - \int_{\mathbb{S}} (\text{Id} - \omega \otimes \omega) J[g^N] g^N \, d\omega \right| \\ &\leq (C_5 + M_0 C_6) \|g_0\|_{\dot{H}^s} e^{\frac{1}{2}(1 + M_0)C_3 T}. \end{aligned}$$

Indeed, again, since any component of  $\text{Id} - \omega \otimes \omega$  is in  $\dot{H}^{-s}$ , we can control the term  $\int_{\mathbb{S}} (\text{Id} - \omega \otimes \omega) g^N \, d\omega$  by any  $\dot{H}^s$  norm of  $g^N$ , uniformly in  $N$  and in  $t \in [0, T]$ .

In summary if we suppose that  $g_0$  is in  $\dot{H}^s(\mathbb{S})$ , for some  $s \in \mathbb{R}$ , we have that  $g^N$  is bounded in  $L^2((0, T), \dot{H}^{s+1}(\mathbb{S})) \cap H^1((0, T), \dot{H}^{s-1}(\mathbb{S}))$ , and that  $J[g^N]$  and  $\frac{d}{dt} J[g^N]$  are uniformly bounded in  $N$  and  $t \in [0, T]$ .

Then, using weak compactness and the Ascoli–Arzela theorem, we can find an increasing sequence  $N_k$ , a function  $g \in L^2((0, T), \dot{H}^{s+1}(\mathbb{S})) \cap H^1((0, T), \dot{H}^{s-1}(\mathbb{S}))$ , and a continuous function  $J : [0, T] \rightarrow \mathbb{R}^n$  such that, as  $k \rightarrow \infty$ ,

- $J[g^{N_k}]$  converges uniformly to  $J$  on  $[0, T]$ , and
- $g^{N_k}$  converges weakly to  $g$  in  $L^2((0, T), \dot{H}^{s+1}(\mathbb{S}))$  and in  $H^1((0, T), \dot{H}^{s-1}(\mathbb{S}))$ .

The limit  $g$  is also bounded by  $M_1$  in  $L^2((0, T), \dot{H}^{s+1}(\mathbb{S})) \cap H^1((0, T), \dot{H}^{s-1}(\mathbb{S}))$ .

Then, since we have  $\int_0^T \int_{\mathbb{S}} \varphi(t) \omega (g^{N_k} - g) \, d\omega \, dt \rightarrow 0$  for any smooth function  $\varphi$ , we get  $\int_0^T \varphi(t) (J[g] - J) \, dt = 0$ , and so  $J = J[g]$ .

For a fixed  $h \in P_M$  passing the weak limit in (15) (for  $N_k \geq M$ ), we get for almost every  $t \in [0, T]$  that

$$\forall h \in P_M, \quad \langle \partial_t g, h \rangle = -\tau \langle \nabla_\omega g, \nabla_\omega h \rangle + (n - 1) J[g] \cdot J[h] + \langle g, J[g] \cdot \nabla_\omega h \rangle.$$

This is valid for any  $M$  (except on a countable union of subsets of  $[0, T]$  of zero measure). By density (and using the first part of Lemma 2.1), we have that  $g$  is a weak solution of our problem.

Now for any  $h \in \dot{H}^{-s+1}(\mathbb{S})$ , we have that  $\langle g^N(t) - \Pi_N(g_0), h \rangle = \int_0^t \langle \partial_t g^N, h \rangle$  is controlled by  $M_1 \sqrt{t} \|h\|_{\dot{H}^{-s+1}}$ , uniformly in  $N$ . So, passing to the limit, we get that  $g(t) \rightarrow g_0$  in  $\dot{H}^{-s+1}(\mathbb{S})$  as  $t \rightarrow 0$ . But since we know that  $g \in C([0, T], H^s(\mathbb{S}))$ , by uniqueness, we get  $g(0) = g_0$ .  $\square$

**PROPOSITION 2.6** (continuity with respect to the initial condition). *Suppose that we set  $T = \frac{1}{C_1} \ln(1 + \frac{1}{1 + 2C_2 K})$  as in (14) and that we have two solutions  $g$  and  $\tilde{g}$ , with  $\|g(0)\|_{\dot{H}^s} \leq K$  and  $\|\tilde{g}(0)\|_{\dot{H}^s} \leq K$ .*

*Then there exists a constant  $M_3$  such that  $g - \tilde{g}$  is bounded in  $L^2((0, T), \dot{H}^{s+1}(\mathbb{S}))$  and in  $H^1((0, T), \dot{H}^{s-1}(\mathbb{S}))$  by  $M_3 \|g(0) - \tilde{g}(0)\|_{\dot{H}^s}$ .*

This automatically gives uniqueness of a weak solution on  $(0, T)$  with initial condition  $g_0$ .

*Proof.* Putting  $h = (-\Delta)^s g \in \dot{H}^{-s+1}$  in (13), we do the same estimations as in the previous proposition. We have the same estimate as (16)–(17):

$$(20) \quad \frac{1}{2} \frac{d}{dt} \|g\|_{\dot{H}^s}^2 + \tau \|g\|_{\dot{H}^{s+1}}^2 \leq C_0 |J[g]| \|g\|_{\dot{H}^s}^2 + (n-1)^s |J[g]|^2$$

$$(21) \quad \leq C_1 \|g\|_{\dot{H}^s}^2 (1 + C_2 \|g\|_{\dot{H}^s}).$$

So if we set  $T = C_1^{-1} \ln(1 + (1 + 2C_2K)^{-1})$ , we can solve this inequality on  $[0, T]$ , exactly as in (18). These solutions are then uniformly bounded in  $L^2((0, T), \dot{H}^{s+1}(\mathbb{S}))$  and in  $H^1((0, T), \dot{H}^{s-1}(\mathbb{S}))$  (by the constant  $M_1$ ).

Taking  $u = g - \tilde{g}$  and using (13) gives an equation for  $u$ : for almost all  $t \in [0, T]$  and for all  $h \in H^{-s}(\mathbb{S})$ ,

$$(22) \quad \langle \partial_t u, h \rangle = -\tau \langle \nabla_\omega u, \nabla_\omega h \rangle + (n-1)J[u] \cdot J[h] + \langle u, J[g] \cdot \nabla_\omega h \rangle + \langle \tilde{g}, J[u] \cdot \nabla_\omega h \rangle.$$

Now we take  $h = (-\Delta)^s u$  and use the first and second parts of Lemma 2.1 to get

$$(23) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \|u\|_{\dot{H}^s}^2 + \tau \|u\|_{\dot{H}^{s+1}}^2 &\leq (1 + M_1)C_3 \|u\|_{\dot{H}^s}^2 + C_7 \|u\|_{\dot{H}^s} \|\tilde{g}\|_{\dot{H}^{s+1}} \|(-\Delta)^s u\|_{\dot{H}^{-s}} \\ &\leq M_2 (1 + \|\tilde{g}\|_{\dot{H}^{s+1}}) \|u\|_{\dot{H}^s}^2. \end{aligned}$$

Grönwall’s lemma then gives the following estimate:

$$\begin{aligned} \|u\|_{\dot{H}^s}^2 + \tau \int_0^T \|u\|_{\dot{H}^{s+1}}^2 &\leq \|u_0\|_{\dot{H}^s}^2 \exp \left( M_2 \int_0^T (1 + \|\tilde{g}\|_{\dot{H}^{s+1}}) \right) \\ &\leq \|u_0\|_{\dot{H}^s}^2 e^{M_2(T+M_1^2)}. \end{aligned}$$

Using (22), we get that  $u$  is bounded in  $L^2((0, T), \dot{H}^{s+1}(\mathbb{S})) \cap H^1((0, T), \dot{H}^{s-1}(\mathbb{S}))$  by a constant  $M_3$  times  $\|u(0)\|_{\dot{H}^s}$ .  $\square$

**PROPOSITION 2.7** (positivity for regular solutions (maximum principle)). *Suppose that  $g_0$  is in  $\dot{H}^s(\mathbb{S})$ , with  $s$  sufficiently large (according to the Sobolev embeddings, so  $s > \frac{n+3}{2}$  is enough) so that the (unique) solution belongs to  $C^0([0, T], C^2(\mathbb{S}))$ . Here  $T$  is defined as in (14), with  $K = \|g_0\|_{\dot{H}^s}$ . We go back to the original formulation  $f = 1 + g$ . Then  $f$  is a classical solution of (3).*

*If  $f_0$  is nonnegative, then  $f$  is positive for any positive time, and, more precisely, we have the following estimates for all  $t \in (0, T]$  and  $\omega \in \mathbb{S}$  (if  $f_0$  is not equal to the constant function 1):*

$$(24) \quad e^{-(n-1) \int_0^t |J[f]|} \min_{\mathbb{S}} f_0 < f(t, \omega) < e^{(n-1) \int_0^t |J[f]|} \max_{\mathbb{S}} f_0.$$

*Proof.* Since the solution is in  $C^0([0, T], C^2(\mathbb{S}))$ , we can do the reverse integration by parts in the weak formulation (12). We get that, as an element of  $L^2((0, T), H^{s-1}(\mathbb{S}))$ , the function  $\partial_t f$  is equal (almost everywhere) to  $\tau \Delta_\omega f - \nabla_\omega \cdot ((\text{Id} - \omega \otimes \omega)J[f]f)$ , which is an element of  $C^0([0, T] \times \mathbb{S})$ . So up to redefining it on a set of measure zero, the function  $f$  belongs to  $C^1([0, T], C(\mathbb{S})) \cap C^0([0, T], C^2(\mathbb{S}))$  and satisfies the PDE.

Applying the chain rule and using the tools given in the beginning of this section, we get another formulation of the PDE (3):

$$(25) \quad \partial_t f = \tau \Delta_\omega f - J[f] \cdot \nabla_\omega f + (n-1)J[f] \cdot \omega f.$$

The next part of the proposition is just a classical strong maximum principle. Here we prove only the left part of the inequality; the other part is very similar, once we have that  $f$  is positive.

Suppose first that  $f_0$  is positive. We denote by  $\tilde{T} > 0$  the first time that the minimum on the unit sphere of  $f$  is zero (or  $\tilde{T} = T$  if  $f$  is always positive).

Then we have for  $t \in [0, \tilde{T}]$  that  $\partial_t f \geq \tau \Delta_\omega f - J[f] \cdot \nabla_\omega f - (n-1)|J[f]|f$ . If we write  $\tilde{f} = f e^{-(n-1) \int_0^t |J[f]|}$ , we get

$$(26) \quad \partial_t \tilde{f} \geq \tau \Delta_\omega \tilde{f} - J[f] \cdot \nabla_\omega \tilde{f}.$$

Then the weak maximum principle (see [9, section 7.1.4, Thm. 8], which is also valid on the sphere) gives us that the minimum of  $\tilde{f}$  on  $[0, \tilde{T}] \times \mathbb{S}$  is reached on  $\{0\} \times \mathbb{S}$ . That means that we have a nonstrict version of the left part of the inequality (24):

$$(27) \quad \forall t \in [0, \tilde{T}], \forall \omega \in \mathbb{S}, \quad f(\omega, t) \geq e^{-(n-1) \int_0^t |J[f]|} \min_{\mathbb{S}} f_0.$$

Consequently, we have that  $\min_{\mathbb{S}} f(\tilde{T}) > 0$ , and so  $\tilde{T} = T$ . If now  $f_0$  is only nonnegative, take  $f_0^\varepsilon = \frac{f_0 + \varepsilon}{1 + \varepsilon}$ , and by continuity with respect to initial condition, inequality (27) is still valid. That gives that  $f$  is nonnegative on  $[0, T]$ , and consequently we have that inequality (26) is valid on  $[0, \tilde{T}]$ .

Now we can use the strong maximum principle (see [9, section 7.1.4, Thm. 11]), which gives that if the inequality (27) is an equality for some  $t > 0$  and  $\omega \in \mathbb{S}$ , then  $\tilde{f}$  is constant on  $[0, t] \times \mathbb{S}$ . So  $f_0$  is the constant function 1.  $\square$

**PROPOSITION 2.8** (global existence, positivity). *Suppose  $f_0$  is a probability measure belonging to  $H^s(\mathbb{S})$  (this is always the case for  $s < -\frac{n-1}{2}$ , according to Sobolev embeddings). Then there exists a global weak solution of (3), which remains a probability measure for any time.*

We remark that the uniqueness of the solution on any time interval remains by Proposition 2.6.

*Proof.* We first prove this proposition in the case  $s > \frac{n+3}{2}$ .

We define a solution by constructing it on a sequence of intervals.

We set  $T_1 = \frac{1}{C_1} \ln(1 + \frac{1}{1+2C_2\|g_0\|_{\dot{H}^s}})$  as in (14). This gives existence to a solution  $g$  in  $C([0, T_1], \dot{H}^s(\mathbb{S}))$ . By induction, we define  $T_{k+1} = T_k + \frac{1}{C_1} \ln(1 + \frac{1}{1+2C_2\|g(T_k)\|_{\dot{H}^s}})$ , which gives existence to a solution  $g \in C([T_k, T_{k+1}], \dot{H}^s(\mathbb{S}))$ .

So we have a solution on  $[0, T]$ , provided that  $T \leq T_k$  for some integer  $k$ .

Now by the previous proposition, this solution  $f = 1 + g$  is nonnegative. We obviously have  $|J[g]| = |J[f]| \leq \int_{\mathbb{S}} |\omega| f = 1$ . Then we can do better estimates, starting from (20):

$$(28) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \|g\|_{\dot{H}^s}^2 + \tau \|g\|_{\dot{H}^{s+1}}^2 &\leq C_0 |J[g]| \|g\|_{\dot{H}^s}^2 + (n-1)^s |J[g]|^2 \\ &\leq C_8 \|g\|_{\dot{H}^s}^2. \end{aligned}$$

Then, Grönwall's lemma gives us that  $\|g(T_k)\|_{\dot{H}^s} \leq \|g_0\|_{\dot{H}^s} e^{C_8 T_k}$ . Suppose now that the sequence  $(T_k)$  is bounded; then  $\|g(T_k)\|_{\dot{H}^s}$  is also bounded. By the definition of  $T_{k+1}$ , the difference  $T_{k+1} - T_k$  does not tend to zero, which implies that the increasing sequence  $(T_k)$  is unbounded, and this is a contradiction. So we have that  $T_k \xrightarrow{k \rightarrow \infty} \infty$ , and the solution is global in time.

Now we do the general case for any  $s$ . Take  $g_0^k$  as a sequence of elements of  $\dot{H}^{\frac{n}{2}+2}$  converging to  $g_0$  in  $\dot{H}^s$  and such that  $f_0^k = 1 + g_0^k$  are positive functions. Let  $g^k$  be the solutions associated with these initial conditions.

Then we have the same estimates as before; since we still have  $|J[g]| \leq 1$ , solving (28) gives

$$\|g^k(t)\|_{\dot{H}^s}^2 + \tau \int_0^t \|g^k(t)\|_{\dot{H}^{s+1}}^2 \leq \|g_0^k\|_{\dot{H}^s} e^{C_s t}.$$

We want to prove that  $g^k$  is a Cauchy sequence, so we study the difference  $u = g^j - g^k$  (in the same way as what was done for  $g - \tilde{g}$  in (22)–(23) to prove uniqueness), which satisfies, for any  $h \in \dot{H}^{-s}(\mathbb{S})$ ,

$$(29) \quad \langle \partial_t u, h \rangle = -\tau \langle \nabla_\omega u, \nabla_\omega h \rangle + (n-1)J[u] \cdot J[h] + \langle u, J[g^j] \cdot \nabla_\omega h \rangle + \langle g^k, J[u] \cdot \nabla_\omega h \rangle.$$

We take  $h = (-\Delta)^s u$  and use the first and second parts of Lemma 2.1 to get

$$(30) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \|u\|_{\dot{H}^s}^2 + \tau \|u\|_{\dot{H}^{s+1}}^2 &\leq C_9 \|u\|_{\dot{H}^s}^2 + C_7 \|u\|_{\dot{H}^s} \|g^k\|_{\dot{H}^{s+1}} \|(-\Delta)^s u\|_{\dot{H}^{-s}} \\ &\leq C_{10} (1 + \|g^k\|_{\dot{H}^{s+1}}) \|u\|_{\dot{H}^s}^2. \end{aligned}$$

If we fix  $T > 0$ , Grönwall’s lemma then gives the following estimate:

$$\begin{aligned} \|u\|_{\dot{H}^s}^2 + \tau \int_0^T \|u\|_{\dot{H}^{s+1}}^2 &\leq \|u_0\|_{\dot{H}^s}^2 \exp\left(C_{10} \int_0^T (1 + \|g^k\|_{\dot{H}^{s+1}})\right) \\ &\leq \|u_0\|_{\dot{H}^s}^2 \exp\left(C_{10}(T + \tau^{-1}\sqrt{T}\|g_0^k\|_{\dot{H}^s} e^{C_s T})\right). \end{aligned}$$

Since  $\|g_0^k\|_{\dot{H}^s}$  is bounded (because  $g_0^k$  converges in  $\dot{H}^s$ ), together with (29), we finally get that  $u$  is bounded in  $L^2((0, T), \dot{H}^{s+1}(\mathbb{S})) \cap H^1((0, T), \dot{H}^{s-1}(\mathbb{S}))$  by a constant  $C_T$  times  $\|u(0)\|_{\dot{H}^s}$ . This gives that  $g^k$  is a Cauchy sequence in that space, and then it converges to a function  $g$ , which is a weak solution of our problem (by Proposition 2.3, we have that  $g(0) = g_0$ ). This is valid for any  $T > 0$ , so this solution is global.

If we take  $\varphi$  in  $C^\infty(\mathbb{S})$ , since  $f^k(t) = 1 + g^k(t)$  is a positive function with mean 1, we have that

$$-\|\varphi\|_\infty = \langle f^k(t), -\|\varphi\|_\infty \rangle \leq \langle f^k(t), \varphi \rangle \leq \langle f^k(t), \|\varphi\|_\infty \rangle = \|\varphi\|_\infty.$$

Passing to the limit gives  $|\langle g(t), \varphi \rangle| \leq \|\varphi\|_\infty$ . Furthermore, we have  $\langle f^k(t), 1 \rangle = 1$  so that  $\langle f(t), 1 \rangle = 1$ , and if  $\varphi$  is a nonnegative function, then  $\langle f^k(t), \varphi \rangle \geq 0$  and we get  $\langle f(t), \varphi \rangle \geq 0$ . This gives that  $f(t)$  is a positive Radon measure with mass 1, which is a probability measure.  $\square$

PROPOSITION 2.9 (instantaneous regularity and boundedness estimates). *If  $f_0$  is a probability measure, then the solution  $f$  belongs to  $C^\infty((0, +\infty) \times \mathbb{S})$  and is positive for any time  $t > 0$ , and we have the following estimates for all  $s \in \mathbb{R}$  and  $m \geq 0$ :*

$$\|f(t)\|_{\dot{H}^{s+m}}^2 \leq C \left(1 + \frac{1}{t^m}\right) \|f_0\|_{\dot{H}^s}^2,$$

where the constant  $C$  depends only on  $\tau, s$ , and  $m$ .

In particular we have that for  $t_0 > 0$ ,  $f$  is uniformly bounded on  $[t_0, +\infty)$  in any  $H^s$  norm.

*Proof.* Suppose  $f_0 \in H^s(\mathbb{S})$ , and fix  $t > 0$ . The solution  $f$  is in  $C([0, +\infty), H^s(\mathbb{S}))$  and in  $L^2((0, t), H^{s+1}(\mathbb{S}))$ . Then there exists  $s < t$  such that  $f(s) \in H^{s+1}(\mathbb{S})$ . So we can construct a solution belonging to  $C([s, +\infty), H^{s+1}(\mathbb{S}))$ . But this solution is also a weak solution in  $L^2((s, T), H^{s+1}(\mathbb{S})) \cap H^{-1}((s, T), H^{s-1}(\mathbb{S}))$  for all  $T > s$ , so by uniqueness it is equal to  $f$ . Then  $f$  belongs to  $C([t, +\infty), H^{s+1}(\mathbb{S}))$ . Since this is true for all  $t > 0$ , then  $f$  belongs to  $C((0, +\infty), H^{s+1}(\mathbb{S}))$ . We can repeat this argument and have that  $f$  belongs to  $C((0, +\infty), H^p(\mathbb{S}))$  for any  $p$ , and is a positive classical solution, by Proposition 2.7. Using the equation, differentiating in time gives that it is also in  $C^k((0, +\infty), H^p(\mathbb{S}))$  for any  $p$  and any  $k$ , so by Sobolev embeddings it is a  $C^\infty$  function of  $(0, +\infty) \times \mathbb{S}$ .

Since we have positivity, we can have estimates for any of the modes of  $f = 1 + g$ . Let us denote by  $f^N$  the orthogonal projection of  $f$  on the  $N$  first eigenspaces of the Laplacian and by  $g^N = f - f^N$  the projection on the other ones (high modes).

We have a Poincaré inequality on this space:  $\|g^N\|_{\dot{H}^s}^2 \leq \frac{1}{(N+1)(N+n-1)} \|g^N\|_{\dot{H}^{s+1}}^2$  (we recall that the eigenvalues of  $-\Delta$  are given by  $\ell(\ell+n-2)$  for  $\ell \in \mathbb{N}$ ). We use the estimate (20):

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|g\|_{\dot{H}^s}^2 + \tau \|g\|_{\dot{H}^{s+1}}^2 &\leq C_0 |J[g]| \|g\|_{\dot{H}^s}^2 + (n-1)^s |J[g]|^2 \\ (31) \qquad \qquad \qquad &\leq \frac{C_0}{(N+1)(N+n-1)} \|g\|_{\dot{H}^{s+1}}^2 + (n-1)^s |J[g]|^2 + C_0 \|f^N - 1\|_{\dot{H}^s}^2. \end{aligned}$$

Now we have, since  $f$  is a probability measure, that

$$\|f^N - 1\|_{\dot{H}^s}^2 = \int_{\mathbb{S}} (-\Delta)^s f^N f d\omega \leq \|(-\Delta)^s f^N\|_{L^\infty} \leq K_N \|f^N - 1\|_{\dot{H}^s},$$

the last inequality being the equivalence between norms in finite dimension. Dividing by this last norm, this gives that the low modes of  $f$  are uniformly bounded in time by a constant  $K_N$ . Then we have, taking  $N$  sufficiently large,

$$\frac{1}{2} \frac{d}{dt} \|g\|_{\dot{H}^s}^2 + \frac{\tau}{2} \|g\|_{\dot{H}^{s+1}}^2 \leq C_{11}.$$

Now multiplying this formula by  $t$  at order  $s + 1$ , we get

$$\frac{1}{2} \frac{d}{dt} (t \|g\|_{\dot{H}^{s+1}}^2) + \frac{\tau}{2} t \|g\|_{\dot{H}^{s+2}}^2 \leq C_{12} t + \frac{1}{2} \|g\|_{\dot{H}^{s+1}}^2,$$

and finally

$$\frac{1}{2} \frac{d}{dt} \left( \|g\|_{\dot{H}^s}^2 + \frac{\tau}{2} t \|g\|_{\dot{H}^{s+1}}^2 \right) + \frac{\tau}{4} \left( \|g\|_{\dot{H}^{s+1}}^2 + \frac{\tau}{2} t \|g\|_{\dot{H}^{s+2}}^2 \right) \leq C_{11} + C_{12} \frac{\tau}{2} t.$$

Together with the Poincaré inequality, solving this inequality gives us

$$\|g\|_{\dot{H}^s}^2 + \frac{\tau}{2} t \|g\|_{\dot{H}^{s+1}}^2 \leq \|g_0\|_{\dot{H}^s}^2 e^{-(n-1)\frac{\tau}{4}t} + C_{13}(1+t).$$

So we have the result for  $\|f\|_{\dot{H}^s}^2 = 1 + \|g\|_{\dot{H}^s}^2$ , and  $m = 1$ :

$$\|f(t)\|_{\dot{H}^{s+1}}^2 \leq C \left( 1 + \frac{1}{t} \right) \|f_0\|_{\dot{H}^s}^2.$$

Then we apply this inequality between 0 and  $\frac{t}{2}$  and the inequality at order  $m$  between  $\frac{t}{2}$  and  $t$  to get the result at order  $m + 1$ . The case where  $m$  is any nonnegative real also works by interpolation.  $\square$

This last proposition ends the proof of Theorem 2.4. Let us here make two small comments concerning the analyticity of the solution and the limit case with no noise:  $\tau = 0$ .

*Remark 2.10* (analyticity of the solution). We can show, as claimed in [4], [5], that at any time  $t > 0$  the solution is analytic in the space variable. The idea is to show, following [5] (based on [3], [11]), that the solution is in some Gevrey class of functions, defined by a parameter depending on time. This class is a subset of the set of real analytic functions on the sphere. More details and a complete proof are given in section A.2. We could have dealt with these classes of functions instead of working in the Sobolev spaces directly, but we will not need these properties of analyticity in the following. In any case, to prove analyticity we need the initial condition to be in  $H^{-\frac{n-1}{2}}(\mathbb{S})$ , so this study of instantaneous regularization was necessary.

*Remark 2.11* (case where  $\tau = 0$ : no noise). The proof is also valid, except that the solution belongs to  $L^\infty((0, T), H^s(\mathbb{S})) \cap H^1((0, T), H^{s-1}(\mathbb{S}))$  if the initial condition is in  $H^s(\mathbb{S})$ . By an optimal regularity argument, we can get that a solution is in fact in  $C([0, T], H^s(\mathbb{S}))$ . The nonnegativity argument is then also valid, and so the solution is global. Obviously, we do not have the instantaneous regularity and boundedness estimates.

**3. Using the free energy.** In this section, we derive the Onsager free energy (6) for Smoluchowski equation (3) and use it to get general results on the steady states.

**3.1. Free energy and steady states.** We rewrite (3) as

$$\partial_t f = Q(f) = \nabla_\omega \cdot (\tau \nabla_\omega f - \nabla_\omega(\omega \cdot J[f])f) = \nabla_\omega \cdot (f \nabla_\omega(\tau \ln f - \omega \cdot J[f])).$$

Since any solution is in  $C^\infty((0, +\infty) \times \mathbb{S})$  and positive for any  $t > 0$ , there is no problem with using  $\ln f$  and doing any integration by parts. We multiply the equation by  $\tau \ln f - \omega \cdot J[f]$  and integrate by parts, and we get

$$\int_{\mathbb{S}} \partial_t f (\tau \ln f - \omega \cdot J[f]) d\omega = - \int_{\mathbb{S}} f |\nabla_\omega(\tau \ln f - \omega \cdot J[f])|^2 d\omega.$$

Since the left part can be recast as a time derivative, this is a conservation relation. We define the free energy  $\mathcal{F}(f)$  and the dissipation term  $\mathcal{D}(f)$  by

$$(32) \quad \mathcal{F}(f) = \tau \int_{\mathbb{S}} f \ln f - \frac{1}{2} |J[f]|^2,$$

$$(33) \quad \mathcal{D}(f) = \int_{\mathbb{S}} f |\nabla_\omega(\tau \ln f - \omega \cdot J[f])|^2,$$

and we have the following energy dissipation relation:

$$(34) \quad \frac{d}{dt} \mathcal{F} + \mathcal{D} = 0.$$

We define a steady state as a (weak) solution which does not depend on time. Here are some characterizations of the steady states.

**PROPOSITION 3.1** (steady states). *The steady states of Smoluchowski equation (3) are the probability measures  $f$  on  $\mathbb{S}$  which satisfy one of the following equivalent conditions:*

1. *Equilibrium:*  $f \in C^2(\mathbb{S})$  and  $Q(f) = 0$ .
2. *No dissipation:*  $f \in C^1(\mathbb{S})$  and  $\mathcal{D}(f) = 0$ .
3. *The probability density  $f \in C^0(\mathbb{S})$  is positive and a critical point of  $\mathcal{F}$  (under the constraint of mean 1).*
4. *There exists  $C \in \mathbb{R}$  such that  $\tau \ln f - J[f] \cdot \omega = C$ .*

*Proof.* By definition, a steady state  $f$  is a solution independent of  $t$ . Since it is a solution, it is positive and  $C^\infty$ , and we get that  $Q(f) = 0$ . By the conservation relation (34), we get that  $\frac{d}{dt}\mathcal{F} = 0$ , so  $\mathcal{D}(f) = 0$ . Since it is positive, we get that  $\nabla_\omega(\tau \ln f - \omega \cdot J[f]) = 0$ , so there exists  $C \in \mathbb{R}$  such that  $\tau \ln f - J[f] \cdot \omega = C$ .

Now we do a variational study of  $\mathcal{F}$  around  $f$ . We take a small perturbation  $f + h$  of  $f$  which remains a probability density function (which means that  $\int_{\mathbb{S}} h = 0$ ).

We can expand the function  $x \mapsto x \ln x$  around  $f$ , since  $f \geq \varepsilon > 0$ , and we have

$$\begin{aligned} \mathcal{F}(f + h) &= \tau \int_{\mathbb{S}} (f \ln f + h \ln f + h) - \frac{1}{2} |J[f]|^2 - J[f] \cdot \int_{\mathbb{S}} \omega h + O(\|h\|_\infty^2) \\ &= \mathcal{F}(f) + \int_{\mathbb{S}} h(\tau \ln f - J[f] \cdot \omega) + O(\|h\|_\infty^2) \\ &= \mathcal{F}(f) + O(\|h\|_\infty^2), \end{aligned}$$

which means that  $f$  is a critical point of  $\mathcal{F}$ . So  $f$  satisfies the four conditions.

Conversely, if  $f \in C^2(\mathbb{S})$  and  $Q(f) = 0$ , then  $f$  is obviously a steady state.

If  $\tau \ln f - J[f] \cdot \omega = C$ , then  $f \in C^2(\mathbb{S})$  and  $Q(f) = 0$ . We will show that the second and third conditions reduce to this fourth condition.

Doing the above computation around a positive  $f \in C^0(\mathbb{S})$  gives that if  $f$  is a critical point for the free energy, then  $\int_{\mathbb{S}} h(\tau \ln f - J[f] \cdot \omega)$  is zero for any  $h$  with mean zero. This is exactly saying that  $\tau \ln f - J[f] \cdot \omega$  is constant.

Finally, if we suppose  $f \in C^1(\mathbb{S})$  and  $\mathcal{D}(f) = 0$ , at any point  $\omega_0 \in \mathbb{S}$  such that  $f(\omega_0) > 0$  we have that  $\nabla(\tau \ln f - J[f] \cdot \omega) = 0$  on a neighborhood of  $\omega_0$ .

The function  $\varphi$  defined by  $\varphi(\omega) = \tau \ln f - J[f] \cdot \omega$  is then locally constant at any point where it is finite, so  $\varphi^{-1}(\{C\})$  is open in  $\mathbb{S}$  for any  $C \in \mathbb{R}$ .

Now if  $\varphi(\omega_k) = C$ , with  $\omega_k$  converging to  $\omega_\infty$ , then  $f(\omega_k) = \exp(\frac{C + J[f] \cdot \omega_k}{\tau})$ . Passing to the limit, we get that  $f(\omega_\infty) = \exp(\frac{C + J[f] \cdot \omega_\infty}{\tau})$ , which gives  $\varphi(\omega_\infty) = C$ . So  $\varphi^{-1}(\{C\})$  is closed.

Since  $f$  is not identically zero, there exists  $C \in \mathbb{R}$  such that  $\varphi^{-1}(\{C\}) \neq \emptyset$ , and by connectedness of the sphere, we get  $\varphi^{-1}(\{C\}) = \mathbb{S}$ , so  $\tau \ln f - J[f] \cdot \omega = C$ .  $\square$

**3.2. LaSalle principle.** We give here an adaptation of LaSalle’s invariance principle to our PDE framework.

**PROPOSITION 3.2 (LaSalle’s invariance principle).** *Let  $f_0$  be a probability measure on the sphere  $\mathbb{S}$ . We denote by  $\mathcal{F}_\infty$  the limit of  $\mathcal{F}(f(t))$  as  $t \rightarrow \infty$ , where  $f$  is the solution to Smoluchowski equation (3) with initial condition  $f_0$ .*

*Then the set  $\mathcal{E}_\infty = \{f \in C^\infty(\mathbb{S}) \text{ such that } \mathcal{D}(f) = 0 \text{ and } \mathcal{F}(f) = \mathcal{F}_\infty\}$  is not empty.*

*Furthermore,  $f(t)$  converges in any  $H^s$  norm to this set of equilibria in the following sense:*

$$\lim_{t \rightarrow \infty} \inf_{g \in \mathcal{E}_\infty} \|f(t) - g\|_{H^s} = 0.$$

*Proof.* First of all,  $\mathcal{F}(f(t))$  is decreasing in time and bounded below by  $-\frac{1}{2}$ , so  $\mathcal{F}_\infty$  is well defined.

Let  $(t_n)$  be an unbounded increasing sequence, and suppose that  $f(t_n)$  converges in  $H^s(\mathbb{S})$  to  $f_\infty$  for some  $s \in \mathbb{R}$ . We first remark that  $f(t_n)$  is uniformly bounded in  $H^{s+2p}(\mathbb{S})$  (using Theorem 2.4), and then by a simple interpolation estimate we get that  $\|f(t_n) - f(t_m)\|_{\dot{H}^{s+p}}^2 \leq \|f(t_n) - f(t_m)\|_{\dot{H}^s} \|f(t_n) - f(t_m)\|_{\dot{H}^{s+2p}}$ , and  $f(t_n)$  also converges in  $H^{s+p}(\mathbb{S})$ . So  $f_\infty$  is in any  $H^s(\mathbb{S})$ .

We want to prove that  $\mathcal{D}(f_\infty) = 0$ . Supposing this is not the case, we write

$$\begin{aligned} \mathcal{D}(f) &= \tau^2 \int_{\mathbb{S}} \frac{|\nabla_\omega f|^2}{f} + J[f] \cdot \int_{\mathbb{S}} (\text{Id} - \omega \otimes \omega) f J[f] - 2\tau J[f] \cdot \int_{\mathbb{S}} \nabla_\omega f \\ (35) \quad &= \tau^2 \int_{\mathbb{S}} \frac{|\nabla_\omega f|^2}{f} + (1 - 2(n - 1)\tau) |J[f]|^2 - \int_{\mathbb{S}} (\omega \cdot J[f])^2 f. \end{aligned}$$

Now we take  $s$  sufficiently large such that  $H^s(\mathbb{S}) \subset L_\infty(\mathbb{S}) \cap H^1(\mathbb{S})$ . If  $f_\infty$  is positive, then  $\mathcal{D}$ , as a function from the nonnegative elements of  $H^s(\mathbb{S})$  to  $[0, +\infty]$ , is continuous at the point  $f_\infty$ . In particular since  $\mathcal{D}(f_\infty) > 0$ , there exist  $\delta > 0$  and  $M > 0$  such that if  $\|f - f_\infty\|_{H^s} \leq \delta$ , then we have  $\mathcal{D}(f) \geq M$ . We want to show the same result in the case where  $f_\infty$  is only nonnegative. We define

$$\mathcal{D}_\varepsilon(f) = \tau^2 \int_{\mathbb{S}} \frac{|\nabla_\omega f|^2}{f + \varepsilon} + (1 - 2(n - 1)\tau) |J[f]|^2 - \int_{\mathbb{S}} (\omega \cdot J[f])^2 f.$$

We have by monotone convergence that  $\mathcal{D}_\varepsilon(f_\infty)$  converges to  $\mathcal{D}(f_\infty)$  as  $\varepsilon \rightarrow 0$ . So there exists  $\varepsilon > 0$  such that  $\mathcal{D}_\varepsilon(f_\infty) > 0$ . Now by continuity of  $\mathcal{D}_\varepsilon$  at the point  $f_\infty$ , we get that there exist  $\delta > 0$  and  $M > 0$  such that if  $\|f - f_\infty\|_{H^s} \leq \delta$ , then  $\mathcal{D}_\varepsilon(f) \geq M$ . The fact that  $\mathcal{D}(f) \geq \mathcal{D}_\varepsilon(f)$  gives the same result as before.

Now since  $\partial_t f$  is uniformly bounded in  $H^s$  (for  $t \geq t_1 > 0$ ), there exists  $\eta > 0$  such that if  $|t - t'| \leq \eta$ , then  $\|f(t) - f(t')\|_{H^s} \leq \frac{\delta}{2}$ . We then take  $N$  sufficiently large such that  $\|f(t_n) - f_\infty\|_{H^s} \leq \frac{\delta}{2}$  for all  $n \geq N$ .

Then we have that for  $n \geq N$ ,  $\mathcal{D}(f) \geq M$  on  $[t_n, t_n + \eta]$ . Up to extracting, we can assume that  $t_{n+1} \geq t_n + \eta$ , so we have

$$\mathcal{F}(f(t_N)) - \mathcal{F}(f(t_{N+p})) = \int_{t_N}^{t_{N+p}} \mathcal{D}(f) \geq p\eta M.$$

Since the left term is bounded by  $\mathcal{F}(f(t_N)) - \mathcal{F}_\infty$ , taking  $p$  sufficiently large gives the contradiction.

Now if we suppose that for a given  $s$  the distance (in  $H^s$  norm) between  $f(t)$  and  $\mathcal{E}_\infty$  does not tend to 0, we get  $\varepsilon > 0$  and a sequence  $t_n$  such that for all  $g \in \mathcal{E}_\infty$ , we have  $\|f(t_n) - g\|_{H^s} \geq \varepsilon$ . Since  $f(t_n)$  is bounded in  $H^{s+1}(\mathbb{S})$ , by a compact Sobolev embedding, up to extracting we can assume that  $f(t_n)$  is converging in  $H^s(\mathbb{S})$  to  $f_\infty$ . By the previous argument,  $f \in C^\infty(\mathbb{S})$  and we have  $\mathcal{D}(f_\infty) = 0$ . Obviously, since  $\mathcal{F}(f)$  is decreasing in time, we have that  $\mathcal{F}(f_\infty) = \mathcal{F}_\infty$ . So  $f_\infty$  belongs to  $\mathcal{E}_\infty$ , and then  $\|f(t_n) - f_\infty\|_{H^s} \geq \varepsilon$  for all  $n$ . This is a contradiction.

Since the distance between  $f(t)$  and  $\mathcal{E}_\infty$  tends to 0, obviously this set is not empty.  $\square$

**3.3. Computation of equilibria.** Define, for a unit vector  $\Omega \in \mathbb{S}$  and  $\kappa \geq 0$ , the Fisher–von Mises distribution with concentration parameter  $\kappa$  and orientation  $\Omega$  by

$$(36) \quad M_{\kappa\Omega}(\omega) = \frac{\exp(\kappa \omega \cdot \Omega)}{\int_{\mathbb{S}} \exp(\kappa v \cdot \Omega) dv}.$$



Note that the denominator depends only on  $\kappa$ . We have that the density of  $M_{\kappa\Omega}$  is 1, and the flux is

$$(37) \quad J[M_{\kappa\Omega}] = \frac{\int_{\mathbb{S}} \omega \exp(\kappa \omega \cdot \Omega) d\omega}{\int_{\mathbb{S}} \exp(\kappa \omega \cdot \Omega) d\omega} = c(\kappa)\Omega,$$

where

$$(38) \quad c(\kappa) = \frac{\int_0^\pi \cos \theta e^{\kappa \cos \theta} \sin^{n-2} \theta d\theta}{\int_0^\pi e^{\kappa \cos \theta} \sin^{n-2} \theta d\theta}.$$

If  $f$  is an equilibrium,  $\tau \ln f - J[f] \cdot \omega$  is constant, and then  $f = C \exp(\tau^{-1} J[f] \cdot \omega)$ . Since  $f$  is a probability density function, we get  $f = M_{\kappa\Omega}$  with  $\kappa\Omega = \tau^{-1} J[f]$  (in the case where  $|J[f]| = 0$ , then  $\kappa = 0$  and we can take any  $\Omega$ ; this is just the uniform distribution). Finally, with (37) we get  $J[f] = c(\kappa)\Omega$ , which gives the following compatibility condition:

$$(39) \quad c(\kappa) = \tau\kappa.$$

We give the solutions of this equation in a proposition.

PROPOSITION 3.3 (compatibility condition).

- If  $\tau \geq \frac{1}{n}$ , there is only one solution to the compatibility condition:  $\kappa = 0$ . The only equilibrium is the constant function  $f = 1$ .
- If  $\tau < \frac{1}{n}$ , the compatibility condition has exactly two solutions:  $\kappa = 0$ , and one unique positive solution that we will denote  $\kappa(\tau)$ . Apart from the constant function  $f = 1$  (the case  $\kappa = 0$ ), the equilibria form a manifold of dimension  $n - 1$ : the functions of the form  $f = M_{\kappa(\tau)\Omega}$ , where  $\Omega \in \mathbb{S}$  is an arbitrary unit vector.

*Proof.* Let us denote  $\tilde{\tau}(\kappa) = \frac{c(\kappa)}{\kappa}$ . A simple Taylor expansion gives  $\tilde{\tau}(\kappa) \xrightarrow{\kappa \rightarrow 0} \frac{1}{n}$ . Since the function  $\tilde{\tau}$  tends to 0 as  $\kappa \rightarrow +\infty$  (because  $c(\kappa) \leq 1$ ), it is sufficient to prove that it is decreasing. Indeed, the function is then a one-to-one correspondence from  $\mathbb{R}_+^*$  to  $(0, \frac{1}{n})$ , and the compatibility condition for  $\kappa > 0$  is exactly solving  $\tau = \tilde{\tau}(\kappa)$ .

But we have (after one integration by parts) that  $\tilde{\tau}'(\kappa) = \frac{1}{\kappa}(1 - n\tilde{\tau}(\kappa) - c(\kappa)^2)$ , which, by the following lemma, is negative for  $\kappa > 0$ .  $\square$

LEMMA 3.4. Define  $\beta = c(\kappa)^2 + n\tilde{\tau}(\kappa) - 1$ . Then for any  $\kappa > 0$ , we have  $\beta > 0$ .

*Proof.* Define  $[\gamma(\cos \theta)]_\kappa = \int_0^\pi \gamma(\cos \theta) e^{\kappa \cos \theta} \sin^{n-2} \theta d\theta$ .

Then we have by definition  $\beta = \frac{\kappa[\cos \theta]_\kappa^2 + n[\cos \theta]_\kappa [1]_\kappa - \kappa [1]_\kappa^2}{\kappa [1]_\kappa^2}$ . So we need only show that the numerator is positive. We will prove in fact that the Taylor expansion of this term in  $\kappa$  has only positive terms.

We have, if we denote  $a_p = \frac{1}{(2p)!} \int_0^\pi \cos^{2p} \theta \sin^{n-2} \theta d\theta \geq 0$ ,

$$[1]_\kappa = \sum_{p=0}^\infty a_p \kappa^{2p} \quad \text{and} \quad [\cos \theta]_\kappa = \sum_{p=0}^\infty (2p+2) a_{p+1} \kappa^{2p+1}.$$

Now doing an integration by parts in the definition of  $a_{p+1}$ , we get

$$a_{p+1} = \frac{2p+1}{n-1} \left( \frac{a_p}{(2p+1)(2p+2)} - a_{p+1} \right),$$

which gives

$$(40) \quad (2p+2) a_{p+1} = \frac{a_p}{2p+n}.$$

We have, for  $\kappa > 0$ ,

$$\begin{aligned} \beta \kappa [1]_{\kappa}^2 &= \sum_{k=0}^{\infty} \left( \sum_{p+q=k-1} (2p+2) a_{p+1} (2q+2) a_{q+1} + \sum_{p+q=k} n(2p+2) a_{p+1} a_q - a_p a_q \right) \kappa^{2k+1} \\ &= \sum_{k=0}^{\infty} \left( \sum_{p+q=k, p \geq 1} 2p a_p \frac{1}{2q+n} a_q + \sum_{p+q=k} \left( \frac{n}{2p+n} - 1 \right) a_p a_q \right) \kappa^{2k+1} \\ &= \sum_{k=0}^{\infty} \left( \sum_{p+q=k} 2p \left( \frac{1}{2q+n} - \frac{1}{2p+n} \right) a_p a_q \right) \kappa^{2k+1} \\ &= \sum_{k=0}^{\infty} \left( \sum_{p+q=k} \left( p \left( \frac{1}{2q+n} - \frac{1}{2p+n} \right) + q \left( \frac{1}{2p+n} - \frac{1}{2q+n} \right) \right) a_p a_q \right) \kappa^{2k+1} \\ &= \sum_{k=0}^{\infty} \left( \sum_{p+q=k} \frac{2(p-q)^2}{(2p+n)(2q+n)} a_p a_q \right) \kappa^{2k+1}. \end{aligned}$$

So we finally get

$$\beta = \left( \sum_{p=0}^{\infty} a_p \kappa^{2p} \right)^{-1} \sum_{k=0}^{\infty} \left( \sum_{p+q=k} \frac{2(p-q)^2}{(2p+n)(2q+n)} a_p a_q \right) \kappa^{2k},$$

which gives that  $\beta > 0$  when  $\kappa > 0$ .  $\square$

*Remark 3.5.* We can do another proof, following an argument of [25], which does not need to compute  $\beta$  explicitly.

The idea is that we compute  $\tilde{\tau}'' = (n-1) \frac{\beta}{\kappa^2} - 2\tilde{\tau}(\tilde{\tau} - \beta)$ , so we see (except in the case  $\kappa = 0$ ) that if  $\tilde{\tau}' = -\frac{\beta}{\kappa} = 0$ , then  $\tilde{\tau}'' < 0$  (indeed, we will easily see in (46) that  $\tilde{\tau} - \beta$  is positive). For the case  $\kappa = 0$ , we can compute the Taylor expansion of  $\tilde{\tau}$  up to order 2:  $\tilde{\tau}(\kappa) = \frac{1}{n} - \frac{1}{n^2(n+2)} \kappa^2 + O(\kappa^4)$ . So we have that any critical point of  $\tilde{\tau}$  is a maximum. Since there is a local maximum at  $\kappa = 0$ , the function is decreasing.

We can have an asymptotic expansion of the order parameter  $c(\kappa(\tau))$  as  $\tau$  reaches the critical value  $\frac{1}{n}$ . Indeed, we have that  $\tau - \frac{1}{n} \sim -\frac{1}{n^2(n+2)} \kappa(\tau)^2$  by the expansion of  $\tilde{\tau}$  in the previous remark. So

$$(41) \quad c(\kappa(\tau)) \sim \frac{1}{n} \kappa(\tau) \sim \sqrt{(n+2) \left( \frac{1}{n} - \tau \right)} \text{ as } \tau \rightarrow \frac{1}{n}.$$

**PROPOSITION 3.6** (minimum of the free energy).

- If  $\tau \geq \frac{1}{n}$ , the minimum of the free energy is 0, reached only by the uniform distribution. Any solution converges to the uniform distribution in any  $H^s$  norm.
- If  $\tau < \frac{1}{n}$ , the minimum of the free energy is negative, reached only by any nonisotropic equilibrium  $M_{\kappa(\tau)\Omega}$ .

*Proof.* By the LaSalle principle (Proposition 3.2), we have that

$$\min_{f \in C^\infty(\mathbb{S}), f > 0} \mathcal{F}(f) = \min_{f \in C^\infty(\mathbb{S}), f > 0, \mathcal{D}(f)=0} \mathcal{F}(f).$$

Indeed, for any positive initial condition  $f$  in  $C^\infty(\mathbb{S})$ , there exists an equilibrium  $f_\infty$  such that  $\mathcal{F}(f_\infty) = \mathcal{F}_\infty \leq \mathcal{F}(f)$ . This gives

$$\inf_{f \in C^\infty(\mathbb{S}), f > 0} \mathcal{F}(f) = \inf_{f \in C^\infty(\mathbb{S}), f > 0, \mathcal{D}(f)=0} \mathcal{F}(f).$$

Since the set of equilibria is compact (either a single point, or one point and a manifold homeomorphic to  $\mathbb{S}$ ), this infimum is a minimum.

Furthermore, if  $f_0$  is not an equilibrium, then  $\mathcal{D}(f_0) > 0$ , and then  $\mathcal{F}(f(t))$  is decreasing in the neighborhood of  $t = 0$ . So the minimum of  $\mathcal{F}$  cannot be reached for  $f_0$ .

In the case  $\tau \geq \frac{1}{n}$ , this gives the result since the only equilibrium is the constant function 1. By the LaSalle principle, we also get that the solution is converging to the constant solution in any  $H^s$  norm.

In the case  $\tau < \frac{1}{n}$ , we have that  $\mathcal{F}(1 + \varepsilon\omega \cdot \Omega) \sim \frac{1}{n}(\tau - \frac{1}{n})\varepsilon^2$  for a fixed unit vector  $\Omega \in \mathbb{S}$ , so there exists  $f_0$  such that  $\mathcal{F}(f_0) < 0$ . Then the uniform distribution cannot be a global minimizer. Since  $\mathcal{F}(M_{\kappa(\tau)\Omega})$  is independent of  $\Omega$ , we get that this value is the minimum.  $\square$

**4. Convergence to equilibrium.** In this section, we establish and study the convergence of the solution to an equilibrium for any initial condition in the three different regimes, depending on whether  $\tau$  is greater, less than, or equal to  $\frac{1}{n}$ .

**4.1. A new entropy, application to the subcritical case  $\tau > \frac{1}{n}$ .** In this section we derive a convex entropy, which shows global decay to the uniform distribution in the case  $\tau > \frac{1}{n}$ .

On  $\dot{H}^{-\frac{n-1}{2}}(\mathbb{S})$  we define the norm  $\|\cdot\|_{\tilde{H}^{-\frac{n-1}{2}}}$  by  $\|g\|_{\tilde{H}^{-\frac{n-1}{2}}}^2 = \int_{\mathbb{S}} g \tilde{\Delta}_{n-1}^{-1} g$ , where the conformal Laplacian  $\tilde{\Delta}_{n-1}$  is defined by (7). This norm is equivalent to  $\|\cdot\|_{\dot{H}^{-\frac{n-1}{2}}}$ . We also define  $\|\cdot\|_{\tilde{H}^{-\frac{n-3}{2}}}$  by  $\|g\|_{\tilde{H}^{-\frac{n-3}{2}}}^2 = \int_{\mathbb{S}} \Delta g \tilde{\Delta}_{n-1}^{-1} g$ , and this norm is equivalent to the  $\|\cdot\|_{\dot{H}^{-\frac{n-3}{2}}}$  norm.

Taking  $h = \tilde{\Delta}_{n-1}^{-1} g$  in the weak formulation (13) and using the last part of Lemma 2.1, we obtain a conservation relation:

$$(42) \quad \frac{1}{2} \frac{d}{dt} \|g\|_{\tilde{H}^{-\frac{n-1}{2}}}^2 = -\tau \|g\|_{\tilde{H}^{-\frac{n-3}{2}}}^2 + \frac{1}{(n-2)!} |J[g]|^2.$$

We remark that this is a conservation law between quadratic quantities, as would be the case for a linear equation.

Since the component of  $g$  on the space of spherical harmonics of degree 1 is given by  $n\omega \cdot J[g]$ , a simple computation shows that this component's contribution to  $\|g\|_{\tilde{H}^{-\frac{n-1}{2}}}^2$  is equal to  $\frac{n}{(n-1)!} |J[g]|^2$ . Then the last term of the conservation relation (42) is bounded by  $\frac{n-1}{n} \|g\|_{\tilde{H}^{-\frac{n-1}{2}}}^2$ . Together with the Poincaré inequality  $\|g\|_{\tilde{H}^{-\frac{n-3}{2}}}^2 \geq (n-1) \|g\|_{\tilde{H}^{-\frac{n-1}{2}}}^2$ , we get the following estimate:

$$\frac{1}{2} \frac{d}{dt} \|g\|_{\tilde{H}^{-\frac{n-1}{2}}}^2 \leq (n-1) \left( \frac{1}{n} - \tau \right) \|g\|_{\tilde{H}^{-\frac{n-1}{2}}}^2.$$

This gives in the case  $\tau > \frac{1}{n}$  an exponential decay of rate  $(n-1)(\tau - \frac{1}{n})$  for the norm  $\|\cdot\|_{\tilde{H}^{-\frac{n-1}{2}}}$ :

$$\|g\|_{\tilde{H}^{-\frac{n-1}{2}}} \leq \|g_0\|_{\tilde{H}^{-\frac{n-1}{2}}} \exp(-(n-1)(\tau - \frac{1}{n})t).$$

In the general case, if  $f_0 \in H^s(\mathbb{S})$  with  $s > -\frac{n-1}{2}$ , we use the estimate (31):

$$\frac{1}{2} \frac{d}{dt} \|g\|_{H^s}^2 + \tau \|g\|_{H^{s+1}}^2 \leq \frac{C_0}{(N+1)(N+n-1)} \|g\|_{H^{s+1}}^2 + (n-1)^s |J[g]|^2 + C_0 \|f^{N-1}\|_{H^s}^2.$$

Now we have, since  $f$  is a probability measure,

$$(n-1)^s |J[g]|^2 + C_0 \|f^N - 1\|_{\dot{H}^s}^2 \leq K_N \|f^N - 1\|_{\dot{H}^{-\frac{n-1}{2}}}^2 \leq K_N \|g_0\|_{\dot{H}^{-\frac{n-1}{2}}}^2 e^{-2(n-1)(\tau-\frac{1}{n})t},$$

the first inequality being the equivalence between norms in finite dimension. For any  $\varepsilon < \frac{1}{n}$ , taking  $N$  sufficiently large, together with the Poincaré inequality we get

$$\frac{1}{2} \frac{d}{dt} \|g\|_{\dot{H}^s}^2 + (n-1)(\tau - \varepsilon) \|g\|_{\dot{H}^s}^2 \leq C \|g_0\|_{\dot{H}^{-\frac{n-1}{2}}}^2 e^{-2(n-1)(\tau-\frac{1}{n})t},$$

where the constant  $C$  depends only on  $s$ .

Solving this equation, we get

$$\|g\|_{\dot{H}^s}^2 \leq \|g_0\|_{\dot{H}^s}^2 e^{-2(n-1)(\tau-\varepsilon)t} + \frac{C}{(n-1)(\frac{1}{n}-\varepsilon)} \|g_0\|_{\dot{H}^{-\frac{n-1}{2}}}^2 e^{-2(n-1)(\tau-\frac{1}{n})t}.$$

Taking, for example,  $\varepsilon = \frac{1}{2n}$ , since  $s > -\frac{n-1}{2}$ , we get

$$\|g\|_{\dot{H}^s}^2 \leq (1 + 2\tilde{C}\frac{n}{n-1}) \|g_0\|_{\dot{H}^s}^2 e^{-2(n-1)(\tau-\frac{1}{n})t}.$$

In summary, we have the following theorem.

**THEOREM 4.1** (new entropy). *For a given probability density function  $f$ , we define the quantities  $\mathcal{H}(f) = \|f - 1\|_{\dot{H}^{-\frac{n-1}{2}}}^2$  and  $\tilde{\mathcal{D}}(f) = 2\tau \|f - 1\|_{\dot{H}^{-\frac{n-3}{2}}}^2 - \frac{2}{(n-2)!} |J[f]|^2$ .*

*We have the following conservation relation for any solution  $f$  of Smoluchowski equation (3):*

$$(43) \quad \frac{d}{dt} \mathcal{H}(f) + \tilde{\mathcal{D}}(f) = 0.$$

*When  $\tau \geq \frac{1}{n}$ , the term  $\tilde{\mathcal{D}}(f)$  is nonnegative, so the new entropy  $\mathcal{H}(f)$  is decreasing in time.*

*Furthermore, if  $\tau > \frac{1}{n}$ , then in any Sobolev space  $H^s(\mathbb{S})$  with  $s \geq -\frac{n-1}{2}$ , we have global exponential decay of the solution to the uniform distribution, with the rate given by  $(n-1)(\tau - \frac{1}{n})$ .*

*More precisely, there is a constant  $C$  depending only on  $s$  such that for any initial condition  $f_0 \in H^s(\mathbb{S})$ , we have*

$$\|f - 1\|_{H^s} \leq C \|f_0 - 1\|_{H^s} e^{-(n-1)(\tau-\frac{1}{n})t}.$$

Let us make a small remark here. Actually this conservation relation is true for any solution, without any positivity condition. We need only that the mean of  $f$  be 1. Since we have existence and uniqueness in small time for any initial condition, with the same instantaneous regularity results (valid only for a short time existence), we get that the solution belongs to  $H^{-\frac{n-1}{2}}(\mathbb{S})$  at some time. But the conservation relation then gives that we have a global solution. So we can state a stronger theorem of existence and uniqueness.

**THEOREM 4.2.** *Given an initial condition  $f_0$  in  $H^s(\mathbb{S})$  (not necessarily nonnegative), there exists a unique weak solution  $f$  of (3) such that  $f(0) = f_0$ . This solution is global in time (Definition 2.2 is valid for any time  $T > 0$ ). Moreover,  $f$  is a classical solution belonging to  $C^\infty((0, +\infty) \times \mathbb{S})$  (and even analytic in space; see section A.2).*

**Remark 4.3.** In this case, we do not have any uniform bound on  $H^s(\mathbb{S})$ , and we can derive the same existence theorem for the case  $\tau = 0$  (see Remark 2.11), but only for the case  $s \geq -\frac{n-1}{2}$  (which does not include all radon signed measures).

Another remark is that if we change the sign in front of the alignment term in Smoluchowski equation (3) (taking  $K(\omega, \bar{\omega}) = \omega \cdot \bar{\omega}$ , every particle tends to go away from the mean direction), then we can derive a conservation relation in the same way. But here the “dissipation term” is  $\tilde{\mathcal{D}}(f) = 2\tau \|f - 1\|_{\tilde{H}^{-\frac{n-3}{2}}}^2 + \frac{2}{(n-2)!} |J[f]|^2 \geq 2\tau(n-1)\mathcal{H}(f)$ , without any condition on  $\tau > 0$ . So in any Sobolev space  $H^s(\mathbb{S})$ , with  $s > -\frac{n-1}{2}$  we have global exponential decay of the solution to the uniform distribution, with rate  $(n-1)\tau$ .

**4.2. Study of the supercritical case  $\tau < \frac{1}{n}$ .** In this section, we fix  $\tau < \frac{1}{n}$  and we study the behavior of a solution as  $t \rightarrow +\infty$ . We will write  $\kappa$  for  $\kappa(\tau)$  and  $c$  for  $c(\kappa(\tau))$ . We first establish that the limit set of equilibria  $\mathcal{E}_\infty$  given by the LaSalle principle (Proposition 3.2) depends only on the fact that  $J[f_0]$  is zero or not.

**PROPOSITION 4.4.** *If  $J[f_0] = 0$ , then  $\mathcal{E}_\infty$  is reduced to the uniform distribution. Equation (3) becomes the heat equation. We have exponential decay to the uniform distribution with rate  $2n\tau$  in any  $H^s(\mathbb{S})$ .*

*If  $J[f_0] \neq 0$ , then  $J[f(t)] \neq 0$ , for all  $t > 0$ . The limit set  $\mathcal{E}_\infty = \{M_{\kappa\Omega}, \Omega \in \mathbb{S}\}$  consists of all the nonisotropic equilibria. Furthermore, we have for any  $s \in \mathbb{R}$ ,*

$$(44) \quad \lim_{t \rightarrow \infty} \|f(t) - M_{\kappa\Omega(t)}\|_{H^s} = 0,$$

where  $\Omega(t) = \frac{J[f(t)]}{|J[f(t)]|}$  is the mean direction of  $f(t)$ .

*Proof.* First of all, we write the equation for  $J[f]$ , multiplying (3) by  $\omega$  and integrating on the sphere. We get

$$(45) \quad \begin{aligned} \frac{d}{dt} J[f] &= -\tau(n-1)J[f] + \left( \int_{\mathbb{S}} (\text{Id} - \omega \otimes \omega) f \, d\omega \right) J[f] \\ &= \left( (1 - (n-1)\tau)\text{Id} - \int_{\mathbb{S}} \omega \otimes \omega f \right) J[f], \end{aligned}$$

which can be viewed as a first-order linear ODE of the form  $\frac{d}{dt} J[f] = M(t)J[f]$ . The matrix  $M$  is a smooth function of time, so we have a global unique solution. Consequently, if  $J[f(t_0)] = 0$  for  $t_0 \geq 0$ , then we have  $J[f(t)] = 0$  for all  $t \geq 0$  and (3) reduces to the heat equation. The distribution  $f$  has no component on the first eigenspace of the Laplace–Beltrami operator, and the second eigenvalue is  $2n$ , so we have exponential decay with rate  $2n\tau$  in any  $H^s$  norm.

Now we suppose that  $J[f_0] \neq 0$ , so by the previous argument we have  $J[f(t)] \neq 0$  for all  $t \geq 0$ . There are two possibilities for the limiting set: either the uniform distribution, or the set  $\{M_{\kappa\Omega}, \Omega \in \mathbb{S}\}$  (by Proposition 3.6, they do not have the same level of free energy).

In the first case, by the LaSalle principle,  $f(t)$  converges to the uniform distribution. Then the matrix  $M(t) = (1 - (n-1)\tau)\text{Id} - \int_{\mathbb{S}} \omega \otimes \omega f$  converges to  $(n-1)\left(\frac{1}{n} - \tau\right)\text{Id}$ . Using the ODE for  $J[f]$ , we get

$$\frac{1}{2} \frac{d}{dt} |J[f]|^2 = J[f] \cdot M(t)J[f] \geq \left( (n-1) \left( \frac{1}{n} - \tau \right) - \varepsilon \right) |J[f]|^2$$

for  $t$  sufficiently large. Taking  $\varepsilon$  sufficiently small, we get that  $|J[f]|$  tends to infinity, which is a contradiction.

So we have that  $\mathcal{E}_\infty = \{M_{\kappa\Omega}, \Omega \in \mathbb{S}\}$ . Now suppose that  $\|f(t) - M_{\kappa\Omega(t)}\|_{H^s}$  does not tend to 0. We take  $t_n$  tending to infinity such that  $\|f(t_n) - M_{\kappa\Omega(t_n)}\|_{H^s} \geq \varepsilon > 0$ .

By our LaSalle principle, there exists  $\Omega_n \in \mathbb{S}$  such that  $\|f(t_n) - M_{\kappa\Omega_n}\|_{H^s} \rightarrow 0$ . Up to extracting, we can suppose that  $\Omega_n \rightarrow \Omega_\infty \in \mathbb{S}$ , so  $f(t_n) \rightarrow M_{\kappa\Omega_\infty}$  in  $H^s(\mathbb{S})$ . In particular we have that  $J[f(t_n)] \rightarrow c(\kappa)\Omega_\infty$ , and then  $\Omega(t_n) \rightarrow \Omega_\infty$ . Then  $M_{\kappa\Omega(t_n)}$  converges to  $M_{\kappa\Omega_\infty}$ , giving that  $\|f(t_n) - M_{\kappa\Omega(t_n)}\|_{H^s} \rightarrow 0$ , which is a contradiction.  $\square$

Now we focus on the case  $J[f_0] \neq 0$ . We define  $\Omega(t)$  as in the previous proposition, and we will expand the solution around  $M_{\kappa\Omega(t)}$ . We first show the convergence in  $L^2(\mathbb{S})$  to a given equilibrium, with exponential rate, assuming conditions on the initial data.

PROPOSITION 4.5. *There exists an “asymptotic rate”  $r_\infty(\tau) > 0$  satisfying the following property.*

*Suppose that  $\|f(t) - M_{\kappa\Omega(t)}\|_{H^s}$  is uniformly bounded on  $[t_0, +\infty)$  by a constant  $K$ , with  $s > \frac{3(n-1)}{2}$ . Then for all  $r < r_\infty(\tau)$ , there exist  $\Omega_\infty \in \mathbb{S}$  and  $\delta, C > 0$  such that if  $\|f(t_0) - M_{\kappa\Omega(t_0)}\|_{L^2} \leq \delta$ , we have*

$$\|f(t) - M_{\kappa\Omega_\infty}\|_{L^2} \leq C\|f(t_0) - M_{\kappa\Omega(t_0)}\|_{L^2}e^{-r(t-t_0)}.$$

The constants  $\delta$  and  $C$  depend only on  $\tau, s, K$ , and  $r$ . Moreover, as  $\tau \rightarrow \frac{1}{n}$ , we have that  $r_\infty(\tau) \geq 2(n-1)(\frac{1}{n} - \tau) + O((\frac{1}{n} - \tau)^{\frac{3}{2}})$ .

*Proof.* We first introduce some notation. When there is no confusion, we write just  $\Omega$  for  $\Omega(t)$ , and we will always assume  $t \geq t_0$ . We write  $\cos \theta = \omega \cdot \Omega$ . We denote by  $\langle \cdot \rangle_{M_{\kappa\Omega}}$  the mean of a function against the probability measure  $M_{\kappa\Omega}$ .

We have the following identities (we recall, by Lemma 3.4, that  $\beta = c^2 + n\tau - 1$  is positive):

$$\begin{aligned} \langle \omega \rangle_{M_{\kappa\Omega}} &= \langle \cos \theta \rangle_{M_{\kappa\Omega}} \Omega = c\Omega, \\ \langle \cos^2 \theta \rangle_{M_{\kappa\Omega}} &= 1 - (n-1)\tau, \\ (46) \quad \langle (\cos \theta - c)^2 \rangle_{M_{\kappa\Omega}} &= 1 - (n-1)\tau - c^2 = \tau - \beta > 0. \end{aligned}$$

We can write  $f = (1+h)M_{\kappa\Omega}$ ; then we have  $\langle h \rangle_{M_{\kappa\Omega}} = 0$ . Since  $\Omega$  is the direction of  $J[f] = \langle (1+h)\omega \rangle_{M_{\kappa\Omega}}$ , we get that  $\langle h\omega \rangle_{M_{\kappa\Omega}} = \langle h \cos \theta \rangle_{M_{\kappa\Omega}} \Omega$ .

So we can do an expansion of the free energy and its dissipation in terms of  $h$ . Since we know that  $M_{\kappa\Omega(t)}$  is a critical point of  $\mathcal{F}$ , we already know that the expansion of  $\mathcal{F}((1+h)M_{\kappa\Omega}) - \mathcal{F}(M_{\kappa\Omega})$  will contain no term of order 0 or 1 in  $h$ . We get, using (32),

$$\mathcal{F}((1+h)M_{\kappa\Omega}) - \mathcal{F}(M_{\kappa\Omega}) = \tau \frac{1}{2} \langle h^2 \rangle_{M_{\kappa\Omega}} - \frac{1}{2} |\langle h\omega \rangle_{M_{\kappa\Omega}}|^2 + O(\|h\|_\infty^3).$$

Using Sobolev embedding and interpolation, we have (writing  $C$  for a generic constant, depending only on  $\tau, s$ , and  $K$ )

$$\|f - M_{\kappa\Omega}\|_\infty \leq C\|f - M_{\kappa\Omega}\|_{H^{\frac{n-1}{2}}} \leq C\|f - M_{\kappa\Omega}\|_{L^2}^{1-\frac{n-1}{2s}} K^{\frac{n-1}{2s}}.$$

So since  $1 - \frac{n-1}{2s} > \frac{2}{3}$  and  $f - M_{\kappa\Omega} = hM_{\kappa\Omega}$ , with  $M_{\kappa\Omega}$  uniformly bounded below and above, we get that  $\|h\|_\infty^3 = o(\langle h^2 \rangle_{M_{\kappa\Omega}})$  (and, more precisely, for any  $\varepsilon > 0$ , there exists  $\eta > 0$  depending only on  $\varepsilon, \tau, s$ , and  $K$  such that  $\|h\|_\infty^3 \leq \varepsilon \langle h^2 \rangle_{M_{\kappa\Omega}}$  as soon as  $\langle h^2 \rangle_{M_{\kappa\Omega}} \leq \eta$ ). We get

$$(47) \quad \mathcal{F}(f) - \mathcal{F}(M_{\kappa\Omega}) = \frac{1}{2} [\tau \langle h^2 \rangle_{M_{\kappa\Omega}} - \langle h \cos \theta \rangle_{M_{\kappa\Omega}}^2] + o(\langle h^2 \rangle_{M_{\kappa\Omega}}).$$

We use the definition (33) of  $\mathcal{D}(f)$ :

$$\begin{aligned} D(f) &= \langle (1+h)|\nabla(\tau \ln(M_{\kappa\Omega}(1+h)) - \langle(1+h)\omega\rangle_{M_{\kappa\Omega}} \cdot \omega)|^2 \rangle_{M_{\kappa\Omega}} \\ &= \langle (1+h)|\nabla(\tau \ln(1+h) - \langle h \cos \theta \rangle_{M_{\kappa\Omega}} \cos \theta)|^2 \rangle_{M_{\kappa\Omega}} \\ &\geq (1 - \|h\|_\infty) \langle |\nabla(\tau \ln(1+h) - \langle h \cos \theta \rangle_{M_{\kappa\Omega}} \cos \theta)|^2 \rangle_{M_{\kappa\Omega}}. \end{aligned}$$

Now we can derive a Poincaré inequality of the form

$$\langle |\nabla g|^2 \rangle_{M_{\kappa\Omega}} \geq \Lambda_\kappa \langle (g - \langle g \rangle_{M_{\kappa\Omega}})^2 \rangle_{M_{\kappa\Omega}}.$$

Indeed, we use the fact that  $M_{\kappa\Omega}$  is positive and bounded:

$$\begin{aligned} \langle |\nabla g|^2 \rangle_{M_{\kappa\Omega}} &\geq \min M_{\kappa\Omega} \int_{\mathbb{S}} |\nabla g|^2 \\ &\geq \min M_{\kappa\Omega} (n-1) \int_{\mathbb{S}} (g - \int_{\mathbb{S}} g)^2 \\ &\geq \frac{\min M_{\kappa\Omega}}{\max M_{\kappa\Omega}} (n-1) \langle (g - \int_{\mathbb{S}} g)^2 \rangle_{M_{\kappa\Omega}} \\ (48) \qquad &\geq (n-1)e^{-2\kappa} \langle (g - \langle g \rangle_{M_{\kappa\Omega}})^2 \rangle_{M_{\kappa\Omega}}. \end{aligned}$$

Actually this is a rough estimate; we have here  $\Lambda_\kappa \geq (n-1)e^{-2\kappa}$ , but a more precise study of  $\Lambda_\kappa$  can be done using separation of variables and is given in the appendix of [6]. The problem then reduces to finding the smallest eigenvalue of a one-dimensional Sturm–Liouville problem, but even in that case, we did not manage to find a better estimate for now.

So we finally get

$$\begin{aligned} \mathcal{D}(f) &\geq (1 - \|h\|_\infty) \Lambda_\kappa \langle [\tau \ln(1+h) - \tau \langle \ln(1+h) \rangle_{M_{\kappa\Omega}} - \langle h \cos \theta \rangle_{M_{\kappa\Omega}} (\cos \theta - c)]^2 \rangle_{M_{\kappa\Omega}} \\ &\geq (1 - \|h\|_\infty) \Lambda_\kappa \langle [\tau h - \langle h \cos \theta \rangle_{M_{\kappa\Omega}} (\cos \theta - c) + O(\|h\|_\infty^2)]^2 \rangle_{M_{\kappa\Omega}} \\ &\geq (1 - \|h\|_\infty) \Lambda_\kappa (\tau^2 \langle h^2 \rangle_{M_{\kappa\Omega}} - (\beta + \tau) \langle h \cos \theta \rangle_{M_{\kappa\Omega}}^2) + O(\|h\|_\infty^3). \end{aligned}$$

With the same argument as before, we get that

$$(49) \qquad \mathcal{D}(f) \geq \Lambda_\kappa (\tau^2 \langle h^2 \rangle_{M_{\kappa\Omega}} - (\beta + \tau) \langle h \cos \theta \rangle_{M_{\kappa\Omega}}^2) + o(\langle h^2 \rangle_{M_{\kappa\Omega}}).$$

The goal is now to express the bounds in (49) and (47) as the sum of positive terms. Indeed, we expect to have a Grönwall inequality which will give a rate of convergence.

We set  $\alpha = \frac{1}{\tau - \beta} \langle h \cos \theta \rangle_{M_{\kappa\Omega}}$ , and we write  $h = \alpha(\cos \theta - c) + g$ . Using (46), we have that  $\alpha$  is well defined since  $\tau - \beta > 0$  and we get  $\langle g \rangle_{M_{\kappa\Omega}} = 0$  and  $\langle g\omega \rangle_{M_{\kappa\Omega}} = 0$ .

Plugging  $\langle h^2 \rangle_{M_{\kappa\Omega}} = (\tau - \beta)\alpha^2 + \langle g^2 \rangle_{M_{\kappa\Omega}}$  into (47) and (49) gives

$$\begin{aligned} (50) \qquad \mathcal{F}(f) - \mathcal{F}(M_{\kappa\Omega}) &= \frac{1}{2} [\beta(\tau - \beta)\alpha^2 + \tau \langle g^2 \rangle_{M_{\kappa\Omega}}] + o(\langle h^2 \rangle_{M_{\kappa\Omega}}), \\ \mathcal{D}(f) &\geq \Lambda_\kappa (\beta^2(\tau - \beta)\alpha^2 + \tau^2 \langle g^2 \rangle_{M_{\kappa\Omega}}) + o(\langle h^2 \rangle_{M_{\kappa\Omega}}) \\ &\geq \Lambda_\kappa \beta (\beta(\tau - \beta)\alpha^2 + \tau \langle g^2 \rangle_{M_{\kappa\Omega}}) + o(\langle h^2 \rangle_{M_{\kappa\Omega}}). \end{aligned}$$

So for all  $r < \Lambda_\kappa \beta$ , if  $\langle h^2 \rangle_{M_{\kappa\Omega}}$  is sufficiently small, we have  $\mathcal{D}(f) \geq r(\mathcal{F}(f) - \mathcal{F}(M_{\kappa\Omega}))$ . Using the conservation relation (43), there exists  $\delta_0 > 0$  (depending only on  $\tau, s, K$ , and  $r$ ) such that if  $\|f(t) - M_{\kappa\Omega(t)}\|_{L^2} \leq \delta_0$ , we have

$$\frac{d}{dt} [\mathcal{F}(f) - \mathcal{F}(M_{\kappa\Omega})] = -\mathcal{D}(f) \leq -2r[\mathcal{F}(f) - \mathcal{F}(M_{\kappa\Omega})].$$

Then we obtain, for all  $T$ , such that  $\|f - M_{\kappa\Omega}\|_{L^2} \leq \delta_0$  on  $[t_0, T]$ ,

$$\mathcal{F}(f(T)) - \mathcal{F}(M_{\kappa\Omega(T)}) \leq [\mathcal{F}(f(t_0)) - \mathcal{F}(M_{\kappa\Omega(t_0)})]e^{-2r(T-t_0)},$$

and then, using the estimate (50), we get that for  $t \in [t_0, T]$ ,

$$(51) \quad \|f - M_{\kappa\Omega}\|_{L^2} \leq C_0\|f(t_0) - M_{\kappa\Omega(t_0)}\|_{L^2}e^{-r(t-t_0)}.$$

So if we take  $\delta < \frac{\delta_0}{C_0} \leq \delta_0$  and we start with  $\|f(t_0) - M_{\kappa\Omega(t_0)}\|_{L^2} \leq \delta$ , we get that  $\|f - M_{\kappa\Omega}\|_{L^2} \leq \delta_0$  on  $[t_0, T]$  for all  $T \geq t_0$ . Otherwise, the largest of such a  $T$  would satisfy  $\delta_0 = \|f(T) - M_{\kappa\Omega(T)}\|_{L^2} \leq C\delta e^{-r(T-t_0)} < \delta_0$ . So the inequality (51) holds for all  $t \in [t_0, +\infty)$ .

It remains to prove that  $\Omega(t)$  converges to some  $\Omega_\infty$  if we want to have strong convergence to a given steady state. This is possible using the ODE satisfied by  $\Omega$ .

Indeed, we have  $J[f] = c\Omega + \langle h\omega \rangle_{M_{\kappa\Omega}} = (c + \alpha(\tau - \beta))\Omega$ , and then

$$\frac{d}{dt}J[f] = (c + \alpha(\tau - \beta))\frac{d}{dt}\Omega + (\tau - \beta)\Omega\frac{d}{dt}\alpha.$$

So applying  $\text{Id} - \Omega \otimes \Omega$  to the ODE (45) gives an ODE for  $\Omega$  in terms of  $\alpha$  and  $g$ . We get

$$\begin{aligned} (\text{Id} - \Omega \otimes \Omega)\frac{d}{dt}J[f] &= -(\text{Id} - \Omega \otimes \Omega)\left(\int_{\mathbb{S}} \omega \otimes \omega f \, d\omega\right)J[f] \\ &= -(c + \alpha(\tau - \beta))(\text{Id} - \Omega \otimes \Omega)[\langle h \cos \theta \omega \rangle_{M_{\kappa\Omega}} + \langle \cos \theta \omega \rangle_{M_{\kappa\Omega}}]. \end{aligned}$$

Since  $\langle (\cos \theta - c) \cos \theta \omega \rangle_{M_{\kappa\Omega}}$  and  $\langle \cos \theta \omega \rangle_{M_{\kappa\Omega}}$  are parallel to  $\Omega$ , we get that

$$(c + \alpha(\tau - \beta))\frac{d\Omega}{dt} = -(c + \alpha(\tau - \beta))(\text{Id} - \Omega \otimes \Omega)\langle g \cos \theta \omega \rangle_{M_{\kappa\Omega}}.$$

Since  $(c + \alpha(\tau - \beta))$  is the norm of  $J[f]$ , it is never zero, and we get (the notation  $C$  standing for a generic constant depending only on  $r, s, \tau$ , and  $K$ )

$$\left|\frac{d\Omega}{dt}\right| \leq C\sqrt{\langle g^2 \rangle_{M_{\kappa\Omega}}} \leq C\|f - M_{\kappa\Omega}\|_{L^2}.$$

So we have exponential decay of  $\frac{d\Omega}{dt}$  with rate  $r$ ; in particular,  $\Omega$  is converging to some  $\Omega_\infty \in \mathbb{S}$ . More precisely,

$$|\Omega(t) - \Omega_\infty| \leq \int_t^\infty \left|\frac{d\Omega}{dt}\right| dt \leq C\|f(t_0) - M_{\kappa\Omega(t_0)}\|_{L^2}e^{-r(t-t_0)}.$$

Now we have that  $\|M_{\kappa\Omega(t)} - M_{\kappa\Omega_\infty}\|_{L^2} \leq C|\Omega(t) - \Omega_\infty|$  (the function  $\Omega \mapsto e^{\kappa\omega \cdot \Omega}$  from  $\mathbb{S}$  to  $\mathbb{R}$  is globally Lipschitz with a constant independent of  $\omega \in \mathbb{S}$ ). So we get the final estimation:

$$\|f - M_{\kappa\Omega_\infty}\|_{L^2} \leq \|f - M_{\kappa\Omega}\|_{L^2} + \|M_{\kappa\Omega(t)} - M_{\kappa\Omega_\infty}\|_{L^2} \leq C\|f(t_0) - M_{\kappa\Omega(t_0)}\|_{L^2}e^{-r(t-t_0)}.$$

So the proposition is true with  $r_\infty(\tau) = \Lambda_\kappa\beta > 0$ . By the estimate (48), we know that  $\Lambda_\kappa \geq (n - 1)e^{-2\kappa}$ . By the expansions of  $c$  and  $\kappa$  as  $\tau \rightarrow \frac{1}{n}$  given in (41), we get that  $r_\infty(\tau) \geq 2(n - 1)(\frac{1}{n} - \tau) + O((\frac{1}{n} - \tau)^{\frac{3}{2}})$ .  $\square$



By Proposition 4.4, we have that  $f(t) - M_{\kappa\Omega(t)}$  tends to zero in any  $H^s(\mathbb{S})$ . So the hypotheses of Proposition 4.5, for any  $r < r_\infty(\tau)$ , are satisfied for some  $t_0 > 0$ .

Once more, by interpolation and uniform boundedness on  $[t_0, +\infty)$  of the  $H^p$  norm, we have

$$\begin{aligned} \|f - M_{\kappa\Omega_\infty}\|_{H^s} &\leq C \|f - M_{\kappa\Omega_\infty}\|_{L^2}^{1-\frac{s}{p}} \|f - M_{\kappa\Omega_\infty}\|_{H^p}^{\frac{s}{p}} \\ &\leq \tilde{C} \|f(t_0 - M_{\kappa\Omega(t_0)})\|_{L^2}^{1-\frac{s}{p}} e^{-r(1-\frac{s}{p})(t-t_0)}, \end{aligned}$$

so taking  $p$  sufficiently large, we also get exponential convergence for the  $H^s$  norm, with rate  $r(1 - \delta)$  for any  $\delta > 0$ .

Finally, we have that for all  $r < r_\infty(\tau)$  and  $s$ , there exist some time  $t_0$  and  $C > 0$  such that  $\|f - M_{\kappa\Omega_\infty}\|_{H^s} \leq Ce^{-rt}$  for  $t \geq t_0$ . We can even get rid of the constant  $C$  since for any  $\tilde{r} < r$  and  $t$  sufficiently large,  $Ce^{-rt} \leq e^{-\tilde{r}t}$ .

**4.3. Study of the critical case  $\tau = \frac{1}{n}$ .** For any  $\tau \in (0, +\infty) \setminus \{\frac{1}{n}\}$ , we have exponential convergence to some equilibrium. However, the rate of convergence tends to 0 when  $\tau$  is close to  $\frac{1}{n}$  (in the case where  $J[f_0] \neq 0$ ). So we do not expect to have a similar rate of convergence in the critical case.

First of all, we know by Proposition 3.6 that the solution converges (in any  $H^s(\mathbb{S})$ ) to the uniform distribution as time goes to infinity. The goal of this section is to estimate the speed of convergence to this equilibrium.

PROPOSITION 4.6. *Suppose that  $\|f(t) - 1\|_{H^s}$  is uniformly bounded on  $[t_0, +\infty)$  by a constant  $K$ , with  $s > \frac{7(n-1)}{2}$ .*

*Then for all  $C > 1$ , there exists  $\delta > 0$  such that if  $\|f(t_0) - 1\|_{L^2} \leq \delta$ , we have, for  $t \geq t_0$ ,*

$$\|f(t) - 1\|_{L^2} \leq \frac{C}{\sqrt{\frac{1}{\sqrt{2(n+2)}\|f(t_0)-1\|_{L^2}} + \frac{2(n-1)}{n(n+2)}(t-t_0)}}.$$

The constant  $\delta$  depends only on  $\tau, s, K$ , and  $C$ .

*Proof.* As in the previous section, we work on  $[t_0, +\infty)$ . We write  $f = 1 + h$ , and as in the previous case, we suppose that  $J[f_0] \neq 0$ . By the same argument used in Proposition 4.4, we have that  $J[f(t)] \neq 0$  for all  $t > 0$ , so we define  $\Omega(t)$  as the unit vector  $\frac{J[f(t)]}{|J[f(t)]|}$ . Similarly we denote  $\langle \cdot \rangle$  for the mean of a function on the unit sphere and  $\cos \theta$  for  $\omega \cdot \Omega$ .

We have  $\langle h \rangle = 0$ . Since  $\Omega$  is the direction of  $J[f] = \langle (1 + h)\omega \rangle = \langle h\omega \rangle$ , we get that  $\langle h\omega \rangle = \langle h \cos \theta \rangle \Omega$ .

We perform an expansion of the free energy and its dissipation in terms of  $h$ . We get, using (32) and taking  $\tau = \frac{1}{n}$ ,

$$\mathcal{F}(1 + h) = \frac{1}{n}(\frac{1}{2}\langle h^2 \rangle - \frac{1}{6}\langle h^3 \rangle + \frac{1}{12}\langle h^4 \rangle) - \frac{1}{2}\langle h \cos \theta \rangle^2 + O(\|h\|_\infty^5).$$

Now we write  $\alpha = n\langle h \cos \theta \rangle$  and we define

$$(52) \quad g = h - \alpha \cos \theta - \frac{1}{2}\alpha^2(\cos^2 \theta - \frac{1}{n}) - \frac{1}{6}\alpha^3(\cos^3 \theta - \frac{3}{n+2} \cos \theta).$$

We have  $\langle \cos^4 \theta \rangle = \frac{3}{n(n+2)}$  (we have used the formula (40) to compute  $\frac{4!a_2}{a_0} = \langle \cos^4 \theta \rangle$ ). Since we have  $\langle \cos^3 \theta \rangle = \langle \cos \theta \rangle = 0$  and  $\langle \cos^2 \theta \rangle = \frac{1}{n}$ , we get  $\langle g \rangle = \langle g \cos \theta \rangle = 0$ . We will see that the terms of order 2 in  $g$  will not vanish in the expansion of the free

energy and the dissipation term. But we will need to expand the free energy in  $\alpha$  up to order 4 and the dissipation term in  $\alpha$  up to order 6. We have

$$(53) \quad \begin{aligned} \frac{1}{2}\langle h^2 \rangle &= \frac{1}{2}\langle g^2 \rangle + \frac{1}{2n}\alpha^2 + \frac{n-1}{4n^2(n+2)}\alpha^4 + \frac{1}{2}\alpha^2\langle g \cos^2 \theta \rangle + O(\alpha^3\|g\|_\infty + \alpha^5), \\ -\frac{1}{6}\langle h^3 \rangle &= -\frac{n-1}{2n^2(n+2)}\alpha^4 - \frac{1}{2}\alpha^2\langle g \cos^2 \theta \rangle + O(\|g\|_\infty^3 + \alpha\|g\|_\infty^2 + \alpha^3\|g\|_\infty + \alpha^5), \\ \frac{1}{12}\langle h^4 \rangle &= \frac{1}{4n(n+2)}\alpha^4 + O(\|g\|_\infty^4 + \alpha\|g\|_\infty^3 + \alpha^2\|g\|_\infty^2 + \alpha^3\|g\|_\infty + \alpha^5). \end{aligned}$$

We finally get

$$(54) \quad \mathcal{F}(1+h) = \frac{1}{2n}\langle g^2 \rangle + \frac{1}{4n^3(n+2)}\alpha^4 + O(\|g\|_\infty^3 + \alpha\|g\|_\infty^2 + \alpha^3\|g\|_\infty + \alpha^5).$$

Using the inequality  $a^p b^q \leq sa^{\frac{p}{s}} + (1-s)b^{\frac{q}{1-s}}$  for  $s \in (0, 1)$ , with  $a = \alpha$  and  $b = \|g\|_\infty$ , we get that  $\alpha\|g\|_\infty^2 \leq \frac{1}{5}\alpha^5 + \frac{4}{5}\|g\|_\infty^{2+\frac{1}{2}}$  and  $\alpha^3\|g\|_\infty \leq \frac{3}{5}\alpha^5 + \frac{2}{5}\|g\|_\infty^{2+\frac{1}{2}}$ .

By Sobolev embedding and interpolation, as in the previous section, we have

$$(55) \quad \|g\|_\infty \leq C\|g\|_{L^2}^{1-\frac{n-1}{2s}}\|g\|_{H^s}^{\frac{n-1}{2s}},$$

with  $1 - \frac{n-1}{2s} > \frac{6}{7}$ .

Since  $\alpha$  is controlled by  $\|h\|_{H^s}$ , using the definition (52) of  $g$ , we have a bound for  $\|g\|_{H^s}$  on  $[t_0, +\infty)$ , depending only on  $s$  and  $K$ . We finally get  $\|g\|_\infty^{2+\frac{1}{2}} \leq C\langle g^2 \rangle^\mu$ , with  $\mu > \frac{1}{2}(2 + \frac{1}{2})\frac{6}{7} > 1$ .

So using (53) and (54), we get that for any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that if  $\|h\|_{L^2} \leq \delta$ , we have

$$(56) \quad \begin{aligned} (1-\varepsilon)(\langle g^2 \rangle + \frac{1}{n}\alpha^2) &\leq \langle h^2 \rangle \leq (1+\varepsilon)(\langle g^2 \rangle + \frac{1}{n}\alpha^2), \\ (1-\varepsilon)(\frac{1}{2n}\langle g^2 \rangle + \frac{1}{4n^3(n+2)}\alpha^4) &\leq \mathcal{F}(1+h) \leq \frac{1+\varepsilon}{4n^3(n+2)}(2n^2(n+2)\langle g^2 \rangle + \alpha^4). \end{aligned}$$

From that, up to taking a smaller  $\delta$ , we obtain

$$(57) \quad \frac{1-\varepsilon}{1+\varepsilon}2n\mathcal{F}(1+h) \leq \langle h^2 \rangle \leq \frac{1+\varepsilon}{\sqrt{1-\varepsilon}}2\sqrt{n(n+2)}\mathcal{F}(1+h).$$

We now estimate the dissipation term. We use the definition (33) of  $\mathcal{D}(f)$  and the Poincaré inequality to get

$$(58) \quad \begin{aligned} \mathcal{D}(f) &= \langle (1+h)|\nabla(\frac{1}{n}\ln(1+h) - \langle (1+h)\omega \rangle \cdot \omega)|^2 \rangle \\ &= \langle (1+h)|\nabla(\frac{1}{n}\ln(1+h) - \langle h \cos \theta \rangle \cos \theta)|^2 \rangle \\ &\geq \frac{n-1}{n^2}(1 - \|h\|_\infty)\underbrace{\langle \ln(1+h) - \langle \ln(1+h) \rangle - n\langle h \cos \theta \rangle \cos \theta \rangle}_{S(h)}. \end{aligned}$$

We have

$$\begin{aligned} S(h) &= \ln(1+h) - \langle \ln(1+h) \rangle - n\langle h \cos \theta \rangle \cos \theta \\ &= h - \langle h \rangle - \alpha \cos \theta - \frac{1}{2}(h^2 - \langle h^2 \rangle) + \frac{1}{3}(h^3 - \langle h^3 \rangle) + O(\|h\|^4). \end{aligned}$$

We compute

$$\begin{aligned} h - \langle h \rangle - \alpha \cos \theta &= g + \frac{1}{2}\alpha^2(\cos^2 \theta - \frac{1}{n}) + \frac{1}{6}\alpha^3(\cos^3 \theta - \frac{3}{n+2}\cos \theta), \\ -\frac{1}{2}(h^2 - \langle h^2 \rangle) &= -\frac{1}{2}(\alpha^2 + \alpha^3 \cos \theta)(\cos^2 \theta - \frac{1}{n}) + O(\|g\|^2 + \alpha\|g\|_\infty + \alpha^4), \\ \frac{1}{3}(h^3 - \langle h^3 \rangle) &= \frac{1}{3}\alpha^3 \cos^3 \theta + O(\|g\|_\infty^3 + \alpha\|g\|_\infty^2 + \alpha^2\|g\|_\infty + \alpha^4). \end{aligned}$$

So

$$(59) \quad \begin{aligned} \langle \mathcal{S}(h)^2 \rangle &= \langle [g + \frac{1}{6}\alpha^3(\frac{3}{n} - \frac{3}{n+2})\cos\theta]^2 \rangle + O(\|g\|^3 + \alpha\|g\|^2 + \alpha^4\|g\|_\infty + \alpha^7) \\ &= \langle g^2 \rangle + \frac{1}{n^3(n+2)^2}\alpha^6 + O(\|g\|_\infty^3 + \alpha\|g\|_\infty^2 + \alpha^4\|g\|_\infty + \alpha^7). \end{aligned}$$

As before, we get that  $\alpha\|g\|_\infty^2 \leq \frac{1}{7}\alpha^7 + \frac{6}{7}\|g\|_\infty^{2+\frac{1}{3}}$  and  $\alpha^4\|g\|_\infty \leq \frac{4}{7}\alpha^7 + \frac{3}{7}\|g\|_\infty^{2+\frac{1}{3}}$ . Using (55), we get  $\|g\|_\infty^{2+\frac{1}{3}} \leq C\langle g^2 \rangle^\mu$ , with  $\mu > \frac{1}{2}(2 + \frac{1}{3})\frac{6}{7} = 1$ . So using (58) and (59), up to taking a smaller  $\delta$ , we have, for  $\|h\|_{L^2} \leq \delta$ ,

$$\mathcal{D}(f) \geq (1 - \varepsilon)\frac{n-1}{n^2}(\langle g^2 \rangle + \frac{1}{n^3(n+2)^2}\alpha^6).$$

Now for any  $C, C' > 0$ , if we take  $\alpha$  and  $g$  sufficiently small (so again up to taking a smaller  $\delta$ ), we have that  $C\langle g^2 \rangle + \alpha^6 \geq (C'\langle g^2 \rangle + \alpha^4)^{\frac{3}{2}}$ . So we get

$$\mathcal{D}(f) \geq (1 - \varepsilon)\frac{n-1}{n^3(n+2)^2}(2n^2(n+2)\langle g^2 \rangle + \alpha^4)^{\frac{3}{2}}.$$

Putting this together with (56) and the conservation relation (43), we get that for any  $0 < \varepsilon < 1$ , there exists  $\delta_0 > 0$  such that, as soon as  $\|h\|_{L^2} \leq \delta_0$ , we have

$$\frac{d}{dt}\mathcal{F}(f) = -\mathcal{D}(f) \leq -\frac{8(n-1)(1-\varepsilon)}{(1+\varepsilon)^{\frac{3}{2}}\sqrt{n(n+2)}}[\mathcal{F}(f)]^{\frac{3}{2}}.$$

Then we obtain, for all  $T$  such that  $\|h\|_{L^2} \leq \delta_0$  on  $[t_0, T]$ ,

$$(60) \quad \mathcal{F}(f(T))^{-\frac{1}{2}} \geq \mathcal{F}(f(t_0))^{-\frac{1}{2}} + \frac{4(n-1)(1-\varepsilon)}{(1+\varepsilon)^{\frac{3}{2}}\sqrt{n(n+2)}}(t - t_0).$$

Then, using (57), we get that for  $t \in [t_0, T]$ ,

$$\|h\|_{L^2}^{-2} \geq \frac{\sqrt{1-\varepsilon}}{(1+\varepsilon)2\sqrt{n(n+2)}} \left[ \sqrt{\frac{2n(1-\varepsilon)}{1+\varepsilon}}\|h(t_0)\|_{L^2}^{-1} + \frac{4(n-1)(1-\varepsilon)}{(1+\varepsilon)^{\frac{3}{2}}\sqrt{n(n+2)}}(t - t_0) \right].$$

We write  $C = \frac{(1+\varepsilon)^{\frac{5}{4}}}{(1-\varepsilon)^{\frac{3}{4}}}$  (a one-to-one correspondence between  $0 < \varepsilon < 1$  and  $C > 1$ ), and we get

$$(61) \quad \|h\|_{L^2} \leq C \left[ \frac{1}{\sqrt{2(n+2)}\|h(t_0)\|_{L^2}} + \frac{2(n-1)}{n(n+2)}(t - t_0) \right]^{-\frac{1}{2}}.$$

So if we take  $\delta < \min(\delta_0, \frac{1}{C^2\sqrt{2(n+2)}}\delta_0^2)$  and  $\|h(t_0)\|_{L^2} \leq \delta$ , we get that  $\|h\|_{L^2} \leq \delta_0$  on  $[t_0, T]$  for all  $T \geq t_0$ . Otherwise, the largest of such a  $T$  would satisfy

$$\delta_0 = \|h(T)\|_{L^2} \leq C \left[ \frac{1}{\sqrt{2(n+2)}\delta} \right]^{-\frac{1}{2}} < \delta_0.$$

So the inequality (61) holds for all  $t \in [t_0, +\infty)$ , which ends the proof.  $\square$

With this proposition, since  $f$  tends to the uniform distribution in any  $H^s(\mathbb{S})$ , we get that for any  $r < \frac{2(n-1)}{n(n+2)}$ , there exists  $t_0$  such that we have  $\|f(t) - 1\|_{L^2} \leq \frac{1}{\sqrt{r(t-t_0)}}$  for  $t \geq t_0$ . We can even get rid of the  $t_0$  in this inequality since for any  $r < \tilde{r} < \frac{2(n-1)}{n(n+2)}$ , for  $t$  sufficiently large, we have  $\frac{1}{\sqrt{\tilde{r}(t-t_0)}} \leq \frac{1}{\sqrt{rt}}$ .

As in the previous section, using interpolation to deal with the other Sobolev norms of the solution would lead, for any  $\eta > 0$  and  $t$  sufficiently large, to an inequality of the form  $\|f(t) - 1\|_{H^p} \leq C_\eta t^{-\frac{1}{2} + \eta}$ . But we can actually do slightly better. Indeed, we have, following the notation of the proof and using (52),

$$\|h\|_{H^s} \leq |\alpha| \|\cos \theta\|_{H^p} + C_2 \alpha^2 + C_3 |\alpha|^3 + \|g\|_{H^p}.$$

We have  $\|\cos \theta\|_{H^p} = (n - 1)^{\frac{p}{2}}$ . We take  $t_0 > 0$  satisfying the conditions of the proposition and such that  $\|h\|_{L^2} \leq \delta$ . We have that  $g$  is uniformly bounded in any  $H^p(\mathbb{S})$ , and so by interpolation, we have  $\|g\|_{H^s} \leq C_\eta \|g\|_{L^2}^{1-\eta}$  for any  $\eta > 0$ . Now using (60) and (56), we get

$$\left(\frac{1}{2n} \langle g^2 \rangle\right) + \frac{1}{4n^3(n+2)} \alpha^4)^{-\frac{1}{2}} \geq \frac{4(n-1)(1-\varepsilon)^{\frac{3}{2}}}{(1+\varepsilon)^{\frac{3}{2}} \sqrt{n(n+2)}} (t - t_0),$$

which gives  $\|g\|_{L^2} = O(t^{-1})$  and  $\alpha^2 \leq \frac{(1+\varepsilon)^{\frac{3}{2}} n(n+2)}{2(n-1)(1-\varepsilon)^{\frac{3}{2}} (t-t_0)}$ . So finally, for any  $\eta > 0$ ,

we have that  $\|h\|_{H^p} \leq (n - 1)^{\frac{p}{2}} \sqrt{\frac{(1+\varepsilon)^{\frac{3}{2}} n(n+2)}{2(n-1)(1-\varepsilon)^{\frac{3}{2}} (t-t_0)}} + O(t^{-1+\eta})$ . This gives that there

exists  $t_1 \geq t_0$  such that for  $t \geq t_1$ , we have  $\|h\|_{H^p} \leq (1+\varepsilon)(n-1)^{\frac{p}{2}} \sqrt{\frac{(1+\varepsilon)^{\frac{3}{2}} n(n+2)}{2(n-1)(1-\varepsilon)^{\frac{3}{2}} (t-t_0)}}$ .

This is true for any  $\varepsilon > 0$ . In conclusion, we have that for any  $r < \frac{2}{n(n-1)^{p-1}(n+2)}$ , there exists  $t_1$  such that for  $t \geq t_1$ , we have  $\|f(t) - 1\|_{H^p} \leq \frac{1}{\sqrt{rt}}$ .

**4.4. Summary.** In summary we can state the following theorem.

**THEOREM 4.7** (convergence to equilibrium). *Suppose  $f_0$  is a probability measure belonging to  $H^s(\mathbb{S})$  (this is always the case for some  $s < -\frac{n-1}{2}$ ).*

*Then there exists a unique weak solution  $f$  to Smoluchowski equation (3) satisfying the initial condition  $f(0) = f_0$ .*

*Furthermore, this is a classical solution, positive for all time  $t > 0$  and belonging to  $C^\infty((0, +\infty) \times \mathbb{S})$ .*

*If  $J[f_0] \neq 0$ , then we have the following three cases, depending on  $\tau$ :*

- *If  $\tau > \frac{1}{n}$ , then  $f$  converges exponentially fast to the uniform distribution, with global rate  $(n - 1)(\tau - \frac{1}{n})$  in any  $H^p$  norm.*

*More precisely, for all  $t_0 > 0$ , there exists a constant  $C > 0$  depending only on  $t_0, s, p, n$ , and  $\tau$  such that for all  $t \geq t_0$ , we have*

$$\|f(t) - 1\|_{H^p} \leq C \|f_0\|_{H^s} e^{-(n-1)(\tau - \frac{1}{n})t}.$$

- *If  $\tau < \frac{1}{n}$ , then there exists  $\Omega \in \mathbb{S}$  such that  $f$  converges exponentially fast to  $M_{\kappa\Omega}$ , with asymptotic rate  $r_\infty(\tau) > 0$  in any  $H^p$  norm.*

*More precisely, for all  $r < r_\infty(\tau)$ , there exists  $t_0 > 0$  (depending on  $f_0$ ) such that for all  $t > t_0$ , we have*

$$\|f(t) - M_{\kappa\Omega}\|_{H^p} \leq e^{-rt}.$$

*When  $\tau$  is close to  $\frac{1}{n}$  we have that  $r_\infty(\tau) \sim 2(n - 1)(\frac{1}{n} - \tau)$ .*

- *If  $\tau = \frac{1}{n}$ , then  $f$  converges to the uniform distribution in any  $H^p$  norm, with asymptotic rate  $\sqrt{\frac{n(n-1)^{p-1}(n+2)}{2t}}$ .*

More precisely, for all  $r < \frac{2}{n(n-1)^{p-1}(n+2)}$ , there exists  $t_0 > 0$  (depending on  $f_0$ ) such that for all  $t > t_0$ , we have

$$\|f(t) - 1\|_{H^p} \leq \frac{1}{\sqrt{rt}}.$$

If  $J[f_0] = 0$ , then the equation reduces to the heat equation on the sphere, so  $f$  converges to the uniform distribution exponentially with global rate  $2n\tau$  in any  $H^p$  norm.

For the subcritical case  $\tau > \frac{1}{n}$ , we used Theorem 4.1. In the case where  $p < -\frac{n-1}{2}$ , a simple embedding gives  $\|f(t) - 1\|_{H^p} \leq \|f(t) - 1\|_{H^{-\frac{n-1}{2}}}$ , so we need only show the result for  $p \geq -\frac{n-1}{2}$ . We get

$$\|f - 1\|_{H^p}^2 \leq C \|f(t_0) - 1\|_{H^p} e^{-(n-1)(\tau - \frac{1}{n})(t-t_0)} \leq C \|f(t_0)\|_{H^p} e^{-(n-1)(\tau - \frac{1}{n})(t-t_0)}.$$

The last inequality comes from the fact that  $f(t_0)$  is a probability density function, so  $f(t_0) - 1$  is the orthogonal projection of  $f(t_0)$  on the space of mean zero functions. Using Proposition 2.9, we get  $\|f(t_0)\|_{H^p} \leq C_{t_0} \|f_0\|_{H^s}$  in the case  $p \geq s$ . Otherwise we use just a simple embedding to first get  $\|f(t_0)\|_{H^p} \leq \|f(t_0)\|_{H^s}$  and then by the same proposition  $\|f(t_0)\|_{H^p} \leq C \|f_0\|_{H^s}$ .

Then the results in the case  $\tau < \frac{1}{n}$  and  $\tau = \frac{1}{n}$  are a summary of the conclusions of the two previous subsections. However, although it gives a clear understanding of how fast the solution converges to the equilibrium, in some sense, this summary is not as accurate as Propositions 4.5 and 4.6, which give a kind of stability: starting close to an equilibrium, the solution stays close.

**5. Conclusion.** In this paper, we have investigated all the possible dynamics in large time for the Smoluchowski equation (3) with dipolar potential. We have obtained a rate of convergence towards the equilibrium given any initial condition and any noise parameter  $\tau > 0$  for any dimension  $n \geq 2$ .

The rate of convergence to the anisotropic steady state, in the case  $\tau < \frac{1}{n}$ , depends on a Poincaré constant which does not seem easy to estimate. A better knowledge of the behavior of this constant, for example as the noise parameter  $\tau$  tends to zero, would be useful for understanding the limiting case  $\tau = 0$ , where we have existence and uniqueness of the solution. In this limit, the steady states are given by the sum of two antipodal Dirac masses  $(1 - \alpha)\delta_\Omega + \alpha\delta_{-\Omega}$  with  $\Omega \in \mathbb{S}$  and  $0 \leq \alpha \leq \frac{1}{2}$ . We conjecture that if the initial condition is continuous (and with nonzero initial momentum), then the solution converges to one of these steady states, with  $\alpha = 0$ .

It should also be possible to get the same kind of rates for the Maier-Saupe potential, but there the classification of the initial conditions leading to a given type of equilibria is much more difficult, in particular in the case where two types of equilibria are stable.

**Appendix A.**

**A.1. Using the spherical harmonics.** For the following we will use the spherical harmonics, so we recall some preliminaries results. We fix  $n \geq 2$  and work on  $\mathbb{R}^n$  and its unit sphere  $\mathbb{S}_{n-1}$ .

DEFINITION A.1. A spherical harmonic of degree  $\ell$  on  $\mathbb{S}_{n-1}$  is the restriction to  $\mathbb{S}_{n-1}$  of a homogeneous polynomial of degree  $\ell$  in  $n$  variables (seen as a function  $\mathbb{R}^n \rightarrow \mathbb{R}$ ) which is a harmonic function (a function  $P$  such that  $\Delta P = 0$ , where  $\Delta$

is the usual Laplace operator in  $\mathbb{R}^n$ ). We denote by  $\mathcal{H}_\ell^{(n)}$  the set of spherical harmonics of degree  $\ell$  on  $\mathbb{S}_{n-1}$  (including 0, so they are vector spaces).

We know that the space of homogeneous polynomials of degree  $\ell$  in  $n$  variables has dimension  $\binom{n+\ell-1}{n-1}$  (the number of  $n$ -tuples  $(i_1, \dots, i_n)$  of sum  $\ell$ ). Writing an arbitrary homogeneous polynomial  $P$  of degree  $\ell$  under the form  $P = \sum_{i=0}^\ell Q_{\ell-i} X_n^i$ , with the polynomials  $Q_i$  being homogeneous of degree  $i$  in the first  $n-1$  variables, and imposing that  $P$  be a harmonic function gives the following conditions (taking the term in  $X_n^{i-2}$ ) for  $i \in \llbracket 0, \ell-2 \rrbracket$ :  $\Delta Q_{\ell-i} + (i+1)(i+2)Q_{\ell-i-2} = 0$ . Finally, the polynomial  $P$  is determined only by the polynomials  $Q_\ell$  and  $Q_{\ell-1}$  in  $n-1$  variables of respective degrees  $\ell$  and  $\ell-1$ . This gives the dimension of the space of spherical harmonics.

PROPOSITION A.2. *The dimension of  $\mathcal{H}_\ell^{(n)}$  is given by*

$$k_\ell^{(n)} = \binom{n+\ell-2}{n-2} + \binom{n+\ell-3}{n-2} = \binom{n+\ell-1}{n-1} - \binom{n+\ell-3}{n-1}.$$

The second expression comes from two successive applications of Pascal’s triangle rule and will be useful in the following. It can also be seen by the following property:<sup>1</sup> every homogeneous polynomial  $P$  of degree  $\ell$  can be decomposed in a unique way as  $H + |X|^2 Q$ , where  $H$  is harmonic of degree  $\ell$  and  $Q$  is homogeneous of degree  $\ell-2$ . Iterating this decomposition, we get

$$P = H_\ell + |X|^2 H_{\ell-2} + |X|^4 H_{\ell-4} + \dots + \begin{cases} |X|^\ell H_0, & \ell \text{ even,} \\ |X|^{\ell-1} H_1, & \ell \text{ odd,} \end{cases}$$

where the polynomials  $H_i$  are harmonic of degree  $i$ . This shows that any restriction of a polynomial on the sphere is equal to a sum of spherical harmonics (the terms  $|X|^{2i}$  are constant when restricted to the sphere). This gives, with the Stone–Weierstrass theorem, that the sums of spherical harmonics are dense in  $L^2(\mathbb{S}_{n-1})$  (since they are dense in the continuous functions). Together with the radial decomposition of the Laplacian  $\Delta = \frac{1}{r^{n-1}} \partial_r (r^{n-1} \partial_r) + \frac{1}{r^2} \Delta_\omega$  (where  $\Delta_\omega$  is the Laplace–Beltrami operator on the sphere  $\mathbb{S}_{n-1}$ , which is self-adjoint in  $L^2(\mathbb{S}_{n-1})$ ), we get the following result.

PROPOSITION A.3. *The spaces  $\mathcal{H}_\ell^{(n)}$ , for  $\ell \in \mathbb{N}$ , are the eigenspaces of the Laplace–Beltrami operator  $\Delta_\omega$  on the sphere  $\mathbb{S}_{n-1}$  for the eigenvalues  $-\ell(\ell+n-2)$ . They are pairwise orthogonal and complete in  $L^2(\mathbb{S}_{n-1})$ .*

We can construct a basis of  $\mathcal{H}_\ell^{(n)}$  by induction on the dimension using the separation of variables. We describe this construction and will use it in the following.

For a given unit vector  $e_n \in \mathbb{R}^n$ , we take an orthonormal basis  $(e_1, \dots, e_n)$  of  $\mathbb{R}^n$ . Any  $\omega \in \mathbb{S}_{n-1} \setminus \{e_n, -e_n\}$  can be written as  $\omega = \cos \theta e_n + \sin \theta v$ , with  $\theta \in (0, \pi)$  and  $v \in \mathbb{S}_{n-2}$ . We identify  $\mathbb{R}^{n-1}$  with the vector space spanned by  $(e_1, \dots, e_{n-1})$ . The special case  $n=2$  works if we consider  $S_0 = \{e_1, -e_1\}$ .

By convention, the only spherical harmonics on  $S_0$  are the constant functions (of degree 0) and the functions  $e_1 \mapsto c, -e_1 \mapsto -c$  (of degree 1).

<sup>1</sup>This can be shown using the appropriate inner product  $(P, Q) \mapsto P(D)Q$  on the space of homogeneous polynomials  $P$  of degree  $\ell$ , where  $P(D)$  is defined as  $\frac{\partial^\ell}{\partial X_1^{\alpha_1} \dots \partial X_n^{\alpha_n}}$  if  $P = X_1^{\alpha_1} \dots X_n^{\alpha_n}$ , and extended by linearity (so, for example, we have that  $|X|^2(D) = \Delta$ ). If we denote by  $E$  the space of polynomials of the form  $P = |X|^2 Q$ , with  $Q$  of degree  $\ell-2$ , then the orthogonal of  $E$  consists of all the polynomials  $P$  such that for all  $Q$  of degree  $\ell-2$ , we have  $(|X|^2 Q)(D)P = Q(D)\Delta P = 0$ , that is, in all the polynomials  $P$  such that  $\Delta P = 0$ . So the claimed decomposition is just the orthogonal decomposition on  $E$  and  $E^\perp$ .

Now, for  $n \geq 1$ , we choose an orthonormal basis  $(Z_m^1, \dots, Z_m^{k_m^{(n-1)}})$  of  $\mathcal{H}_m^{(n-1)}$  for any  $m \in \mathbb{N}$  and we have the following result.

**PROPOSITION A.4.** *There exist polynomials  $Q_{\ell,m}$  of degree  $\ell - m$  such that if we denote  $Y_{\ell,m}^k(\omega) = Q_{\ell,m}(\cos \theta) \sin^m \theta Z_m^k(v)$ , then the  $Y_{\ell,m}^k$  for  $m \in \llbracket 0, \ell \rrbracket, k \in \llbracket 1, k_m^{(n-1)} \rrbracket$  form an orthonormal basis of  $H_\ell^{(n)}$ .*

*Proof.* Writing  $Y_{\ell,m}^k(\omega) = Q_{\ell,m}(\cos \theta) \sin^m \theta Z_m^k(v)$  and asking it to be a spherical harmonic is equivalent to the following linear ODE for  $Q_{\ell,m}$  (we recall that the Laplace–Beltrami operator is given by  $\sin^{2-n} \theta \partial_\theta (\sin^{n-2} \theta \partial_\theta) + \frac{1}{\sin^2 \theta} \Delta_v$  in these coordinates):

$$\begin{aligned} \sin^{2-n} \partial_\theta ( - \sin^{n+m-1} \theta Q'_{\ell,m}(\cos \theta) + m \cos \theta \sin^{n+m-3} \theta Q_{\ell,m}(\cos \theta) ) \\ - m(m+n-3) Q_{\ell,m}(\cos \theta) \sin^{m-2} \theta = -\ell(\ell+n-2) Q_{\ell,m}(\cos \theta) \sin^m \theta. \end{aligned}$$

We write  $x = \cos \theta$ , and this equation transforms into

$$(1-x^2)Q''_{\ell,m} - (n+2m-1)xQ'_{\ell,m} + (\ell-m)(\ell+n+m-2)Q_{\ell,m} = 0.$$

This equation is a particular form of the Jacobi differential equation, where the two parameters  $\alpha$  and  $\beta$  are equal (also called the Gegenbauer differential equation). One solution of this differential equation is a polynomial, called the ultraspherical polynomial (a particular case of the Jacobi polynomials, also called the Gegenbauer polynomials) and denoted by  $P_i^{(\lambda)}$  following the notation of Szegő in [20]. Precisely, it satisfies the differential equation

$$(1-x^2)y'' - (2\lambda+1)xy' + i(i+2\lambda)y = 0.$$

Taking  $\lambda = m - 1 + \frac{n}{2}$  and  $i = \ell - m$ , we get a solution  $Q_{\ell,m} = \alpha_{\ell,m} P_{\ell-m}^{(m-1+\frac{n}{2})}$ , where  $\alpha_{\ell,m}$  is a positive constant of normalization such that  $Y_{\ell,m}^k$  is of norm 1 in  $L^2(\mathbb{S}_{n-1})$ . We have to be careful here because  $P_i^{(\lambda)}$  is not defined for  $\lambda = 0$ , and so the only special case is  $n = 2, m = 0$ , for which we have a solution  $Q_{\ell,0} = \sqrt{2}T_\ell$ , where  $T_\ell(\cos \theta) = \cos \ell \theta$  (the Chebyshev polynomial of first order of degree  $\ell$ ).

So for a fixed  $\ell$ , we have constructed a family of spherical harmonics  $Y_{\ell,m}^k$  of degree  $\ell$  for  $m \in \llbracket 0, \ell \rrbracket, k \in \llbracket 1, k_m^{(n-1)} \rrbracket$ . They are pairwise orthogonal in  $L^2(\mathbb{S}_{n-1})$  since the  $Z_m^k$  are pairwise orthogonal in  $L^2(\mathbb{S}_{n-2})$ . The size of this family is exactly

$$(62) \quad \sum_{m=0}^{\ell} k_m^{(n-1)} = \sum_{m=0}^{\ell} \binom{n+m-2}{n-2} - \binom{n+m-4}{n-2} = \binom{n+\ell-2}{n-2} + \binom{n+\ell-3}{n-2} = k_\ell^{(n)},$$

which is the dimension of  $H_\ell^{(n)}$ , so we get that the  $Y_{\ell,m}^k$  for  $m \in \llbracket 0, \ell \rrbracket, k \in \llbracket 1, k_m^{(n-1)} \rrbracket$  form an orthonormal basis of  $H_\ell^{(n)}$ .  $\square$

From now on, we will use the construction done in the proof. We have that, for a fixed  $m \geq 0$ , the polynomials  $Q_{\ell,m}$  for  $\ell \geq m$  are a family of orthogonal polynomials for the inner product  $(P, Q) \mapsto \int_{-1}^1 P(x)Q(x)(1-x^2)^{m-1+\frac{n-1}{2}} dx$ .

We will use three properties on the Gegenbauer polynomials (see [20]) for the

following for  $i \geq 0$ ,  $\lambda \neq 0$ , and  $\lambda > -\frac{1}{2}$  (with the convention  $P_{-1}^{(\lambda)} = 0$ ):

$$(63) \quad \int_{-1}^1 (P_i^{(\lambda)}(x))^2 (1-x^2)^{\lambda-\frac{1}{2}} dx = \frac{2^{1-2\lambda} \pi \Gamma(i+2\lambda)}{(i+\lambda) \Gamma^2(\lambda) \Gamma(i+1)},$$

$$(64) \quad (i+1)P_{i+1}^{(\lambda)} = 2(i+\lambda)XP_i^{(\lambda)} - (i+2\lambda-1)P_{i-1}^{(\lambda)},$$

$$(65) \quad (1-X^2)(P_i^{(\lambda)})' = \frac{1}{2(i+\lambda)} \left( (i+2\lambda-1)(i+2\lambda)P_{i-1}^{(\lambda)} - i(i+1)P_{i+1}^{(\lambda)} \right).$$

We have the following normalization for the  $Q_{\ell,m}$ :

$$\int_{-1}^1 Q_{\ell,m}^2(x) (1-x^2)^{m-1+\frac{n-1}{2}} dx = \int_{-1}^1 (1-x^2)^{\frac{n-1}{2}-1} dx.$$

This gives the following relation, together with (63):

$$(66) \quad \alpha_{\ell+1,m}^2 = \frac{(\ell+\frac{n}{2})(\ell+1-m)}{(\ell+\frac{n}{2}-1)(\ell+m+n-2)} \alpha_{\ell,m}^2.$$

By the previous construction, we can decompose  $g = \sum_{k,\ell,m} c_{\ell,m}^k Y_{\ell,m}^k$  and we have  $\int_{\mathbb{S}_{n-1}} g^2 = \sum_{k,\ell,m} |c_{\ell,m}^k|^2$ . Since  $g$  is of mean zero, we have  $c_{0,0}^1 = 0$  (the only spherical harmonic of degree 0 is the constant function 1). So from now on, the indices  $k, \ell, m$  of the sum will mean  $\ell > 0, m \in \llbracket 0, \ell \rrbracket, k \in \llbracket 1, k_m^{(n-1)} \rrbracket$ .

We decompose  $h = \sum_{k,\ell,m} d_{\ell,m}^k Y_{\ell,m}^k$  in the same way. We give a first formula in the form of a lemma.

LEMMA A.5. *We have*

$$(67) \quad e_n \cdot \int_{\mathbb{S}_{n-1}} g \nabla h = \frac{1}{2} \sum_{k,\ell,m} b_{\ell,m} [(\ell+n-1)c_{\ell,m}^k d_{\ell+1,m}^k - \ell c_{\ell+1,m}^k d_{\ell,m}^k],$$

where  $b_{\ell,m} = \frac{\sqrt{\ell-m+1}\sqrt{\ell+m+n-2}}{\sqrt{\ell+\frac{n}{2}-1}\sqrt{\ell+\frac{n}{2}}} \leq 1$ .

*Proof.* We have

$$e_n \cdot \nabla Y_{\ell,m}^k = -\sin \theta \partial_\theta Y_{\ell,m}^k = [(1-X^2)Q'_{\ell,m} - mXQ_{\ell,m}] (\cos \theta) \sin^m \theta Z_m^k(v),$$

and using the inductions formulas (64), (65), and (66), we get

$$(68) \quad (1-X^2)Q'_{\ell,m} - mXQ_{\ell,m} = \frac{1}{2} [b_{\ell-1,m}(\ell+n-2)Q_{\ell-1,m} - b_{\ell,m}\ell Q_{\ell+1,m}],$$

where  $b_{\ell,m}$  is given in the statement of the lemma. In the special case  $n = 2, m = 0$ , using the formula  $Q_{\ell,0}(\cos \theta) = \cos \ell \theta$  gives the same formula as (68), with  $b_{\ell,0} = 1$ .

So we have that  $\int_{\mathbb{S}_{n-1}} e_n \cdot \nabla Y_{\ell,m}^k Y_{\ell',m'}^{k'}$  can be nonzero only if  $m = m', k = k'$ , and  $\ell = \ell' \pm 1$ . By bilinearity, together with the fact that  $Y_{\ell,m}^k$  form an orthonormal basis, this gives the claimed formula.  $\square$

Now we have all the tools to prove Lemma 2.1 (we recall it here).

LEMMA 2.1 (estimates on the sphere).

1. *If  $h$  is in  $\dot{H}^{-s+1}(\mathbb{S})$  and  $g$  is in  $\dot{H}^s(\mathbb{S})$ , the following integral is well defined and we have*

$$\left| \int_{\mathbb{S}} g \nabla h \right| \leq C \|g\|_{\dot{H}^s} \|h\|_{\dot{H}^{-s+1}},$$

where the constant depends only on  $s$  and  $n$ .



2. We have the following estimation for any  $g \in \dot{H}^{s+1}(\mathbb{S})$ :

$$\left| \int_{\mathbb{S}} g \nabla (-\Delta)^s g \right| \leq C \|g\|_{\dot{H}^s}^2,$$

where the constant depends only on  $s$  and  $n$ .

3. We have the following identity for any  $g \in \dot{H}^{-\frac{n-3}{2}}$ :

$$\int_{\mathbb{S}} g \nabla \tilde{\Delta}_{n-1}^{-1} g = 0.$$

*Proof.* Using Lemma A.5, we get

$$\begin{aligned} e_n \cdot \int_{\mathbb{S}_{n-1}} g \nabla h &\leq \frac{1}{2} \sum_{k,\ell,m} \sqrt{\frac{\ell+n-1}{\ell+1}} \left(\frac{\lambda_{\ell+1}}{\lambda_\ell}\right)^{\frac{s}{2}} |\lambda_\ell^{\frac{s}{2}} c_{\ell,m}^k| |\lambda_{\ell+1}^{-\frac{s+1}{2}} d_{\ell+1,m}^k| \\ &\quad + \frac{1}{2} \sum_{k,\ell,m} \sqrt{\frac{\ell}{\ell+n-2}} \left(\frac{\lambda_\ell}{\lambda_{\ell+1}}\right)^{\frac{s}{2}} |\lambda_{\ell+1}^{\frac{s}{2}} c_{\ell+1,m}^k| |\lambda_\ell^{-\frac{s+1}{2}} d_{\ell,m}^k| \\ &\leq C \|g\|_{\dot{H}^s} \|h\|_{\dot{H}^{-s+1}}, \end{aligned}$$

where  $\lambda_\ell = \ell(\ell + n - 2)$  (the eigenvalue of  $-\Delta$  for the spherical harmonics of degree  $\ell$ ). The last line comes from the fact that the sequences  $\sqrt{\frac{\ell+n-1}{\ell+1}} \left(\frac{\lambda_{\ell+1}}{\lambda_\ell}\right)^{\frac{s}{2}}$  and  $\sqrt{\frac{\ell}{\ell+n-2}} \left(\frac{\lambda_\ell}{\lambda_{\ell+1}}\right)^{\frac{s}{2}}$  are bounded (they tend to 1), together with a Cauchy–Schwarz inequality. This gives the first part of the lemma, since this is true for any unit vector  $e_n$ .

Now we take  $h = (-\Delta)^s g$ , which is replacing  $d_{\ell,m}^k$  by  $\lambda_\ell^s c_{\ell,m}^k$  in Lemma A.5. We get

$$\begin{aligned} e_n \cdot \int_{\mathbb{S}_{n-1}} g \nabla (-\Delta)^s g &= \sum_{k,\ell,m} \frac{1}{2} b_{\ell,m} c_{\ell+1,m}^k c_{\ell,m}^k [(\ell + n - 1) \lambda_{\ell+1}^s - \ell \lambda_\ell^s] \\ &\leq \sum_{k,\ell,m} |\lambda_{\ell+1}^{\frac{s}{2}} c_{\ell+1,m}^k| |\lambda_\ell^{\frac{s}{2}} c_{\ell,m}^k| |(\ell + n - 1) \left(\frac{\lambda_{\ell+1}}{\lambda_\ell}\right)^{\frac{s}{2}} - \ell \left(\frac{\lambda_\ell}{\lambda_{\ell+1}}\right)^{\frac{s}{2}}| \\ &\leq C \|g\|_{\dot{H}^s}^2. \end{aligned}$$

Indeed, we have that  $\frac{\lambda_{\ell+1}}{\lambda_\ell} = 1 - \frac{2}{\ell} + O(\ell^{-2})$ , so  $|(\ell + n - 1) \left(\frac{\lambda_{\ell+1}}{\lambda_\ell}\right)^{\frac{s}{2}} - \ell \left(\frac{\lambda_\ell}{\lambda_{\ell+1}}\right)^{\frac{s}{2}}|$  is bounded (it tends to  $(n - 1) + 2s$ ). Since this computation is now valid for any unit vector  $e_n$ , this gives the second part of the lemma.

The last part is straightforward by taking  $h = \tilde{\Delta}_{n-1}^{-1} g$  with Lemma A.5. According to the definition given in (8), we have  $d_{\ell,m}^k = \frac{1}{\ell(\ell+1)\dots(\ell+n-2)} c_{\ell,m}^k$ . We get

$$e_n \cdot \int_{\mathbb{S}_{n-1}} g \nabla \tilde{\Delta}^{-1} g = \sum_{k,\ell,m} \frac{1}{2} b_{\ell,m} c_{\ell+1,m}^k c_{\ell,m}^k \left[ \frac{\ell+n-1}{(\ell+1)\dots(\ell+n-1)} - \frac{\ell}{\ell(\ell+1)\dots(\ell+n-2)} \right] = 0,$$

which is true for any unit vector  $e_n$ . □

**A.2. Analyticity of the solution.** Following [5], we will show that the solution belongs to a special Gevrey class. We define the space  $G_r$  as the set of functions  $g$  (with mean zero) such that  $\tilde{\Delta}_{n-1}^{-\frac{1}{2}} e^{r(-\Delta)^{\frac{1}{2}}} g$  is in  $L^2(\mathbb{S})$ . Using the notation of the

previous proof, this is a Hilbert space associated with the inner product

$$\langle g, h \rangle_{G_r^s} = \sum_{k,\ell,m} \frac{e^{2r\sqrt{\ell(\ell+n-2)}}}{\ell(\ell+1)\dots(\ell+n-2)} c_{\ell,m}^k d_{\ell,m}^k.$$

The norm on this Hilbert space will be written as  $\|\cdot\|_{G_r}$ .

**THEOREM A.6.** *We define  $r(t) = \delta \min\{1, t\}$ .*

*If  $\delta > 0$  is sufficiently small, then for any solution of Smoluchowski equation (3) of the form  $f = 1 + g$ , with  $g(0) \in \dot{H}^{-\frac{n-1}{2}}(\mathbb{S})$ , we have that  $g(t)$  is bounded in  $G_r(t)$  uniformly for  $t \geq 0$ .*

Before giving a proof, we remark that the condition  $g(0) \in \dot{H}^{-\frac{n-1}{2}}(\mathbb{S})$  is not very strong since, by instantaneous regularization (Proposition 2.9), we have it for any time  $t > 0$ . The shape of  $r(t)$  is not optimal, and we will provide a more precise condition in the proof. Now since  $G_r$ , for  $r > 0$ , is a subset of the set of analytical functions on the sphere, we get that any solution becomes instantaneously analytic in space.

*Proof.* We take  $r$  as an arbitrary function of  $t$ , and we will denote its time derivative by  $\dot{r}$ . For a given solution  $f = 1 + g$ , we put  $h = \tilde{\Delta}_{n-1}^{-1} e^{2r(-\Delta)^{\frac{1}{2}}} g$  in (13).

The left-hand side is

$$\begin{aligned} \langle \partial_t g, \tilde{\Delta}_{n-1}^{-1} e^{2r(-\Delta)^{\frac{1}{2}}} g \rangle &= \sum_{k,\ell,m} \frac{e^{2r\sqrt{\ell(\ell+n-2)}}}{\ell(\ell+1)\dots(\ell+n-2)} c_{\ell,m}^k \frac{d}{dt} c_{\ell,m}^k \\ &= \sum_{k,\ell,m} \frac{1}{2} \frac{d}{dt} \left( \frac{e^{2r\sqrt{\ell(\ell+n-2)}}}{\ell(\ell+1)\dots(\ell+n-2)} |c_{\ell,m}^k|^2 \right) - \dot{r} \frac{e^{2r\sqrt{\ell(\ell+n-2)}} |c_{\ell,m}^k|^2}{\sqrt{\ell(\ell+1)\dots(\ell+n-1)} \sqrt{\ell+n-2}} \\ &= \frac{1}{2} \frac{d}{dt} \|g\|_{G_r}^2 - \dot{r} \|(-\Delta)^{\frac{1}{4}} g\|_{G_r}^2. \end{aligned}$$

Using Lemma A.5, we get

$$\begin{aligned} e_n \cdot \langle g, \nabla \tilde{\Delta}_{n-1}^{-1} e^{2r(-\Delta)^{\frac{1}{2}}} g \rangle &= \frac{1}{2} \sum_{k,\ell,m} b_{\ell,m} c_{\ell+1,m}^k c_{\ell,m}^k \frac{e^{2r\sqrt{(\ell+1)(\ell+n-1)}} - e^{2r\sqrt{\ell(\ell+n-2)}}}{(\ell+1)\dots(\ell+n-2)} \\ &\leq \frac{1}{2} \sum_{k,\ell,m} \frac{\sqrt[4]{(\ell+1)(\ell+n-1)} e^{r\sqrt{(\ell+1)(\ell+n-1)}}}{\sqrt{(\ell+1)\dots(\ell+n-2)}(\ell+n-1)} |c_{\ell+1,m}^k| \frac{\sqrt[4]{\ell(\ell+n-2)} e^{r\sqrt{\ell(\ell+n-2)}}}{\sqrt{\ell(\ell+1)\dots(\ell+n-2)}} |c_{\ell,m}^k| \\ &\times \frac{\sqrt[4]{\ell(\ell+n-1)}}{\sqrt[4]{(\ell+1)(\ell+n-2)}} \left( e^{r(\sqrt{(\ell+1)(\ell+n-1)} - \sqrt{\ell(\ell+n-2)})} - e^{-r(\sqrt{(\ell+1)(\ell+n-1)} - \sqrt{\ell(\ell+n-2)})} \right) \\ &\leq \sinh(r(\sqrt{2n} - \sqrt{n-1})) \|(-\Delta)^{\frac{1}{4}} g\|_{G_r}^2. \end{aligned}$$

Indeed, the expression  $\sqrt{(\ell+1)(\ell+n-1)} - \sqrt{\ell(\ell+n-2)}$  is a decreasing function of  $\ell \geq 0$ . Since this is valid for any unit vector  $e_n$ , we get

$$\left| J[g] \cdot \langle g, \nabla \tilde{\Delta}_{n-1}^{-1} e^{2r(-\Delta)^{\frac{1}{2}}} g \rangle \right| \leq \sinh(r(\sqrt{2n} - \sqrt{n-1})) \|(-\Delta)^{\frac{1}{4}} g\|_{G_r}^2.$$

Now since  $\|(-\Delta)^{\frac{1}{4}} g\|_{G_r}^2 \leq \frac{1}{\sqrt{n-1}} \|(-\Delta)^{\frac{1}{2}} g\|_{G_r}^2$  and  $|J[h]| \leq \frac{e^{2r\sqrt{n-1}}}{(n-1)!} |J[g]|$ , we finally get

$$\frac{1}{2} \frac{d}{dt} \|g\|_{G_r}^2 + \left[ \tau - \frac{1}{\sqrt{n-1}} (\dot{r} + \sinh(r(\sqrt{2n} - \sqrt{n-1}))) \right] \|(-\Delta)^{\frac{1}{2}} g\|_{G_r}^2 \leq \frac{e^{2r\sqrt{n-1}}}{(n-2)!}.$$

As soon as  $\dot{r} + \sinh(r(\sqrt{2n} - \sqrt{n-1})) < (\tau - \varepsilon)\sqrt{n-1}$  and  $r$  is bounded in time (for example, the shape given in the statement of the theorem,  $r(t) = \delta \min(1, t)$ , for  $\delta$  sufficiently small), using the Poincaré inequality, we have that  $\|g\|_{G_r}^2$  satisfies an inequality of the form  $\dot{y} + ay \leq b$  with some positive constants  $a$  and  $b$ . Therefore this quantity is uniformly bounded, provided  $g(0)$  is in  $G_{r(0)}$ . So if we have  $r(0) = 0$ , we need only that  $g(0)$  be in  $\dot{H}^{-\frac{n-1}{2}}(\mathbb{S})$ .  $\square$

**Acknowledgments.** The authors would like to thank Pierre Degond for initiating this project, suggesting to work on it, and engaging in many stimulating discussions. The authors are also thankful for the support of the Mathematical Sciences Center at Tsinghua University.

## REFERENCES

- [1] W. BECKNER, *Sharp Sobolev inequalities on the sphere and the Moser–Trudinger inequality*, Ann. of Math. (2), 138 (1993), pp. 213–242.
- [2] F. BOLLEY, J. A. CAÑIZO, AND J. A. CARRILLO, *Mean-field limit for the stochastic Vicsek model*, Appl. Math. Lett., 25 (2012), pp. 339–343.
- [3] C. CAO, M. A. RAMMAHA, AND E. S. TITI, *Gevrey regularity for nonlinear analytic parabolic equations on the sphere*, J. Dynam. Differential Equations, 12 (2000), pp. 411–433.
- [4] P. CONSTANTIN, I. KEVREKIDIS, AND E. S. TITI, *Remarks on a Smoluchowski equation*, Discrete Contin. Dyn. Syst., 11 (2004), pp. 101–112.
- [5] P. CONSTANTIN, E. S. TITI, AND J. VUKADINOVIC, *Dissipativity and Gevrey regularity of a Smoluchowski equation*. Indiana Univ. Math. J., 54 (2005), pp. 949–969.
- [6] P. DEGOND, A. FROUELLE, AND J.-G. LIU, *Macroscopic limits and phase transition in a system of self-propelled particles*, J. Nonlinear Sci., submitted.
- [7] P. DEGOND AND S. MOTSCH, *Continuum limit of self-driven particles with orientation interaction*, Math. Models Methods Appl. Sci., 18 (2008), pp. 1193–1215.
- [8] M. DOI, *Molecular dynamics and rheological properties of concentrated solutions of rodlike polymers in isotropic and liquid crystalline phases*, J. Polymer Sci. Polymer Phys. Ed., 19 (1981), pp. 229–243.
- [9] L. C. EVANS, *Partial Differential Equations*, Grad. Stud. Math. 19, American Mathematical Society, Providence, RI, 1998.
- [10] I. FATKULLIN AND V. SLASTIKOV, *Critical points of the Onsager functional on a sphere*, Nonlinearity, 18 (2005), pp. 2565–2580.
- [11] C. FOIAS AND R. TEMAM, *Gevrey class regularity for the solutions of the Navier–Stokes equations*, J. Funct. Anal., 87 (1989), pp. 359–369.
- [12] A. FROUELLE, *A continuum model for alignment of self-propelled particles with anisotropy and density-dependent parameters*, Math. Models Methods Appl. Sci., to appear.
- [13] E. P. HSU, *Stochastic Analysis on Manifolds*, Grad. Stud. Math. 38, American Mathematical Society, Providence, RI, 2002.
- [14] G. JI, Q. WANG, P. ZHANG, H. WANG, AND H. ZHOU, *Steady states and their stability of homogeneous, rigid, extended nematic polymers under imposed magnetic fields*, Commun. Math. Sci., 5 (2007), pp. 917–950.
- [15] G. JI, Q. WANG, P. ZHANG, AND H. ZHOU, *Study of phase transition in homogeneous, rigid extended nematics and magnetic suspensions using an order-reduction method*, Phys. Fluids, 18 (2006), 123103.
- [16] H. LIU, H. ZHANG, AND P. ZHANG, *Axial symmetry and classification of stationary solutions of Doi–Onsager equation on the sphere with Maier–Saupe potential*, Commun. Math. Sci., 3 (2005), pp. 201–218.
- [17] W. MAIER AND A. SAUPE, *Eine einfache molekulare Theorie des nematischen kristallin-flüssigen Zustandes*, Z. Naturforsch., 13 (1958), pp. 564–566.
- [18] H. P. MCKEAN, *Propagation of chaos for a class of non-linear parabolic equations*, in Stochastic Differential Equations (Lecture Series in Differential Equations 7, Catholic University, Washington, D.C., 1967), Air Force Office of Scientific Research, Arlington, VA, 1967, pp. 41–57.
- [19] L. ONSAGER, *The effects of shape on the interaction of colloidal particles*, Ann. New York Acad. Sci., 51 (1949), pp. 627–659.
- [20] G. SZEGÖ, *Orthogonal Polynomials*, 4th ed., Colloq. Publ. 23, American Mathematical Society,

- Providence, RI, 1975.
- [21] A.-S. SZNITMAN, *Topics in propagation of chaos*, in *École d'Été de Probabilités de Saint-Flour XIX—1989*, Lecture Notes in Math. 1464, Springer, Berlin, 1991, pp. 165–251.
  - [22] T. VICSEK, A. CZIRÓK, E. BEN-JACOB, I. COHEN, AND O. SHOCHET, *Novel type of phase transition in a system of self-driven particles*, Phys. Rev. Lett., 75 (1995), pp. 1226–1229.
  - [23] G. S. WATSON, *Distributions on the circle and sphere*, J. Appl. Probab., 19 (1982), pp. 265–280.
  - [24] H. ZHANG AND P. W. ZHANG, *Stable dynamic states at the nematic liquid crystals in weak shear flow*, Phys. D, 232 (2007), pp. 156–165.
  - [25] H. ZHOU, H. WANG, M. G. FOREST, AND Q. WANG, *A new proof on axisymmetric equilibria of a three-dimensional Smoluchowski equation*, Nonlinearity, 18 (2005), pp. 2815–2825.
  - [26] H. ZHOU, H. WANG, Q. WANG, AND M. G. FOREST, *Characterization of stable kinetic equilibria of rigid, dipolar rod ensembles for coupled dipole–dipole and Maier–Saupe potentials*, Nonlinearity, 20 (2007), pp. 277–297.