

\mathbb{A}^1 -EULER CLASSES: SIX FUNCTORS FORMALISMS, DUALITIES, INTEGRALITY AND LINEAR SUBSPACES OF COMPLETE INTERSECTIONS

TOM BACHMANN AND KIRSTEN WICKELGREN

ABSTRACT. We equate various Euler classes of algebraic vector bundles, including those of [BM00], [KW21], [DJK18], and one suggested by M.J. Hopkins, A. Raksit, and J.-P. Serre. We establish integrality results for this Euler class, and give formulas for local indices at isolated zeros, both in terms of 6-functor formalism of coherent sheaves and as an explicit recipe in commutative algebra of Scheja and Storch. As an application, we compute the Euler classes enriched in bilinear forms associated to arithmetic counts of d -planes on complete intersections in \mathbb{P}^n in terms of topological Euler numbers over \mathbb{R} and \mathbb{C} .

CONTENTS

1. Introduction	2
1.1. Sketch proof and generalizations	3
1.2. Applications	4
1.3. Acknowledgements	4
1.4. Notation and conventions	4
2. Equality of coherent duality and Poincaré–Hopf Euler numbers	5
2.1. Coherent duality Euler Number	5
2.2. Local indices for $n^{\text{GS}}(V, \sigma, \rho)$	8
2.3. Scheja–Storch and coherent duality	10
2.4. The Poincaré–Hopf Euler number with respect to a section	11
3. Cohomology theories for schemes	13
3.1. Introduction	13
3.2. Features of cohomology theories	14
3.3. Some cohomology theories	14
3.4. The yoga of Euler numbers	15
4. Cohomology theories represented by motivic spectra	16
4.1. Aspects of the six functors formalism	16
4.2. Cohomology groups and Gysin maps	18
4.3. Orientations	21
4.4. SL^c -orientations	23
5. Euler classes for representable theories	23
5.1. Tautological Euler class	23
5.2. Integrally defined Euler numbers	23
5.3. Refined Euler classes and numbers	26
5.4. Refined Euler classes and the six functors formalism	27
6. d -Dimensional planes on complete intersections in projective space	28
6.1. Some Euler numbers of symmetric powers on Grassmannians	28
6.2. An arithmetic count of the d -planes on a complete intersection in projective space	33
7. Indices of sections of vector bundles and \mathbb{A}^1 -degrees	35
7.1. \mathbb{A}^1 -degrees	35
7.2. Main result	36
8. Euler numbers in KO-theory and applications	36
Appendix A. KO via framed correspondences	38
Appendix B. Miscellaneous	39
B.1. Cotangent complexes and dualizing complexes	39
B.2. Grothendieck–Witt rings and Witt rings	40
B.3. Regular sequences	41

1. INTRODUCTION

For algebraic vector bundles with an appropriate orientation, there are Euler classes and numbers enriched in bilinear forms. We will start over a field k , and then discuss more general base schemes, obtaining integrality results. Let $\mathrm{GW}(k)$ denote the Grothendieck–Witt group of k , defined to be the group completion of the semi-ring of non-degenerate, symmetric, k -valued, bilinear forms, see e.g. [Lam05]. Let $\langle a \rangle$ in $\mathrm{GW}(k)$ denote the class of the rank 1 bilinear form $(x, y) \mapsto axy$ for a in k^* .

For a smooth, proper k -scheme $f : X \rightarrow \mathrm{Spec} k$ of dimension n , coherent duality defines a trace map $\eta_f : \mathrm{H}^n(X, \omega_{X/k}) \rightarrow k$, which can be used to construct the following Euler number in $\mathrm{GW}(k)$. Let $V \rightarrow X$ be a rank n vector bundle equipped with a relative orientation, meaning a line bundle \mathcal{L} on X and an isomorphism

$$\rho : \det V \otimes \omega_{X/k} \rightarrow \mathcal{L}^{\otimes 2}.$$

For $0 \leq i, j \leq n$, let $\beta_{i,j}$ denote the perfect pairing

$$(1) \quad \beta_{i,j} : \mathrm{H}^i(X, \wedge^j V^* \otimes \mathcal{L}) \otimes \mathrm{H}^{n-i}(X, \wedge^{n-j} V^* \otimes \mathcal{L}) \rightarrow k$$

given by the composition

$$\mathrm{H}^i(X, \wedge^j V^* \otimes \mathcal{L}) \otimes \mathrm{H}^{n-i}(X, \wedge^{n-j} V^* \otimes \mathcal{L}) \xrightarrow{\cup} \mathrm{H}^n(X, \wedge^n V^* \otimes \mathcal{L}^{\otimes 2}) \xrightarrow{\rho} \mathrm{H}^n(X, \omega_{X/k}) \xrightarrow{\eta_f} k.$$

For $i = n-i$ and $j = n-j$, note that $\beta_{i,j}$ is a bilinear form on $\mathrm{H}^i(X, \wedge^j V^* \otimes \mathcal{L})$. Otherwise, $\beta_{i,j} \oplus \beta_{n-i, n-j}$ determines the bilinear form on $\mathrm{H}^i(X, \wedge^j V^* \otimes \mathcal{L}) \oplus \mathrm{H}^{n-i}(X, \wedge^{n-j} V^* \otimes \mathcal{L})$. The alternating sum

$$n^{\mathrm{GS}}(V) := \sum_{0 \leq i, j \leq n} (-1)^{i+j} \beta_{i,j}$$

thus determines an element of $\mathrm{GW}(k)$, which we will call the Grothendieck–Serre duality or coherent duality Euler number. Note that $\beta_{i,j} \oplus \beta_{n-i, n-j}$ in $\mathrm{GW}(k)$ is an integer multiple of h where h denotes the hyperbolic form $h = \langle 1 \rangle + \langle -1 \rangle$, with Gram matrix

$$h = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

This notion of Euler number was suggested by M.J. Hopkins, J.-P. Serre and A. Raksit, and developed by M. Levine and Raksit for the tangent bundle in [LR20].

For a relatively oriented vector bundle V equipped with a section σ with only isolated zeros, an Euler number $n^{\mathrm{PH}}(V, \sigma)$ was defined in [KW21, Section 4] as a sum of local indices

$$n^{\mathrm{PH}}(V, \sigma) = \sum_{x: \sigma(x)=0} \mathrm{ind}_x^{\mathrm{PH}} \sigma.$$

The index $\mathrm{ind}_x^{\mathrm{PH}} \sigma$ can be computed explicitly with a formula of Scheja–Storch [SS75] or Eisenbud–Levine/Khimshiashvili [EL77] [Khi77] (see Sections 2.4 and 2.3) and is also a local degree [KW19] [BBM⁺21] (this is discussed further in Section 7). For example, when x is a simple zero of σ with $k(x) = k$, the index is given by a well-defined Jacobian $\mathrm{Jac} \sigma$ of σ ,

$$\mathrm{ind}_x^{\mathrm{PH}} \sigma = \langle \mathrm{Jac} \sigma(x) \rangle,$$

illustrating the relation with the Poincaré–Hopf formula for topological vector bundles. (For the definition of the Jacobian, see the beginning of Section 6.2.) In [KW21, Section 4, Corollary 36], it was shown that $n^{\mathrm{PH}}(V, \sigma) = n^{\mathrm{PH}}(V, \sigma')$ when σ and σ' are in a family over \mathbb{A}_L^1 of sections with only isolated zeros, where L is a field extension with $[L : k]$ odd. We strengthen this result by equating $n^{\mathrm{PH}}(V, \sigma)$ and $n^{\mathrm{GS}}(V)$; this is the main result of §2.

Theorem 1.1 (see §2.4). *Let k be a field, and $V \rightarrow X$ be a relatively oriented, rank n vector bundle on a smooth, proper k -scheme of dimension n . Suppose V has a section σ with only isolated zeros. Then*

$$n^{\mathrm{PH}}(V, \sigma) = n^{\mathrm{GS}}(V).$$

In particular, $n^{\mathrm{PH}}(V, \sigma)$ is independent of the choice of σ .

Remark 1.2. Theorem 1.1 strengthens Theorem 1 of [BKW20], removing hypothesis (2) entirely. It also simplifies the proofs of [KW21, Theorem 1] and [SW21, Theorems 1 and 2]: it is no longer necessary to show that the sections of certain vector bundles with non-isolated isolated zeros are codimension 2, as in [KW21, Lemmas 54,56,57], and [SW21, Lemma 1], because $n^{\mathrm{PH}}(V, \sigma)$ is independent of σ .

1.1. Sketch proof and generalizations. The proof of the above theorem proceeds in three steps.

- (0) For a section σ of V , we define an Euler number relative to the section using coherent duality and denote it by $n^{\text{GS}}(V, \sigma, \rho)$. If $\sigma = 0$, we recover the absolute Euler number $n^{\text{GS}}(V, \rho)$, essentially by construction.
- (1) For two sections σ_1, σ_2 , we show that $n^{\text{GS}}(V, \sigma_1, \rho) = n^{\text{GS}}(V, \sigma_2, \rho)$. To prove this, one can use homotopy invariance of Hermitian K -theory, or show that $n^{\text{GS}}(V, \sigma_1, \rho) = n^{\text{GS}}(V, \rho)$ by showing an instance of the principle that alternating sums, like Euler characteristics, are unchanged by passing to the homology of a complex.
- (2) If a section σ has isolated zeros, then $n^{\text{GS}}(V, \sigma, \rho)$ can be expressed as a sum of local indices $\text{ind}_{Z/S}(\sigma)$, where Z is (a clopen component of) the zero scheme of σ .
- (3) For Z a local complete intersection in affine space, i.e. in the presence of coordinates, we compute the local degree explicitly, and identify it with the *Scheja–Storch* form [SS75, 3].

Taken together, these steps show that $n^{\text{GS}}(V, \rho)$ is a sum of local contributions given by Scheja–Storch forms, which is essentially the definition of $n^{\text{PH}}(V, \rho)$.

These arguments can be generalized considerably, replacing the Grothendieck–Witt group GW by a more general *cohomology theory* E . We need E to admit transfers along proper lci morphisms of schemes, and an SL^c -orientation (see §3 for more details). Then for step (0) one can define an Euler class $e(V, \sigma, \rho)$ as $z^* \sigma_*(1)$, where z is the zero section. Step (2) is essentially formal; the main content is in steps (1) and (3). Step (1) becomes formal if we assume that E is \mathbb{A}^1 -invariant. In particular, steps (0)–(2) can be performed for SL -oriented cohomology theories represented by motivic spectra; this is explained in §§3,4,5.

It remains to find a replacement for step (3). We offer two possibilities: in §7 we show that, again in the presence of coordinates, the local indices can be identified with appropriate \mathbb{A}^1 -degrees. On the other hand, in §8 we show that for $E = \text{KO}$ the motivic spectrum corresponding to Hermitian K -theory, the local indices are again given by Scheja–Storch forms. This implies the following.

Corollary 1.3 (see Corollary 8.2 and Definition 3.10). *Let $S = \text{Spec}(k)$, where k is a field of characteristic $\neq 2$.¹ Let $\pi : X \rightarrow k$ be smooth and V/X a relatively oriented vector bundle with a non-degenerate section σ . Write $\varpi : Z = Z(\sigma) \rightarrow k$ for the vanishing scheme (which need not be smooth). Then*

$$n^{\text{PH}}(V, \sigma) = \varpi_*(1) \in \text{KO}^0(k) = \text{GW}(k).$$

Here we have used the lci pushforward

$$\varpi_* : \text{KO}^0(Z) \xrightarrow{\rho} \text{KO}^{L_\infty}(Z) \rightarrow \text{KO}^0(k)$$

of Déglise–Jin–Khan [DJK18]. If moreover X is proper then $\varpi_*(1)$ also coincides with $\pi_* z^* z_*(1)$, where $z : X \rightarrow V$ is the zero section (see Corollary 5.21, Corollary 5.18 and Proposition 5.19). This provides an alternative proof that $n^{\text{PH}}(V, \sigma)$ is independent of the choice of σ (under our assumption on k).

Another important example is when E is taken to be the motivic cohomology theory representing Chow–Witt groups. This recovers the Barge–Morel Euler class [BM00] $e^{\text{BM}}(V)$ in $\widetilde{\text{CH}}^r(X, \det V^*)$, which is defined for a base field of characteristic not 2. Suppose that ρ is a relative orientation of V and $\pi : X \rightarrow \text{Spec } k$ is the structure map.

Corollary 1.4. *Let k be a field of characteristic $\neq 2$. Then $\pi_* e^{\text{BM}}(V, \rho) = n^{\text{GS}}(V, \rho)$ in $\text{GW}(k)$.*

Proof. We have $e^{\text{BM}}(V, \rho) = e(V, \rho, H\tilde{Z})$; indeed by Proposition 5.19 $e(V, \rho, H\tilde{Z})$ can be computed in terms of pushforward along the zero section of V , and the exactly the same is true for e^{BM} by definition [BM00, §2.1]. We also have $n^{\text{GS}}(V, \rho) = n(V, \rho, \text{KO})$; indeed the right hand side is represented by the natural symmetric bilinear form on the cohomology of the Koszul complex by Example 8.1, and this is essentially the definition of $n^{\text{GS}}(V, \rho)$.

It thus suffices to prove that $n(V, \rho, H\tilde{Z}) = n(V, \rho, \text{KO}) \in \text{GW}(k)$. Consider the span of ring spectra $H\tilde{Z} \leftarrow \hat{f}_0 \text{KO} \rightarrow \text{KO}$ as in the proof of Proposition 5.4. It induces an isomorphism on $\pi_0(-)(k)$, namely with $\text{GW}(k)$ in all cases. The desired equality follows from naturality of the Euler numbers.

(An alternative argument proceeds as follows. It suffices to prove that $\pi_* e^{\text{BM}}(V, \rho)$ and $n^{\text{GS}}(V, \rho)$ have the same image in $\text{W}(k)$ and \mathbb{Z} . The image of $n^{\text{GS}}(V, \rho)$ in $\text{W}(k)$ is given by $n(V, \rho, \text{KW})$; for this we need only show that $e(V, \rho, \text{KW})$ is represented by the Koszul complex, which is Example 5.20. It will

¹Here and many times in the text, we restrict to bases S with $1/2 \in \mathcal{O}_S$ in order for the classical constructions of Hermitian K -theory to be well-behaved. Forthcoming work by other authors is expected to produce well-behaved Hermitian K -theory spectra in all characteristics, and then all our assumptions on the characteristic can be removed.

thus be enough to show that $n(V, \rho, H\mathbb{Z}) = n(V, \rho, KGL)$ and $n(V, \rho, \underline{W}[\eta^\pm]) = n(V, \rho, KW)$; this follows as before by considering the spans $H\mathbb{Z} \leftarrow \text{kg}l \rightarrow KGL$ and $\underline{W}[\eta^\pm] \leftarrow KW_{\geq 0} \rightarrow KW$.² \square

The left hand side is the Euler class studied by M. Levine in [Lev20]. We do not compare these Euler classes with the obstruction theoretic Euler class of [Mor12, Chapter 8]. Asok and Fasel show that the latter agrees with $\pi_* e^{\text{BM}}(V, \rho)$ up to a unit in $\text{GW}(k)$ [AF16].

1.2. Applications. It is straightforward to see that Euler numbers for cohomology theories are stable under base change (see Corollary 5.3). This implies that when considering vector bundles on varieties which are already defined over e.g. $\text{Spec}(\mathbb{Z}[1/2])$, then the possible Euler numbers are constrained to live in $\text{GW}(\mathbb{Z}[1/2]) = \mathbb{Z}\langle -1, \langle 2 \rangle \rangle \subset \text{GW}(\mathbb{Q})$. Using novel results on Hermitian K-theory [CDH⁺20] allows one to use the base scheme $\text{Spec} \mathbb{Z}$ as well. Proposition 5.4 contains both of these cases, and the $\mathbb{Z}[1/2]$ case is independent of [CDH⁺20]. It follows that for relatively oriented bundles over \mathbb{Z} the Euler numbers can be read off from topological computations (Proposition 5.9). Over $\mathbb{Z}[1/2]$ the topological Euler numbers of the associated real and complex vector bundles together with one further algebraic computation over some field in which 2 is not a square determine the Euler number (and this is again independent of [CDH⁺20]). See Theorem 5.11.

We use this to compute a weighted count of d -dimensional hyperplanes in a general complete intersection

$$\{f_1 = \dots = f_j\} \hookrightarrow \mathbb{P}_k^n$$

over a field k . This count depends only on the degrees of the f_i and not the polynomials f_i themselves: it is determined by associated real and complex counts, for any d and degrees so that the expected variety of d -planes is 0-dimensional and the associated real count is defined. This is Corollary 6.9. For example, combining with results of Finashin–Kharlamov over \mathbb{R} , we have that $160, 839\langle 1 \rangle + 160, 650\langle -1 \rangle$ and

$$32063862647475902965720976420325\langle 1 \rangle + 32063862647475902965683320692800\langle -1 \rangle$$

are arithmetic counts of the 3-planes in a 7-dimensional cubic hypersurface and in a 16-dimensional degree 5-hypersurface respectively. See Example 6.13. This builds on results of Finahin–Kharlamov [FK13], J.L. Kass and the second-named author [KW21], M. Levine [Lev19] [Lev20], S. McKean [McK21], Okonek–Teleman [OT14], S. Pauli [Pau20], J. Solomon [Sol06], P. Srinivasan and the second-named author [SW21], and M. Wendt [Wen20].

1.3. Acknowledgements. We warmly thank M. J. Hopkins for suggesting the definition of the Euler class using coherent duality. We likewise wish to thank A. Ananyevskiy and I. Panin for the reference to [Knu91] giving the existence of Nisnevich coordinates, as well as A. Ananyevskiy, M. Hoyois and M. Levine for useful discussions.

Kirsten Wickelgren was partially supported by National Science Foundation Awards DMS-1552730 and 2001890.

1.4. Notation and conventions.

Grothendieck duality. We believe that if $f : X \rightarrow Y$ is a morphism of schemes which is locally of finite presentation, then there is a well-behaved adjunction

$$f_! : D_{\text{qcoh}}(X) \rightleftarrows D_{\text{qcoh}}(Y) : f^!$$

between the associated derived (∞ -)categories of unbounded complexes of \mathcal{O}_X -modules with quasi-coherent homology sheaves. Unfortunately we are not aware of any references in this generality. Instead, whenever mentioning a functor $f^!$, we implicitly assume that X, Y are separated and of finite type over some noetherian scheme S . In this situation, the functor $f^!$ is constructed for homologically bounded above complexes in [Sta18, Tag 0A9Y] (see also [Har66, Con00]), and this is all we will use.

Vector bundles. We identify locally free sheaves and vector bundles *covariantly*, via the assignment

$$\mathcal{E} \leftrightarrow \text{Spec}(\text{Sym}(\mathcal{E}^*)).$$

While it can be convenient to (not) pass to duals here (as in e.g. [DJK18]), we do not do this, since it confuses the first named author terribly.

²We include this alternative argument because we feel that Example 5.20 is more fully justified in this paper than Example 8.1.

Regular sequences and immersions. Following e.g. [BGI⁺71], by a regular immersion of schemes we mean what is called a Koszul-regular immersion in [Sta18, Tag 0638], i.e. a morphism which is locally a closed immersion cut out by a Koszul-regular sequence. Moreover, by a regular sequence we will always mean a Koszu-regular sequence [Sta18, Tag 062D], and we reserve the term *strongly regular sequence* for the usual notion. A strongly regular sequence is regular [Sta18, Tag 062F], whence a strongly regular immersion is regular. In locally noetherian situations, regular immersions are strongly regular [Sta18, Tags 063L].

Cotangent complexes. For a morphism $f : X \rightarrow Y$, we write L_f for the cotangent complex. Recall that if f is smooth then $L_f \simeq \Omega_f$, whereas if f is a regular immersion then $L_f \simeq C_f[1]$, where C_f denotes the conormal bundle.

Graded determinants. We write $\widetilde{\det} : K(X) \rightarrow \text{Pic}(D(X))$ for the determinant morphism from Thomason–Trobough K -theory to the groupoid of graded line bundles. If C is a perfect complex, then we write $\widetilde{\det}C$ for the determinant of the associated K -theory point. We write $\det C \in \text{Pic}(X)$ for the ungraded determinant.

Given an lci morphism f , we put $\omega_f = \det L_f$ and $\widetilde{\omega}_f = \widetilde{\det}L_f$.

We systematically use graded determinants throughout the text. For example we have the following compact definition of a relative orientation.

Definition 1.5. Let $\pi : X \rightarrow S$ be an lci morphism and V a vector bundle on X . By a *relative orientation* of $V/X/S$ we mean a choice of line bundle \mathcal{L} on X and an isomorphism

$$\rho : \underline{\text{Hom}}(\widetilde{\det}V^*, \widetilde{\omega}_{X/S}) \xrightarrow{\simeq} \mathcal{L}^{\otimes 2}.$$

Note that if π is smooth, this just means that the locally constant functions $x \mapsto \text{rank}(V_x)$ and $x \mapsto \dim \pi^{-1}(\pi(x))$ on X agree, and that we are given an isomorphism $\mathcal{L}^{\otimes 2} \simeq \omega_{X/S} \otimes \det V$. Hence we recover the definition from [KW21, Definition 17].

2. EQUALITY OF COHERENT DUALITY AND POINCARÉ–HOPF EULER NUMBERS

We prove Theorem 1.1 in this section.

2.1. Coherent duality Euler Number. Let $f : X \rightarrow \text{Spec } k$ be a smooth proper k -scheme of dimension n , and let V be a rank n vector bundle, relatively oriented by the line bundle \mathcal{L} on X and isomorphism $\rho : \det V \otimes \omega_{X/k} \rightarrow \mathcal{L}^{\otimes 2}$. Let $\sigma : X \rightarrow V$ be a section. Let $K(\sigma)^\bullet$ denote the Koszul complex

$$0 \rightarrow \wedge^n V^* \rightarrow \wedge^{n-1} V^* \rightarrow \dots \rightarrow V^* \rightarrow \mathcal{O} \rightarrow 0,$$

with \mathcal{O} in degree 0 and differential of degree +1 given by

$$d(v_1 \wedge v_2 \wedge \dots \wedge v_j) = \sum_{i=1}^j (-1)^{i-1} v_i(\sigma) v_1 \wedge \dots \wedge v_{i-1} \wedge v_{i+1} \wedge \dots \wedge v_j.$$

This choice of $K(\sigma)^\bullet$ is $\text{Hom}_{\mathcal{O}}(-, \mathcal{O})$ applied to the Koszul complex of [Eis95, 17.2]. $K(\sigma)^\bullet$ carries a canonical multiplication

$$(2) \quad m : K(\sigma)^\bullet \otimes K(\sigma)^\bullet \rightarrow K(\sigma)^\bullet$$

defined in degree $-p$ by $m = \oplus_{i+j=p} 1_{\wedge^i V^*} \wedge 1_{\wedge^j V^*}$. Composing m with the projection $p : K(\sigma)^\bullet \rightarrow \det V^*[n]$ defines a non-degenerate bilinear form

$$\beta_{(V,\sigma)} : K(V,\sigma) \otimes K(V,\sigma) \rightarrow \det V^*[n],$$

$$\beta_{(V,\sigma)} = pm.$$

Tensoring $\beta_{(V,\sigma)}$ by $\mathcal{L}^{\otimes 2}$ and reordering the tensor factors of the domain, we obtain a non-degenerate symmetric bilinear form on $K(V,\sigma) \otimes \mathcal{L}$ valued in $(\det V^* \otimes \mathcal{L}^{\otimes 2})[n]$. The orientation ρ determines an isomorphism $(\det V^* \otimes \mathcal{L}^{\otimes 2})[n] \rightarrow \omega_{X/k}[n]$. Composing $\beta_{(V,\sigma)} \otimes \mathcal{L}^{\otimes 2}$ with this isomorphism produces a non-degenerate bilinear form

$$\beta_{(V,\sigma,\rho)} : (K(V,\sigma) \otimes \mathcal{L}) \otimes (K(V,\sigma) \otimes \mathcal{L}) \rightarrow \omega_{X/k}[n].$$

Let $D(X)$ denote the derived category of quasi-coherent \mathcal{O}_X -modules. Serre duality determines an isomorphism $Rf_*\omega_{X/k}[n] \cong \mathcal{O}_k$ [Har13, III Corollary 7.2 and Theorem 7.6]. Since Rf_* is lax symmetric monoidal (being right adjoint to a symmetric monoidal functor), we obtain a symmetric morphism

$$Rf_*\beta_{(V,\sigma,\rho)} : [Rf_*(K(V,\sigma) \otimes \mathcal{L})]^{\otimes 2} \rightarrow Rf_*\omega_{X/k}[n] \simeq \mathcal{O}_k,$$

in $D(k)$, which is non-degenerate by Serre duality.

The derived category $D(k)$ is equivalent to the category of graded k -vector spaces, by taking cohomology³. If V is a (non-degenerate) symmetric bilinear form in graded k -vector spaces, denote by $V^{(n)} = V_n \oplus V_{-n}$ (for $n \neq 0$) and $V^{(0)} = V_0$ the indicated subspaces; observe that they also carry (non-degenerate) symmetric bilinear forms.

Definition 2.1. For a relatively oriented rank n vector bundle $V \rightarrow X$ with section σ and orientation ρ , over a smooth and proper variety $f : X \rightarrow k$ of dimension n , the *Grothendieck–Serre-duality Euler number with respect to σ* is

$$n^{\text{GS}}(V, \sigma, \rho) = \sum_{i \geq 0} (-1)^i [(Rf_* \beta_{(V, \sigma, \rho)})^{(i)}] \in \text{GW}(k).$$

Remark 2.2. In order to not clutter notation unnecessarily, we also write the above definition as

$$n^{\text{GS}}(V, \sigma, \rho) = \sum_i (-1)^i [(Rf_* \beta_{(V, \sigma, \rho)})_i].$$

We shall commit to this kind of abuse of notation from now on.

Recall that $n^{\text{GS}}(V, \rho) \in \text{GW}(k)$ was defined in the introduction, in terms of the symmetric bilinear form on $\bigoplus_{i,j} H^i(X, \Lambda^j V^* \otimes \mathcal{L})$.

Proposition 2.3. *For any section σ we have $n^{\text{GS}}(V, \sigma, \rho) = n^{\text{GS}}(V, \rho) \in \text{GW}(k)$.*

To prove Proposition 2.3, we use the hypercohomology spectral sequence $E_r^{i,j}(K^\bullet)$ associated to a complex K^\bullet of locally free sheaves on X with

$$E_1^{i,j}(K^\bullet) := H^j(X, K^i) \Rightarrow R^{i+j} f_* K^\bullet$$

Let F_i denote the resulting filtration on $R^* f_* K^\bullet$, so that

$$\dots \supseteq F_i = \text{Im}(H^*(X, K^{\bullet \geq i}) \rightarrow H^*(X, K^\bullet)) \supseteq F_{i+1} \supseteq \dots$$

Given a perfect symmetric pairing of chain complexes $\beta : K^\bullet \otimes K^\bullet \rightarrow \omega_{X/k}[n]$, cup product induces pairings

$$\beta' : R^* f_* K^\bullet \otimes R^* f_* K^\bullet \rightarrow R^* f_* \omega_{X/k}[n] \rightarrow k$$

and

$$\beta_1 : E_1^{*,*}(K^\bullet) \otimes E_1^{*,*}(K^\bullet) \rightarrow k.$$

The following properties hold:

- (1) Placing the k in the codomain of β_1 in bidegree $(-n, n)$, β_1 is a map of bigraded vector spaces and satisfies the Leibniz rule with respect to d_1 . It thus induces $\beta_2 : E_2^{*,*}(K^\bullet) \otimes E_2^{*,*}(K^\bullet) \rightarrow k$. Then β_2 satisfies the Leibniz rule with respect to d_2 and hence induces β_3 , and so on.
- (2) All the pairings β_i are perfect.
- (3) The pairing β' is compatible with the filtration in the sense that $\beta'(F_i, F_k) = 0$ if $i + k > -n$.
- (4) It follows that β' induces a pairing on $\text{gr}_\bullet R^* f_* K^\bullet$. Under the isomorphism $\text{gr}_\bullet \simeq E_\infty$, it coincides with β_∞ .
- (5) β' is perfect in the filtered sense: the induced pairing $F_i \otimes R^* f_* K^\bullet / F_{-n-i+1} \rightarrow k$ is perfect. (In particular the pairing β' is perfect.)

Remark 2.4. We do not know a reference for these facts, and proving them would take us too far afield. The main idea is that we have a sequence of duality-preserving functors

$$C^{\text{perf}}(X) \xrightarrow{\sigma_\bullet} D(X)^{\text{fil}} \xrightarrow{\pi_*} D(k)^{\text{fil}}.$$

Here $C^{\text{perf}}(X)$ denotes the category of bounded chain complexes of vector bundles, $D(X)^{\text{fil}}$ is the filtered derived category [GP18], and σ_\bullet is the “stupid truncation” functor (composed with forgetting to the filtered derived category). The first duality is with respect to $\underline{\text{Hom}}(-, \omega[n])$, the second with respect to $\underline{\text{Hom}}(-, \sigma_\bullet(\omega[n])) = \underline{\text{Hom}}(-, \omega[n](-n))$, and the third with respect to $\underline{\text{Hom}}(-, k[0](-n))$. There are further duality preserving functors

$$(-)^{\text{gr}} : D(k)^{\text{fil}} \rightarrow D(k)^{\text{gr}} \text{ and } U : D(k)^{\text{fil}} \rightarrow D(k),$$

where $D(X)^{\text{gr}} = \text{Fun}(\mathbb{Z}, D(X))$, with \mathbb{Z} viewed as a discrete category. Hence any perfect pairing $C \otimes C \rightarrow k[0](-n) \in D(k)^{\text{fil}}$ induces a perfect pairing on $H_* C^{\text{gr}} \otimes H_* C^{\text{gr}} \rightarrow k(-n, n)$, satisfying (1), and a pairing $H_* UC \otimes H_* UC \rightarrow k$, satisfying (3, 5). Moreover there is a spectral sequence $E_1 = H_* C^{\text{gr}} \Rightarrow H_* UC$,

³In this section, we treat all categories as 1-categories, i.e. ignore the higher structure of $D(k)$ as an ∞ -category.

satisfying (1) and (4). (2) is obtained from the fact that passage to homology is a duality preserving functor.

We apply this to $K^\bullet \in C^{\text{perf}}(X)$; then $\text{gr}_i \sigma_\bullet K^\bullet = K^i[i]$ and hence $\text{gr}_i(\pi_* \sigma_\bullet K^\bullet) = \pi_* K^i[i]$.

Lemma 2.5. *Let X be a graded k -vector space with a finite decreasing filtration*

$$X \supset \dots \supset X_\bullet \supset X_{\bullet+1} \supset \dots$$

Suppose $X \otimes X \rightarrow k$ is a perfect symmetric bilinear pairing, which is compatible with the filtration in the sense of (3) and (5). Let X^i denote the i th graded subspace of X and X_\bullet^i denote the i th graded subspace of X_\bullet . Then in $\text{GW}(k)$, there is equality

$$\sum_i (-1)^i [X^i] = \sum_i (-1)^i [\text{gr}_\bullet X^i].$$

Proof. Note that (5) implies that the pairing $\text{gr}_\bullet X$ is non-degenerate, so the statement makes sense (recall Remark 2.2). On any graded symmetric bilinear form, the degree i and $-i$ part for $i \neq 0$ assemble into a metabolic space, with Grothendieck-Witt class determined by the rank (see Lemma B.2). It is clear that the ranks on both sides of our equation are the same; hence it suffices to prove the Lemma in the case where $X^i = 0$ for $i \neq 0$. We may thus ignore the gradings.

Let N be maximal with the property that $X_N \neq 0$. We have a perfect pairing

$$X_{N+1} \otimes X/X_{-n-N} \rightarrow k.$$

Since $X_{N+1} = 0$ we deduce that $X_{-n-N} = X$ and hence $X_j = X$ for all $j \leq -n - N$. If $-n - N \geq N$ then $X = X_N(N)$ and there is nothing to prove; hence assume the opposite.

We have the perfect pairing

$$X_N/X_{N+1} \otimes X_{-n-N}/X_{-n-N+1} \simeq X_N \otimes X/X_{-n-N+1} \rightarrow k.$$

Pick a sequence of subspaces $X \supset X'_{-n-N+1} \supset \dots \supset X'_{N-1}$ such that $X'_i \subset X_i$ and the canonical projection $X'_i \rightarrow X_i/X_N$ is an isomorphism. Extend the filtration X' by zero on the left and constantly on the right. By construction, $X'_i{}^{\text{gr}} = X_i{}^{\text{gr}}$ for $i \neq N, -n - N$, and the pairing on $X' \subset X$ is perfect in the filtered sense. By [MH73, Lemma I.3.1], we have $X = X' \oplus (X')^\perp$. By induction on N , we have $[X'] = [\text{gr}_\bullet X']$. It thus suffices to show that $[(X')^\perp] = [\text{gr}_N X \oplus \text{gr}_{-n-N} X]$. This holds since both sides are metabolic of the same rank: X_{-n-N} is an isotropic subspace of half rank on either side (see again Lemma B.2). \square

Lemma 2.6. *Let E^\bullet be a chain complex with a non-degenerate, symmetric bilinear form $E^\bullet \otimes E^\bullet \rightarrow k[0]$. Then*

$$\sum_i (-1)^i [H^i(E)] = \sum_i [E^i] \in \text{GW}(k).$$

Proof. Since passing to homology is a duality preserving functor, the statement makes sense. Both sides have the same rank, so it suffices to prove equality in $\text{W}(k)$ (see Lemma B.2). We have a perfect pairing $C^i \otimes C^{-i} \rightarrow k$ and similarly for homology. Both are metabolic unless $i = 0$. We can choose a splitting

$$C^0 = H \oplus C';$$

here $H \subset \ker(C^0 \rightarrow C^1)$ maps isomorphically to $H^0(C)$. The restriction of the pairing on C^0 to H is perfect by construction, and hence $C^0 = H \oplus H^\perp$. It suffices to show that H^\perp is metabolic. Compatibility of the pairing with the differential shows that $d(C^{-1}) \subset C^0$ is an isotropic subspace. Self-duality shows that

$$\text{im}(d : C^{-1} \rightarrow C^0) \simeq \text{im}(d^\vee : (C^1)^\vee \rightarrow (C^0)^\vee) \simeq \text{im}(d : C^0 \rightarrow C^1)^\vee$$

which implies that $d(C^{-1}) \subset H^\perp$ is of half rank. This concludes the proof. \square

Proof of Proposition 2.3. Let $K^\bullet = K(V, \sigma)^\bullet \otimes \mathcal{L}$. We compute

$$\begin{aligned} n^{\text{GS}}(V, \rho) &\stackrel{\text{def.}}{=} \sum (-1)^{i+j} [E_1^{i,j}(K^\bullet)] \\ &\stackrel{\text{L.2.6}}{=} \sum (-1)^{i+j} [E_\infty^{i,j}(K^\bullet)] \\ &\stackrel{\text{L.2.5}}{=} \sum_i (-1)^i [R^i f_* K^\bullet] \\ &\stackrel{\text{def.}}{=} n^{\text{GS}}(V, \sigma, \rho). \end{aligned}$$

This is the desired result. \square

Remark 2.7. Admitting a version of Hermitian K -theory which is \mathbb{A}^1 -invariant on regular schemes and has proper pushforwards, one can give an alternative proof of Proposition 2.3 by considering the Koszul complex with respect to the section $t\sigma$ on $\mathbb{A}^1 \times X$. While we believe such a theory exists, at the time of writing there is no reference for this in characteristic 2, so we chose to present the above argument instead.

2.2. Local indices for $n^{\text{GS}}(V, \sigma, \rho)$. Suppose that σ is a section with only isolated zeros. Let i denote the closed immersion $i : Z = Z(\sigma) \hookrightarrow X$ given by the zero locus of σ . We express $n^{\text{GS}}(V, \sigma, \rho)$ as a sum over the points z of Z of a local index at z . To do this, we use a pushforward in a suitable context and show that $\beta_{(V, \sigma)}$ is a pushforward from Z .

For a line bundle \mathcal{L} on a scheme X , denote by $\text{BL}_{\text{naive}}(D(X), \mathcal{L}[n])$ the set of isomorphism classes of non-degenerate symmetric bilinear forms on the derived category of perfect complexes on X , with respect to the duality $\underline{\text{Hom}}(-, \mathcal{L}[n])$. For a proper, lci map $f : X' \rightarrow X$, coherent duality supplies us with a *trace map* $\eta_{f, \mathcal{L}} : f_* f^! (\mathcal{L}) \rightarrow \mathcal{L}$. We can use this to build a pushforward (see [CH09, Theorem 4.2.9])

$$f_* : \text{BL}_{\text{naive}}(D(X'), f^! \mathcal{L}) \rightarrow \text{BL}_{\text{naive}}(D(X), \mathcal{L}),$$

$$[E \otimes E \xrightarrow{\phi} f^! \mathcal{L}] \mapsto [f_* E \otimes f_* E \rightarrow f_*(E \otimes E) \xrightarrow{f_* \phi} f_*(f^! \mathcal{L}) \xrightarrow{\eta_{f, \mathcal{L}}} \mathcal{L}].$$

Remark 2.8. There is a canonical weak equivalence $f^! \mathcal{L} \simeq f^! \mathcal{O}_X \otimes f^* \mathcal{L}$ and $\eta_{f, \mathcal{L}}$ is given by the composition

$$f_*(f^! \mathcal{L}) \simeq f_*(f^! \mathcal{O}_X \otimes f^* \mathcal{L}) \simeq f_* f^! \mathcal{O}_X \otimes \mathcal{L} \xrightarrow{\eta_{f, \mathcal{L}}} \mathcal{L},$$

where $\eta_f = \eta_{f, \mathcal{O}_S}$ [Sta18, Lemma 47.17.8].

Example 2.9. Consider the case of a relatively oriented vector bundle V on a smooth proper variety $f : X \rightarrow \text{Spec}(k)$. Note that elements of $\text{BL}^{\text{naive}}(k)$ are just isomorphism classes of symmetric bilinear forms on graded vector spaces. The orientation supplies us with an equivalence

$$f^!(\mathcal{O}_k) \simeq \omega_{X/k}[n] \simeq \det V^*[n] \otimes \mathcal{L}^{\otimes 2}.$$

Under the induced pushforward map we have

$$f_*[\beta_{(V, \sigma, \rho)}] = n^{\text{GS}}(V, \sigma, \rho) \in \text{BL}^{\text{naive}}(k),$$

where $\beta_{(V, \sigma, \rho)} \in \text{BL}^{\text{naive}}(X, \det V^*[n] \otimes \mathcal{L}^{\otimes 2})$ is the form on $K(V, \sigma) \otimes \mathcal{L}$ defined in §2.1.

Remark 2.10. A symmetric bilinear form ϕ on the derived category $D(S)$ is usually not a very sensible notion. We offer three ways around this.

- (1) If $1/2 \in S$, we could look at the image of ϕ in the Balmer-Witt group of S .
- (2) If ϕ happens to be concentrated in degree zero, it corresponds to a symmetric bilinear form on a vector bundle on S , which is a sensible invariant.
- (3) If $S = \text{Spec}(k)$ is the spectrum of a field, then $D(S)$ is equivalent to the category of graded vector spaces, and we can split ϕ into components by degree and consider

$$cl(\phi) := [H^0(\phi)] + \sum_{i>0} (-1)^i [H^i(\phi) \oplus H^{-i}(\phi)] \in \text{GW}(k).$$

Let 1_Z denote the element of $\text{BL}_{\text{naive}}(D(Z), \mathcal{O}_Z[0])$ represented by $\mathcal{O}_Z \otimes \mathcal{O}_Z \rightarrow \mathcal{O}_Z$.

Proposition 2.11. *Let X be a scheme, V a vector bundle, and $\sigma \in \Gamma(X, V)$ a section locally given by a regular sequence. Write $i : Z = Z(\sigma) \hookrightarrow X$ for the inclusion of the zero scheme. Proposition B.1 yields a canonical equivalence $i^! \det(V^*)[n] \simeq \mathcal{O}_Z[0]$, where n is the rank of V ; under the induced map*

$$i_* : \text{BL}_{\text{naive}}(D(Z), \mathcal{O}_Z[0]) \rightarrow \text{BL}_{\text{naive}}(D(X), \det(V^*)[n])$$

we have $i_*(1_Z) = \beta_{(V, \sigma)}$, where

$$\beta_{(V, \sigma)} : K(V, \sigma) \otimes K(V, \sigma) \rightarrow \det(V^*)[n]$$

is the canonical pairing on the Koszul complex as in §2.1.

Proof. Because σ locally corresponds to a regular sequence, the canonical map $r : K(V, \sigma)^\bullet \rightarrow i_* \mathcal{O}_Z$ is an equivalence in $D(X)$. The canonical projection $i_* \mathcal{O}_Z \simeq K(V, \sigma) \rightarrow \det(V^*)[n]$ induces by adjunction a map $\mathcal{O}_Z \rightarrow i^! \det(V^*)[n]$. We claim that this is the equivalence of Proposition B.1. The proof of said proposition shows that the problem is local on Z , so we may assume that V is trivial. Then this map is precisely the isomorphism constructed in [Har66, Proposition III.7.2 and preceding pages], which is also the isomorphism employed in the proof of Proposition B.1.

Now we prove that $i_*(1_Z) = \beta_{(V,\sigma)}$. Consider the following diagram

$$\begin{array}{ccc} K(V,\sigma) \otimes K(V,\sigma) & \xrightarrow{r \otimes r} & i_* \mathcal{O}_Z \otimes^L i_* \mathcal{O}_Z \\ m_K \downarrow & & m_Z \downarrow \\ K(V,\sigma) & \xrightarrow{r} & i_* \mathcal{O}_Z \simeq i_* i^! \det(V^*)[n] \xrightarrow{\text{tr}} \det(V^*)[n]. \end{array}$$

The map $m_K : K(V,\sigma) \otimes K(V,\sigma) \rightarrow K(V,\sigma)$ is the canonical multiplication (see (2) §2.1), and $m_Z : i_* \mathcal{O}_Z \otimes^L i_* \mathcal{O}_Z \rightarrow i_* \mathcal{O}_Z \otimes i_* \mathcal{O}_Z \rightarrow i_* \mathcal{O}_Z$ is equivalently given by either the multiplication in \mathcal{O}_Z or the lax monoidal witness transformation of i_* . The former interpretation shows that the left hand square commutes. The pairing $i_*(1)$ is given by the composite from the top right hand corner to the bottom right hand corner. To prove the claim it suffices to show that the bottom row composite $K(V,\sigma) \rightarrow \det(V^*)[n]$ is the canonical projection. This follows by adjunction from our choice of equivalence $\mathcal{O}_Z \simeq i^! \det(V^*)[n]$.

This concludes the proof. \square

Proposition 2.11 is an example of a more general phenomenon given in Meta-Theorem 3.9.

Lemma 2.12 ([CH09], Theorem 5.1.9). *Let $g : Z \rightarrow Y$ and $f : Y \rightarrow X$ be proper maps.⁴ Given equivalences $f^! \mathcal{L} \simeq \mathcal{M}[n]$ and $g^! \mathcal{M}[n] \simeq \mathcal{N}$, the canonical equivalence $(fg)^! \simeq g^! f^!$ produces a weak equivalence $(fg)^! \mathcal{L} \simeq \mathcal{N}$, and consequently push forward maps*

$$\begin{aligned} g_* &: \text{BL}_{\text{naive}}(D(Z), \mathcal{N}) \rightarrow \text{BL}_{\text{naive}}(D(Y), \mathcal{M}[n]) \\ f_* &: \text{BL}_{\text{naive}}(D(Y), \mathcal{M}[n]) \rightarrow \text{BL}_{\text{naive}}(D(X), \mathcal{L}) \\ (fg)_* &: \text{BL}_{\text{naive}}(D(Z), \mathcal{N}) \rightarrow \text{BL}_{\text{naive}}(D(X), \mathcal{L}). \end{aligned}$$

There is a canonical equivalence $(fg)_* \simeq f_* g_*$.

Proof. The main point is that $\eta_{f,\mathcal{L}} \circ f_*(\eta_{g,\mathcal{M}[n]}) = \eta_{fg,\mathcal{L}}$. The categorical details are worked out in the reference. \square

Now we get back to our Euler numbers. Let X/k be smooth and proper, V a relatively oriented vector bundle, σ a section of V with only isolated zeros. Write $i : Z = Z(\sigma) \hookrightarrow X$ for the inclusion of the zero scheme of σ . Let $\varpi : Z \rightarrow \text{Spec } k$ and $f : X \rightarrow \text{Spec } k$ denote the structure maps, so that $\varpi = fi$.

The weak equivalence $i^! \det(V^*)[n] \simeq \mathcal{O}_Z[0]$ of Proposition 2.11 and Remark 2.8 produce a weak equivalence $i^!(\det V^*[n] \otimes \mathcal{L}^{\otimes 2}) \cong i^* \mathcal{L}^{\otimes 2}$. The orientation ρ gives an isomorphism $\det V^*[n] \otimes \mathcal{L}^{\otimes 2} \cong \omega_{X/k}[n]$. Combining, we have a chosen weak equivalence

$$i^!(\omega_{X/k}[n]) \cong i^* \mathcal{L}^{\otimes 2}.$$

Since also $f^! \mathcal{O}_k \simeq \omega_{X/k}[n]$ (see e.g. Proposition B.1), we therefore obtain a canonical equivalence

$$\varpi^! \mathcal{O}_k \simeq i^* \mathcal{L}^{\otimes 2}.$$

We use this equivalence to define

$$\varpi_* : \text{BL}^{\text{naive}}(Z, i^* \mathcal{L}^{\otimes 2}) \rightarrow \text{BL}^{\text{naive}}(k).$$

Corollary 2.13. *With this notation, we have*

$$n^{\text{GS}}(V, \sigma, \rho) = \varpi_*(i^* \mathcal{L} \otimes i^* \mathcal{L} \rightarrow i^* \mathcal{L}^{\otimes 2}).$$

Proof. By Lemma 2.12 we have $\varpi_* = f_* i_*$. Proposition 2.11 and the projection formula imply that $i_*(i^* \mathcal{L} \otimes i^* \mathcal{L} \rightarrow i^* \mathcal{L}^{\otimes 2}) = \beta_{V,\sigma,\rho}$. We conclude by Example 2.9. \square

Suppose that σ has isolated zeros, or in other words that the support of σ is a disjoint union of points. Then $n^{\text{GS}}(V, \sigma, \rho)$ can be expressed as a sum of local contributions. Namely, for each point z of Z , let $i_z : Z_z \hookrightarrow X$ denote the chosen immersion coming from the connected component of Z given by z . Let $\varpi_z : Z_z \rightarrow \text{Spec } k$ denote the structure map. Then

$$n^{\text{GS}}(V, \sigma, \rho) = \sum_{z \in Z} \varpi_{z*}(i_z^* \mathcal{L} \otimes i_z^* \mathcal{L} \rightarrow i_z^* \mathcal{L}^{\otimes 2}).$$

In light of this we propose the following.

Definition 2.14. For a relatively oriented vector bundle with a section as above, and $z \in Z(\sigma)$, we define

$$\text{ind}_z(\sigma) = \text{ind}_z(V, \sigma, \rho) = \varpi_{z*}(i_z^* \mathcal{L} \otimes i_z^* \mathcal{L} \rightarrow i_z^* \mathcal{L}^{\otimes 2}) \in \text{BL}^{\text{naive}}(k).$$

⁴Recall our convention that since we are invoking a functor $f^!$, Z, Y, X are also finite type and separated over a noetherian base S . Without this we should add the hypothesis that f, g are locally (so globally) of finite presentation.

The above formula then reads

$$(3) \quad n^{\text{GS}}(V, \sigma, \rho) = \sum_{z \in Z} \text{ind}_z(V, \sigma, \rho).$$

In the next two subsections, we compute the local contributions $\text{ind}_z(\sigma)$ as an explicit bilinear form constructed by Scheja and Storch [SS75], appearing in the Eisenbud–Levine–Khimshiashvili signature theorem [EL77] [Khi77], and used as the local index of the Euler class constructed in [KW21, Section 4].

2.3. Scheja–Storch and coherent duality. Let S be a scheme, $\pi : X \rightarrow S$ a smooth scheme of relative dimension n , and $Z \subset X$ closed with $\varpi : Z \rightarrow S$ finite. Suppose given the following data:

- (1) Sections $T_1, \dots, T_n \in \mathcal{O}(X)$ such that $T_i \otimes 1 - 1 \otimes T_i$ generate the ideal of $Z \subset X \times_S X$.
- (2) Sections $f_1, \dots, f_n \in \mathcal{O}(X)$ such that $Z = Z(f_1, \dots, f_n)$.

Remark 2.15. Since $Z \rightarrow X$ is quasi-finite, Lemma B.5 shows that f_1, \dots, f_n is a regular sequence and $Z \rightarrow X$ is flat, so finite locally free (being finite and finitely presented [Sta18, Tag 02KB]).

Choose $a_{ij} \in \mathcal{O}(X \times_S X)$ such that

$$f_i \otimes 1 - 1 \otimes f_i = \sum_j a_{ij}(T_j \otimes 1 - 1 \otimes T_j).$$

Let $\Delta \in \mathcal{O}(Z \times_S Z)$ be the image of the determinant of a_{ij} . Since ϖ is finite locally free, Δ determines an element $\tilde{\Delta}$ of

$$\text{Hom}_{\mathcal{O}_S}(\mathcal{O}_S, (\varpi \times_S \varpi)_* \mathcal{O}_{Z \times_S Z}) \simeq \text{Hom}_{\mathcal{O}_S}(\mathcal{O}_S, \varpi_* \mathcal{O}_Z \otimes \varpi_* \mathcal{O}_Z) \simeq \text{Hom}_{\mathcal{O}_S}((\varpi_* \mathcal{O}_Z)^*, \varpi_* \mathcal{O}_Z).$$

Remark 2.16. We can make $\tilde{\Delta}$ explicit: if $\Delta = \sum_i b_i \otimes b'_i$, then

$$\tilde{\Delta}(\alpha) = \sum_i \alpha(b_i) b'_i.$$

Remark 2.17. By construction, the pullback of Δ along the diagonal $\delta : Z \rightarrow Z \times_S Z$ is the determinant of the differentiation map $C_{Z/X} \rightarrow \Omega_X|_Z$ with respect to the canonical bases. In other words this is the *Jacobian*:

$$\delta^*(\Delta) = \text{Jac}^F := \det \left(\frac{\partial f_i}{\partial T_j} \right)_{i,j=1}^n.$$

Theorem 2.18. *Under the above assumptions, the map*

$$\tilde{\Delta} : (\varpi_* \mathcal{O}_Z)^* \rightarrow \varpi_* \mathcal{O}_Z$$

is a symmetric isomorphism and hence determines a symmetric bilinear structure on $\varpi_ \mathcal{O}_Z$. This is the same structure as $\varpi_*(1)$, i.e.*

$$\varpi_*(\mathcal{O}_Z) \otimes \varpi_*(\mathcal{O}_Z) \rightarrow \varpi_*(\mathcal{O}_Z) \simeq \varpi_*(\varpi^! \mathcal{O}_S) \xrightarrow{\eta_\varpi} \mathcal{O}_S.$$

Here the isomorphism $\varpi^! \mathcal{O}_Z \simeq \mathcal{O}_Z$ arises from

$$\varpi^!(\mathcal{O}_Z) \simeq \widetilde{\det} L_\varpi \simeq \omega_{Z/X} \otimes \omega_{X/S} \simeq \mathcal{O},$$

with the first isomorphism given by Proposition B.1, and the third isomorphism given by the sections (T_i) and (f_i) .

Remark 2.19. The theorem asserts in particular that the isomorphism $\tilde{\Delta}$, and hence the section Δ , are independent of the choice of the a_{ij} .

We begin with some preliminary observations before delving into the proof. The problem is local on S , so we may assume that $S = \text{Spec}(A)$; then $Z = \text{Spec}(B)$. Since ϖ is finite, there is a canonical isomorphism [Har66, III §8 Theorem 8.7 (3), or Ideal Theorem (3) p. 6]

$$\varpi^! \simeq \underline{\text{Hom}}_A(B, -) : D(A) \rightarrow D(B).$$

In particular

$$\varpi^!(A) \simeq \underline{\text{Hom}}_A(B, A)$$

and the trace map takes the form [Har66, Ideal theorem 3) pg 7]

$$\varpi_* \varpi^! A \simeq \text{Hom}_A(B, A) \rightarrow A, \eta \mapsto \eta(1).$$

Proof of Theorem 2.18. The isomorphisms

$$B^* = \underline{\mathrm{Hom}}_A(B, A) \simeq \varpi^!(A) \simeq B$$

determine an element $\Delta' \in \mathrm{Hom}_A(B^*, B)$. The theorem is equivalent to showing that $\tilde{\Delta} = \Delta'$.

We thus need to make explicit the isomorphism

$$\underline{\mathrm{Hom}}_A(B, A) \simeq \varpi^!(\mathcal{O}_A) \simeq i^! \pi^!(\mathcal{O}_A) \simeq \omega_{Z/X} \otimes \omega_X \simeq \mathcal{O}.$$

Tracing through the definitions (including the proof of [Har66, III Proposition 8.2]), one finds that this isomorphism arises by computing

$$\mathrm{Ext}_X^n(B, \mathcal{O}_X)$$

in two ways. On the one hand, the kernel of the surjection $\mathcal{O}_X \rightarrow B$ is generated by f_1, \dots, f_n , which is a regular sequence by Remark 2.15; let $K_A(f)^\bullet$ denote the corresponding Koszul complex. On the other hand we can consider the embedding $Z \hookrightarrow X \times Z$; its ideal is generated by the strongly regular sequence $T_i - t_i$, where t_i is the image of T_i in B . We thus obtain a resolution $K_B(T-t)^\bullet \rightarrow B$ over $X \times Z$. Since $p : X \times Z \rightarrow X$ is finite, $p_* K_B(T-t)^\bullet \rightarrow p_* B = B$ is still a resolution. We shall conflate $K_B(T-t)$ and $p_* K_B(T-t)$ notationally. We can thus compute

$$\mathrm{Ext}_X^n(B, \mathcal{O}_X) \simeq \mathrm{coker}(\mathrm{Hom}_X(K_B(T-t)^{n-1}, \mathcal{O}_X) \rightarrow \mathrm{Hom}_X(K_B(T-t)^n, \mathcal{O}_X)).$$

Since $\mathrm{Hom}_X(B \otimes \mathcal{O}_X, \mathcal{O}_X) \simeq \mathrm{Hom}_A(B, \mathcal{O}_X) \simeq \mathrm{Hom}_A(B, A) \otimes_A \mathcal{O}_X$, there is a natural map $\xi : \mathrm{Hom}_A(B, A) \rightarrow \mathrm{Hom}_X(K_B(T-t)^n, \mathcal{O}_X)$ (sending α to $\alpha \otimes 1$). One checks that this induces $\mathrm{Hom}_A(B, A) \simeq \mathrm{coker}(\dots) \simeq \mathrm{Ext}_X^n(B, \mathcal{O}_X)$.

We can write down a map of resolutions $\zeta : K_A(f) \rightarrow K_B(T-t)$ as follows. The kernel of $B \otimes \mathcal{O}_X \rightarrow B$ is by construction generated by $\{T_i - t_i\}_i$, but it also contains f_i . Note that $f_i = \sum_j \bar{a}_{ij}(T_j - t_j)$, where we write \bar{a}_{ij} for the image of a_{ij} in $\mathcal{O}_X \otimes B$. Letting $K_A(f)$ be the exterior algebra on $\{e_1, \dots, e_n\}$ and $K_B(T-t)$ the exterior algebra on $\{e'_1, \dots, e'_n\}$, the map ζ is specified by $\zeta(e_i) = \sum_j \bar{a}_{ij} e'_j$. The isomorphism

$$\mathrm{Hom}_A(B, A) \simeq h^n \mathrm{Hom}_X(K_B(T-t)^\bullet, \mathcal{O}_X) \simeq h^n \mathrm{Hom}_X(K_A(f)^\bullet, \mathcal{O}_X) \simeq B$$

is thus given by

$$\begin{aligned} \mathrm{Hom}_A(B, A) &\xrightarrow{\xi} \mathrm{Hom}_X(K_B(T-t)^n, \mathcal{O}_X) \xrightarrow{\det(a_{ij})^*} \mathrm{Hom}_X(K_A(f)^n, \mathcal{O}_X) \\ &\simeq \mathrm{Hom}_X(\mathcal{O}_X, \mathcal{O}_X) \simeq \mathcal{O}_X \rightarrow B. \end{aligned}$$

Write the image of $\det(a_{ij})$ in $B \otimes B$ as $\sum_k b_k \otimes b'_k$. Tracing through the definitions, we find that the above composite sends $\alpha \in \mathrm{Hom}_A(B, A)$ to $\sum_k \alpha(b_k) b'_k$. By Remark 2.16, this is precisely $\tilde{\Delta}$.

This concludes the proof. \square

Definition 2.20. If $X = U \subset \mathbb{A}_S^n$, (T_i) are the standard coordinates, and $F = (f_1, \dots, f_n)$, we denote the symmetric bilinear form constructed above by

$$\langle - | - \rangle^{\mathrm{SS}} = \langle - | - \rangle^{\mathrm{SS}}(U, F, S).$$

This form was first constructed, without explicitly using coherent duality, by Scheja and Storch [SS75, 3].

Example 2.21. Suppose that $Z \rightarrow S$ is an isomorphism (where $Z = Z(F)$ as above), so that the diagonal $\delta : Z \rightarrow Z \times_S Z$ is also an isomorphism. Then $\langle - | - \rangle^{\mathrm{SS}}$ is just the rank 1 bilinear form corresponding to multiplication by $\delta^*(\Delta) \in \mathcal{O}_Z \simeq \mathcal{O}_S$. In other words, using Remark 2.17, $\langle - | - \rangle^{\mathrm{SS}}$ identifies with $(x, y) \mapsto (\mathrm{Jac} F)xy$.

2.4. The Poincaré–Hopf Euler number with respect to a section. In this subsection, we recall the Euler class defined in [KW21, Section 4] and prove Theorem 1.1. To distinguish this Euler class from the others under consideration, here we call it the *Poincaré–Hopf Euler number*, because it is a sum of local indices as in the Poincaré–Hopf theorem for the Euler characteristic of a manifold. It is defined using local coordinates.

Let k be a field, and let X be an n -dimensional smooth k -scheme. Let z be a closed point of X .

Definition 2.22. (cf. [KW21, Definition 17]) By a system of *Nisnevich coordinates around z* we mean a Zariski open neighborhood U of z in X , and an étale map $\varphi : U \rightarrow \mathbb{A}_k^n$ such that the extension of residue fields $k(\varphi(z)) \subseteq k(z)$ is an isomorphism.

Proposition 2.23. *When $n > 0$, there exists a system of Nisnevich coordinates around every closed point z of X .*

Proof. When k is infinite, this follows from [Knu91, Chapter 8. Proposition 3.2.1]. When $k(z)/k$ is separable, for instance when k is finite, this is [KW21, Lemma 18]. \square

As above, let V be a relatively oriented, rank n vector bundle on X . Let σ be a section with only isolated zeros, and let $Z \hookrightarrow X$ denote the closed subscheme given by the zero locus of σ . Let z be a point of Z . The Poincaré–Hopf local index or degree

$$\mathrm{ind}_z^{\mathrm{PH}} \sigma \in \mathrm{GW}(k)$$

was defined in [KW21, Definition 30] as follows. Choose a system of Nisnevich coordinates $\varphi : U \rightarrow \mathbb{A}_k^n$ around z . After possibly shrinking U , the restriction of V to U is trivial and we may choose an isomorphism $\psi : V|_U \rightarrow \mathcal{O}_U^n$ of V . The local trivialization ψ induces a distinguished section of $\det V(U)$. The system of local coordinates φ induces a distinguished section of $\det T_X(U)$, and we therefore have a distinguished section of $(\det V \otimes \omega_X)(U)$. As in [KW21, Definition 19], the local coordinates ϖ and local trivialization ψ are said to be *compatible with the relative orientation* if the distinguished element is the tensor square of a section of $\mathcal{L}(U)$. By multiplying ψ by a section in $\mathcal{O}(U)$, we may assume this compatibility.

Under ψ , the section σ can be identified with an n -tuple (f_1, \dots, f_n) of regular functions, $\psi(\sigma) = (f_1, \dots, f_n) \in \oplus_{i=1}^n \mathcal{O}_U$. Let m denote the maximal ideal of \mathcal{O}_U corresponding to z . Since z is an isolated zero, there is an integer n such that $m^n = 0$ in $\mathcal{O}_{Z,z}$. For any N , it is possible to choose (g_1, \dots, g_n) in $\oplus_{i=1}^n m^N$ such that $(f_i + g_i)|_U$ is in the image of $\varphi^* : \mathcal{O}_{\mathbb{A}_k^n} \rightarrow \varphi_* \mathcal{O}_U$ after possibly shrinking U [KW21, Lemma 22]. For $N = 2n$, choose such (g_1, \dots, g_n) and let F_i in $\mathcal{O}_{\mathbb{A}_k^n}(\mathbb{A}_k^n)$ be the functions such that $\varphi^*(F_i) = f_i + g_i$. Then φ induces an isomorphism $\mathcal{O}_{Z,z} \cong k[t_1, \dots, t_n]_{m_{\varphi(z)}} / (F_1, \dots, F_n)$ [KW21, Lemma 25], and $\mathrm{ind}_z^{\mathrm{PH}} \sigma$ is defined to be the associated form of Scheja–Storch $\langle - | - \rangle^{\mathrm{SS}}(\varphi(U), F, k)$ (see §2.3 and Definition 2.20 for the definition of $\langle - | - \rangle^{\mathrm{SS}}(\varphi(U), F, k)$). The local index $\mathrm{ind}_z^{\mathrm{PH}} \sigma$ is well-defined by [KW21, Lemma 26]. Then the Poincaré–Hopf Euler number is defined to be the sum of the local indices:

Definition 2.24. The *Poincaré–Hopf Euler number* $n^{\mathrm{PH}}(V, \sigma)$ of V with respect to σ is $n^{\mathrm{PH}}(V, \sigma) = \sum_{z \text{ such that } \sigma(z)=0} \mathrm{ind}_z^{\mathrm{PH}} \sigma$.

Proof of Theorem 1.1. By Proposition 2.3 we have $n^{\mathrm{GS}}(V) = n^{\mathrm{GS}}(V, \sigma)$, where the orientation has been suppressed from the notation, but is indeed present. Using Formula (3), it is thus enough to show that $\mathrm{ind}_z(\sigma) = \mathrm{ind}_z^{\mathrm{PH}}(\sigma)$. This follows from Theorem 2.18. One needs to be careful about the trivializations used in defining the various pushforward maps; this is ensured precisely by the condition that the tautological section is a square. The details of this argument are spelled out more carefully in the proof in Proposition 3.13 in the next section. \square

One can extend the comparison of local degrees $\mathrm{ind}_z^{\mathrm{PH}} \sigma = \mathrm{ind}_z \sigma$ to work over a more general base scheme S . This was done for $S = \mathbb{A}_k^1$ with k a field in [KW21, Lemma 33], but in more generality, it is useful to pick the local coordinates using knowledge of both σ and X as follows.

Definition 2.25. Let X be a scheme, V a vector bundle on X and σ a section of V .

- (1) We call σ *non-degenerate* if it locally corresponds to a regular sequence.
- (2) Given another scheme S and a morphism $\pi : X \rightarrow S$, we call σ *very non-degenerate* (with respect to π) if it non-degenerate and the zero locus $Z(\sigma)$ is finite and locally free over S .

Remark 2.26. Suppose that X is smooth over S and $\mathrm{rk}(V) = \dim X/S$.

- (1) If $S = \mathrm{Spec}(k)$ is the spectrum of a field, then $Z(\sigma) \rightarrow \mathrm{Spec}(k)$ is quasi-finite if and only if it is finite locally free, if and only if σ is locally given by a regular sequence. In other words σ is non-degenerate if and only if it is very non-degenerate, if and only if $Z(\sigma) \rightarrow Z$ is quasi-finite.
- (2) In general, σ is non-degenerate as soon as $Z(\sigma) \rightarrow S$ is quasi-finite, and very non-degenerate if and only if $Z(\sigma) \rightarrow S$ is finite. See Lemma B.5.

Example 2.27. If σ is a non-degenerate section, then precomposition with σ induces an isomorphism $\mathrm{Hom}(V, \mathcal{O}) \simeq C_{Z/X}$. In particular $N_{Z/X} \simeq V$ and $L_{Z/X} \simeq V^*[1]$.

Definition 2.28. Let X be a smooth S -scheme, and let $V \rightarrow X$ be a vector bundle, relatively oriented by ρ . Let σ be a very non-degenerate section of V , and let Z be a closed and open subscheme of the zero locus $Z(\sigma)$ of σ . By a *system of coordinates for (V, X, σ, ρ, Z)* we mean an open neighbourhood U of Z in X , an étale map $\varphi : U \rightarrow \mathbb{A}_S^n$, a trivialization $\psi : V|_U \simeq \mathcal{O}_U^n$, and a section $\sigma' \in \mathcal{O}_{\mathbb{A}_S^n}(\varphi(U))$, such that the following conditions hold:

- (1) $Z = Z(\sigma|_U) \simeq Z(\sigma')$,

- (2) $\det(\sigma|_Z) = \det(\varphi^* \sigma'|_Z) \in \det N_{Z/X}$, and
- (3) the canonical section of $\omega_{X/S} \otimes \det V|_Z \cong \mathcal{L}^{\otimes 2}|_Z$ determined by ψ and φ corresponds to the square of a section of $\mathcal{L}|_Z$.

Here for (2) and (3) we used Example 2.27.

Suppose that X is dimension n over S , so that the rank of V is also n . Let $Z \subset Z(\sigma)$ be a clopen component and write $\varpi : Z \rightarrow S$ for the structure map. The local index generalizes straightforwardly from Definition 2.14:

Definition 2.29. We call

$$\text{ind}_Z(\sigma) = \text{ind}_Z(V, \sigma, \rho) = \varpi_*(i^* \mathcal{L} \otimes i^* \mathcal{L} \rightarrow i^* \mathcal{L}^{\otimes 2}) \in \text{BL}^{\text{naive}}(S)$$

the local index at Z .

Remark 2.30. Since ϖ is finite locally free, ϖ_* preserves vector bundles. In particular, $\text{ind}_Z(\sigma) \in \text{BL}^{\text{naive}}(S)$ is a symmetric bilinear form on a vector bundle, as opposed to just on a complex up to homotopy. (See also Remark 2.10.)

A system of coordinates for (V, X, σ, ρ, Z) determines a presentation $Z = Z(\sigma|_U) = Z(\sigma') \subset \mathbb{A}_S^n$, where $\sigma' : \mathbb{A}_S^n \supset \phi(U) \rightarrow \mathbb{A}^n$. Hence Definition 2.20 supplies us with a symmetric bilinear form $\langle -|-\rangle^{\text{SS}}(\varphi(U), \sigma, S) \in \text{BL}^{\text{naive}}(S)$.

Proposition 2.31. *The form $\langle -|-\rangle^{\text{SS}}(\varphi(U), \sigma', S)$ coincides (up to isomorphism) with $\text{ind}_Z(\sigma)$. In particular its isomorphism class is independent of the choice of coordinates.*

Proof. The argument is the same as in the proofs of Theorem 1.1 and Proposition 3.13. \square

In contrast, our proof that the Euler number (sum of indices) is independent of the choice of section (i.e. Proposition 2.3) does not generalize immediately; in fact this will not hold in $\text{BL}^{\text{naive}}(S)$ but rather in some quotient (like $\text{GW}(k)$ in the case of fields). As indicated in Remark 2.7, one situation in which it is easy to see this independence is if the quotient group satisfies homotopy invariance. This suggests studying Euler numbers valued in more general homotopy invariant cohomology theories for algebraic varieties, which is what the remainder of this work is concerned with.

3. COHOMOLOGY THEORIES FOR SCHEMES

3.1. Introduction. In order to generalize the results from the previous sections, we find it useful to introduce the concept of a *cohomology theory twisted by K -theory*. We do not seek here to axiomatize all the relevant data, but just introduce a common language for similar phenomena.

Definition 3.1. Let S be a scheme and $\mathcal{C} \subset \text{Sch}_S$ a category of schemes. Denote by \mathcal{C}^L the category of pairs (X, ξ) where $X \in \mathcal{C}$ and $\xi \in K(X)$ (i.e. a point in the K -theory space of X), and morphisms those maps of schemes compatible with the K -theory points.⁵ By a *cohomology theory E* over S (for schemes in \mathcal{C}) we mean a presheaf of sets on \mathcal{C}^L , i.e. a functor

$$E : (\mathcal{C}^L)^{\text{op}} \rightarrow \text{Set}, (X, \xi) \mapsto E^\xi(X).$$

To illustrate the flavor of cohomology theory we have in mind, we begin with two examples.

Example 3.2. We can put

- (1) $E^\xi(X) = \text{CH}^{r^{k(\xi)}}(X)$, the Chow group of algebraic cycles up to rational equivalence of the appropriate codimension, or
- (2) $E^\xi(X) = \text{GW}(X, \widetilde{\det} \xi)$, the Grothendieck-Witt group of symmetric bilinear perfect complexes for the duality $\underline{\text{Hom}}(-, \widetilde{\det} \xi)$ (see e.g. [Sch10]).

Warning 3.3. For cohomology theories with values in a 1-category (like sets), in the above definition we can safely replace $K(X)$ by its truncation $K(X)_{\leq 1}$, i.e. the ordinary 1-groupoid of virtual vector bundles. However, we can in general *not* replace it by just the set $K_0(X)$. In other words, if (say) V is a vector bundle on X and ϕ an automorphism of V , then there is an induced automorphism

$$E(\phi) : E^V(X) \rightarrow E^V(X)$$

which may or may not be trivial. For example, in the case $E = \text{GW}$ as above, if $V = \mathcal{O}$ is trivial and ϕ corresponds to $a \in \mathcal{O}^\times(X)$, then $E(\phi)$ is given by multiplication by $\langle a \rangle \in \text{GW}(X)$.

⁵Technically speaking, this means “coherently compatible”, so \mathcal{C}^L is an ∞ -category. However we will only need its homotopy 1-category, so for us compatible means “together with a homotopy class of paths joining the two K -theory points”.

3.2. Features of cohomology theories. Many cohomology theories that occur in practice satisfy additional properties beyond the basic ones of the above definition, and many come with more data. We list here some of those relevant to the current paper.

Morphisms of theories: Cohomology theories form a category in an evident way, with morphisms given by natural transformations.

Trivial bundles: We usually abbreviate $E^{\mathcal{O}^n}(X)$ to $E^n(X)$.

Additive and multiplicative structure: Often, E takes values in abelian groups. Moreover, often $E^0(X)$ is a ring and $E^\xi(X)$ is a module over $E^0(X)$. Typically all of this structure is preserved by the pullback maps.

Disjoint unions: Usually E converts finite disjoint unions into products, i.e. $E(\emptyset) = *$ and $E(X \coprod Y) = E(X) \times E(Y)$. If E takes values in abelian groups, this is usually written as $E(X \coprod Y) = E(X) \oplus E(Y)$.

Orientations: In many cases, the cohomology theory E factors through a quotient of the category \mathcal{C}^L , built using a quotient $q : K(X) \rightarrow K'(X)$ of the K -theory groupoid. In other words, one has canonical isomorphisms $E^\xi(X) \simeq E^{\xi'}(X)$ for certain K -theory points ξ, ξ' . More specifically:

GL-orientations: In the above situation, if $K'(X) = \mathbb{Z}$ and q is the rank map, then we speak of a GL-orientation. In other words, in this situation we canonically have $E^\xi(X) \simeq E^{rk(\xi)}(X)$. In particular, Warning 3.3 does not apply: all automorphisms of vector bundles act trivially on E . This happens for example if $E = \text{CH}$ (see Example 3.2(1)).

SL-orientations: If instead $K'(X) = \text{Pic}(D(X))$ via the determinant, then we speak of an SL-orientation. In other words, in this situation $E^\xi(X)$ only depends on the rank and (ungraded) determinant of ξ . We write $E^{rk(\xi)}(X, \det(\xi))$ for this common group. This happens for example if $E = \text{GW}$ (see Example 3.2(1)).

SL^c-orientations: This is a further strengthening of the concept of an SL-orientation, where in $K'(X) = \text{Pic}(D(X))$ we mod out (in the sense of groupoids) by the squares of line bundles. In other words, if $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3$ are line bundles on X , then any isomorphism $\mathcal{L}_1 \simeq \mathcal{L}_2 \otimes \mathcal{L}_3^{\otimes 2}$ induces

$$E^n(X, \mathcal{L}_1) \simeq E^n(X, \mathcal{L}_2).$$

Note that in particular then $E^n(X, \mathcal{L}) \simeq E^n(X, \mathcal{L}^*)$. This also happens for $E = \text{GW}$, essentially by construction.

Supports: Often, for $Z \subset X$ closed there is a cohomology with support, denoted $E_Z^\xi(X)$. It enjoys further functorialities which we do not list in detail here.

Transfers: In many theories, for appropriate morphisms $p : X \rightarrow Y$ and $\xi \in K(Y)$, there exists $tw(p, \xi) \in K(X)$ and a transfer map

$$p_* : E^{tw(p, \xi)}(X) \rightarrow E^\xi(Y),$$

compatible with composition. Typically p is required to be lci, and

$$tw(p, \xi) = p^*\xi + L_p,$$

where L_p is the *cotangent complex* [Ill71]. Furthermore, typically p is required to be proper, or else we need to fix $Z \subset X$ closed and proper over Y and obtain $p_* : E_Z^{tw(p, \xi)}(X) \rightarrow E^\xi(Y)$. Finally, usually E takes values in abelian groups and satisfies the disjoint union property, and transfer from a disjoint union is just the sum of the transfers.

Remark 3.4. (1) We have defined a morphism of cohomology theories as a natural transformation of functors valued in sets. Whether or not such a transformation respects additional structure (abelian group structures, orientations, transfers etc.) must be investigated in each case.

(2) In many cases (in particular in the presence of homotopy invariance), SL-oriented theories are also canonically SL^c-oriented. See Proposition 4.19.

3.3. Some cohomology theories. We now introduce a number of cohomology theories that can be used in this context.

Hermitian K -theory GW: This is the theory from Example 3.2(2). It is SL^c-oriented. We believe that it has transfers for (at least) smooth proper morphisms and regular immersions, but we are not aware of a reference for this in adequate generality. If X is regular and $1/2 \in X$ one can use the comparison with KO-theory (see below).

Naive derived bilinear forms BL_{naive}: See §2.2.

Cohomology theories represented by motivic spectra: Let $\mathcal{SH}(S)$ denote the motivic stable ∞ -category. Then any $E \in \mathcal{SH}(S)$ defines a cohomology theory on Sch_S , automatically satisfying many good properties; for example they always have transfers along smooth and proper morphisms, as well as regular immersions. For a lucid introduction, see [EHK⁺20a]. We recall some of the main points in §4.

Orthogonal K -theory spectrum KO: This spectrum is defined and stable under arbitrary base change if $1/2 \in S$ [PW18, ST15]. Over regular bases, it represents Hermitian K -theory GW; in general it represents a homotopy invariant version.

Generalized motivic cohomology $H\mathbb{Z}$: This can be defined as $\pi_0^{\text{eff}}(\mathbb{1})$; see for example [Bac17]. Over fields (of characteristic not 2) it represents generalized motivic cohomology in the sense of Calmès-Fasel [CF17b, CF17a, BF17]; it is unclear if this theory is useful in this form over more general bases.

3.4. The yoga of Euler numbers. Let E be a cohomology theory.

Definition 3.5. We will say that E has Euler classes if, for each scheme X over S and each vector bundle V on X we are supplied with a class

$$e(V, E) \in E^{V^*}(X).$$

Remark 3.6. The twist by V^* (instead of V) in the above definition may seem peculiar. This ultimately comes from our choice of covariant (instead of contravariant) equivalence between locally free sheaves and vector bundles, whereas a contravariant equivalence is used in the motivic Thom spectrum functor and hence in the definition of twists.

Typically, the Euler classes will satisfy further properties, such as stability under base change; we do not formalize this here.

Now suppose that $\pi : X \rightarrow S$ is smooth and proper, V is relatively oriented, E has transfers for smooth proper maps and is SL^c -oriented. In this case we have a transfer map

$$E^{V^*}(X) \simeq E^n(X, \det V) \xrightarrow{\rho} E^n(X, \omega_{X/S}) \simeq E^{L_\pi}(X) \xrightarrow{\pi_*} E^0(S).$$

Definition 3.7. In the above situation, we call

$$n(V, \rho, E) = \pi_* e(V, E)$$

the *Euler number* of V in E with respect to the relative orientation ρ .

Example 3.8. Let $E = \text{GW}$. We can define a family of Euler classes by

$$e(V) = [K(V, 0)] \in \text{GW}(X, \widetilde{\det V}) \simeq \text{GW}^{V^*}(X);$$

here we use the Koszul complex from §2.1. This depends initially on a choice of section, but we shall show that the Grothendieck-Witt class often does not. In any case, here we chose the 0-section for definiteness. Assuming that GW has transfers (of the expected form) in this context, we find that

$$n(V, \rho, \text{GW}) = n^{\text{GS}}(V, 0, \rho).$$

Now let σ be a non-degenerate section of V (in the sense of Definition 2.25) and write $i : Z = Z(\sigma) \hookrightarrow X$ for the inclusion of the zero-scheme. Thus i is a regular immersion. In this case one has (see Example 2.27)

$$[L_i] \simeq -[N_{Z/X}^*] \simeq -[V^*|_Z]$$

and consequently, if E has pushforwards along regular immersions, there is a transfer map

$$i_* : E^0(Z) \simeq E^{[V^*]|_Z - [V^*|_Z]}(Z) \simeq E^{i^*V^* + L_i}(Z) \rightarrow E^{V^*}(X).$$

The following result is true in all cases that we know of; but of course it cannot be proved from the weak axioms that we have listed.

Meta-Theorem 3.9. *Let σ be a non-degenerate section of a vector bundle V over a scheme X . Let E be a cohomology theory with Euler classes and pushforwards along regular immersions, such that $E^0(S)$ has a distinguished element 1 (e.g. is a ring). Then*

$$e(V, E) = i_*(1),$$

where we use the identification above for the pushforward.

Going back to the situation where X is smooth and proper over S , V is relatively oriented and E is SL^c -oriented and has transfers along proper lci morphisms, we also have the pushforward

$$\varpi_* : E^0(Z) \simeq E^0(Z, \mathcal{L}^{\otimes 2}|_Z) \xrightarrow{\cong} E^0(Z, \det V^* \otimes \omega_{X/S}|_Z) \simeq E^{L\varpi}(Z) \rightarrow E^0(S).$$

More generally, if $Z' \subset Z$ is a clopen component, then we have a similar transfer originating from $E^0(Z')$.

Definition 3.10. For V, σ, X, E as above, for any clopen component $Z' \subset Z$ we denote by

$$\mathrm{ind}_{Z'}(\sigma, \rho, E) = \varpi'_*(1) \in E^0(S)$$

the local index of σ around Z' in E . Here $\varpi' : Z' \rightarrow S$ is the restriction of ϖ to Z' .

Meta-Corollary 3.11. *Let σ be a non-degenerate section of a relatively oriented vector bundle V over $\pi : X \rightarrow S$. Let E be an SL^c -oriented cohomology theory with Euler classes and pushforwards along proper lci morphisms, such that Meta-Theorem 3.9 applies. Then*

$$n(V, \rho, E) = \sum_{Z'} \mathrm{ind}_{Z'}(\sigma, \rho, E).$$

Proof. By assumption, transfers are compatible with composition, and additive along disjoint unions. The result follows. \square

Example 3.12. If $S = \mathrm{Spec}(k)$ is the spectrum of a field, then Z is zero-dimensional, and hence decomposes into a finite disjoint union of “fat points”. In particular, the Euler number is expressed as a sum of local indices, in bijection with the zeros of our non-degenerate section.

Recall the notion of coordinates from Definition 2.28. The following result states that indices may be computed in local coordinates.

Proposition 3.13. *Let E be an SL^c -oriented cohomology theory with Euler classes and pushforwards along proper lci morphisms. Let $(\psi, \varphi, \sigma_2)$ be a system of coordinates for $(V, X, \sigma_1, \rho_1, Z)$. Then*

$$\mathrm{ind}_Z(\sigma_1, \rho_1, E) = \mathrm{ind}_Z(\sigma_2, \rho_2, E),$$

where ρ_2 is the canonical relative orientation of $\mathcal{O}_{\mathbb{A}^n}/\mathbb{A}^n$.

Proof. Let $\varpi : Z \rightarrow S$ denote the canonical map. Then both sides are obtained as $\varpi_*(1)$, but conceivably the orientations used to define the transfer could be different; we shall show that they are not. In other words, we are given two isomorphisms

$$\widetilde{\det} L_\varpi \xrightarrow{\cong} \mathcal{L}_i^{\otimes 2}$$

and we need to exhibit $\mathcal{L}_1 \xrightarrow{\cong} \mathcal{L}_2$ such that $\alpha_1^{-1}\alpha_2 = \beta^{\otimes 2}$. The isomorphisms α_i arise as

$$\det L_\varpi \simeq \det N_{Z/X} \otimes \omega_{X/S}|_Z \xrightarrow{\cong} \det V \otimes \omega_{X/S}|_Z \xrightarrow{\cong} \mathcal{L}_i^{\otimes 2}|_Z;$$

here $\mathcal{L}_2 = \mathcal{O}$, and for $i = 2$ we implicitly use φ and ψ as well. We first check that the two isomorphisms $\det N_{Z/X} \simeq \det V|_Z$ are the same. Indeed $V|_U \simeq \mathcal{O}^n$ via ψ , and up to this trivialization the isomorphism is given by the trivialization of $C_{Z/X}$ by σ_i ; these are the same by assumption (2). Now we deal with the

second half of the isomorphism. By construction we have an isomorphism $\mathcal{O} \simeq \mathcal{O}^{\otimes 2} \xrightarrow{\alpha_1^{-1}\alpha_2} \mathcal{L}_1^{\otimes 2}$; what we need to check is the corresponding global section of $\mathcal{L}_1^{\otimes 2}$ is a tensor square. Unwinding the definitions, this follows from assumption (3). \square

4. COHOMOLOGY THEORIES REPRESENTED BY MOTIVIC SPECTRA

We recall some background material about motivic extraordinary cohomology theories, i.e. theories represented by motivic spectra. We make essentially no claim to originality.

4.1. Aspects of the six functors formalism. We recall some aspects of the six functors formalism for the motivic stable categories $\mathcal{SH}(-)$, following the exposition in [EHK⁺20a].

4.1.1. *Adjunctions.* For every scheme X , we have a symmetric monoidal, stable category $\mathcal{SH}(X)$. For every morphism $f : X \rightarrow Y$ of schemes we have an adjunction

$$f^* : \mathcal{SH}(Y) \rightleftarrows \mathcal{SH}(X) : f_*.$$

If no confusion can arise, we sometimes write $E_Y := f^*E$. If f is smooth, there is a further adjunction

$$f_{\#} : \mathcal{SH}(X) \rightleftarrows \mathcal{SH}(Y) : f^*.$$

If f is locally of finite type, then there is the exceptional adjunction

$$f_! : \mathcal{SH}(X) \rightleftarrows \mathcal{SH}(Y) : f^!$$

There is a natural transformation $\alpha : f_! \rightarrow f_*$. If f is proper, then α is an equivalence.

The assignments $f \mapsto f^*, f_*, f^!, f_!, f_{\#}$ are functorial. In particular, given composable morphisms f, g of the appropriate type, we have equivalences $(fg)_* \simeq f_*g_*$, and so on.

4.1.2. *Exchange transformations.* Suppose given a commutative square of categories

$$\begin{array}{ccc} \mathcal{C} & \xleftarrow{F} & \mathcal{D} \\ G \uparrow & & G' \uparrow \\ \mathcal{C}' & \xleftarrow{F'} & \mathcal{D}' \end{array},$$

i.e. a natural isomorphism $\gamma : FG' \simeq GF'$. If the functors G', G have right adjoints H', H , then we have the natural transformation

$$F'H' \xrightarrow{\text{unit}} HGF'H' \xrightarrow{\gamma} HFG'H' \xrightarrow{\text{counit}} HF$$

called the associated *exchange transformation*. Similarly if G', G have left adjoints K', K , then we have the exchange transformation

$$KF \xrightarrow{\text{unit}} KFG'K' \xrightarrow{\gamma} KGFK' \xrightarrow{\text{counit}} FK'.$$

Suppose we have a commutative square of schemes

$$(4) \quad \begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ g' \downarrow & & g \downarrow \\ X & \xrightarrow{f} & Y. \end{array}$$

Then we have an induced commutative square of categories

$$\begin{array}{ccc} \mathcal{SH}(X') & \xleftarrow{f'^*} & \mathcal{SH}(Y') \\ g'^* \uparrow & & g^* \uparrow \\ \mathcal{SH}(X) & \xleftarrow{f^*} & \mathcal{SH}(Y). \end{array}$$

Passing to the right adjoints, we obtain the exchange transformation

$$\text{Ex}_*^* : f^*g_* \rightarrow g'_*f'^*.$$

Similarly there is $\text{Ex}_{\#}^* : g'_{\#}f'^* \rightarrow f^*g_{\#}$ (for g smooth; this is in fact an equivalence if (4) is cartesian), and so on.

4.1.3. *Exceptional exchange transformation.* Given a cartesian square of schemes as in (4), with g (and hence g') locally of finite type, there is a canonical equivalence

$$\text{Ex}_!^* : f^*g_! \simeq g'_!f'^*.$$

Passing to right adjoints, we obtain

$$\text{Ex}^{*!} : f'^*g^! \rightarrow g^!f^*.$$

4.1.4. *Thom transformation.* Given a perfect complex \mathcal{E} of vector bundles on X , the motivic J-homomorphism $K(X) \rightarrow \text{Pic}(\mathcal{SH}(X))$ [BH17, §16.2] provides us with an invertible spectrum $\Sigma^{\mathcal{E}} \mathbb{1} \in \mathcal{SH}(X)$. We denote by $\Sigma^{\mathcal{E}} : \mathcal{SH}(X) \rightarrow \mathcal{SH}(X)$, $E \mapsto E \wedge \Sigma^{\mathcal{E}} \mathbb{1}$ the associated invertible endofunctor. If \mathcal{E} is a vector bundle (concentrated in degree zero), then $\Sigma^{\mathcal{E}} \mathbb{1}$ is the suspension spectrum on the Thom space $\mathcal{E}^*/\mathcal{E}^* \setminus 0$.⁶

Lemma 4.1. *The functor $f^!$ commutes with Thom transforms.*

Proof. This follows from the projection formula [CD19, A.5.1(6)] and invertibility of $\Sigma^{(-)} \mathbb{1}$:

$$f^!(\Sigma^V X) \simeq f^! \underline{\text{Hom}}(\Sigma^{-V} \mathbb{1}, X) \simeq \underline{\text{Hom}}(f^* \Sigma^{-V} \mathbb{1}, f^! X) \simeq \Sigma^{f^* V} f^! X.$$

□

4.1.5. *Purity transformation.* Let $f : X \rightarrow Y$ be a smoothable [EHK⁺20a, §2.1.21] lci morphism. Then the cotangent complex L_f is perfect, and there exists a canonical *purity transformation*

$$\mathfrak{p}_f : \Sigma^{L_f} f^* \rightarrow f^!.$$

4.2. Cohomology groups and Gysin maps.

4.2.1. Let S be a scheme and $E \in \mathcal{SH}(S)$. Given $(\pi : X \rightarrow S) \in \text{Sch}_S$, $i : Z \hookrightarrow X$ closed, $\xi \in K(Z)$, we define the ξ -twisted E -cohomology of X with support in Z as

$$E_Z^\xi(X) = [\mathbb{1}, \pi_* i_! \Sigma^\xi i^! \pi^* E]_{\mathcal{SH}(S)}.$$

This assignment forms a cohomology theory in the sense of §3.1. It takes values in abelian groups, has supports, and satisfies the disjoint union property. We shall see that it has transfers for proper lci maps. It need not be orientable in general.

If $Z = X$, we may omit it from the notation and just write $E^\xi(X)$. As before if ξ is a trivial virtual vector bundle of rank $n \in \mathbb{Z}$ then we also write $E_Z^n(X)$ instead of $E_Z^\xi(X)$.

Example 4.2. Suppose that $\xi = i^* V$, where V is a vector bundle on X . We have

$$E_Z^\xi(X) = [\mathbb{1}, \pi_* i_! \Sigma^\xi i^! \pi^* E]_{\mathcal{SH}(S)} \simeq [\mathbb{1}, i^! \Sigma^\xi \pi^* E]_{\mathcal{SH}(Z)} \simeq [i_* \mathbb{1}, \Sigma^\xi \pi^* E]_{\mathcal{SH}(X)},$$

where we have used that $i_* \simeq i_!$ and Lemma 4.1. Using the localization sequence $j_{\#} j^* \rightarrow \text{id} \rightarrow i_* i^*$ [Hoy17, Theorem 6.18(4)] to identify $i_* \mathbb{1} \simeq X/X \setminus Z$ we find that

$$E_Z^\xi(X) \simeq [X/(X \setminus Z), \Sigma^\xi \pi^* E]_{\mathcal{SH}(X)}.$$

Remark 4.3. The final expression in the above example only depends on $Z \subset X$ as a subset, not subscheme. This also follows directly from the definition, since $\mathcal{SH}(Z) \simeq \mathcal{SH}(Z_{\text{red}})$; this is another consequence of localization.

4.2.2. *Functoriality in E .* If $\alpha : E \rightarrow F \in \mathcal{SH}(S)$ is any morphism, then there is an induced morphism $\alpha_* : E_Z^\xi(X) \rightarrow F_Z^\xi(X)$. This just follows from the fact that π^* etc. are functors.

4.2.3. *Contravariant functoriality in X .* Let $f : X' \rightarrow X \in \text{Sch}_S$. Then there is a pullback map

$$f^* : E_Z^\xi(X) \rightarrow E_{f^{-1}(Z)}^{f^* \xi}(X'),$$

coming from the morphism

$$\begin{aligned} \pi_* i_! \Sigma^\xi i^! \pi^* E &\xrightarrow{\text{unit}} \pi_* f_* f^* i_! \Sigma^\xi i^! \pi^* E \simeq \pi'_* f'_* i'_! \Sigma^\xi i'^! \pi'^* E \xrightarrow{\text{Ex}_1^*} \pi'_* i'_! f'^* \Sigma^\xi i'^! \pi'^* E \\ &\simeq \pi'_* i'_! \Sigma^{f^* \xi} f'^* i'^! \pi'^* E \xrightarrow{\text{Ex}^{*!}} \pi'_* i'_! \Sigma^{f^* \xi} i'^* f^* \pi^* E \simeq \pi'_* i'_! \Sigma^{f^* \xi} i'^* \pi'^* E. \end{aligned}$$

Here the Ex_1^* and $\text{Ex}^{*!}$ come from the cartesian square

$$\begin{array}{ccc} Z' & \xrightarrow{f'} & Z \\ i' \downarrow & & i \downarrow \\ X' & \xrightarrow{f} & X \end{array}$$

⁶Indeed, in [EHK⁺20a, §2.1.2], the transformation $\Sigma^{\mathcal{E}}$ is built out of $\text{Spec}(\text{Sym}(\mathcal{E}))$ —which is the vector bundle corresponding to \mathcal{E}^* , in our convention.

Lemma 4.4. *Let $f : X' \rightarrow X \in \text{Sch}_X$, $Z \subset X$, $\xi \in K(Z)$ and $\alpha : E \rightarrow F \in \mathcal{SH}(S)$. The following square commutes*

$$\begin{array}{ccc} E_{Z'}^{\xi'}(X') & \xrightarrow{\alpha_*} & F_{Z'}^{\xi'}(X') \\ f^* \uparrow & & f^* \uparrow \\ E_Z^\xi(X) & \xrightarrow{\alpha_*} & F_Z^\xi(X'), \end{array}$$

where $\xi' = f^*(\xi)$ and $Z' = f^{-1}(Z)$.

Proof. This is just an expression of the fact that the exchange transformations used to build f^* are indeed natural transformations. \square

4.2.4. *Covariant functoriality in X .* Suppose given a commutative square in Sch_S

$$(5) \quad \begin{array}{ccc} Z_1 & \xrightarrow{i} & X \\ g \downarrow & & f \downarrow \\ Z_2 & \xrightarrow{k} & Y, \end{array}$$

where f is smoothable lci, i, k are closed immersions and g is proper. For every $\xi \in K(Z_2)$, there is the *Gysin map*

$$f_* : E_{Z_1}^{g^* \xi + i^* L_f}(X) \rightarrow E_{Z_2}^\xi(Y)$$

coming from the morphism

$$f_* i_! \Sigma^{g^* \xi + i^* L_f} i^! f^* E_Y \xrightarrow{p_f} f_* i_! \Sigma^{g^* \xi} i^! f^! E_Y \simeq k_! g_! \Sigma^{g^* \xi} g^! k^! E_Y \simeq k_! g_! g^! \Sigma^\xi k^! E_Y \xrightarrow{\text{counit}} k_! \Sigma^\xi k^! E_Y,$$

where we used Lemma 4.1 to move Σ^{L_f} through $i^!$ and Σ^ξ through $g^!$, and also used $i_* \simeq i_!$, $k_* \simeq k_!$, $g_* \simeq g_!$.

Remark 4.5. In [EHK⁺20a], the Gysin map is denoted by $f_!$, to emphasize that it involves the purity transform. We find the notation f_* more convenient.

Lemma 4.6. *The following hold.*

(1) *Suppose given a square as in (5) and $\alpha : E \rightarrow F \in \mathcal{SH}(S)$. Then the following square commutes*

$$\begin{array}{ccc} E_{Z_1}^{g^* \xi + i^* L_f}(X) & \xrightarrow{\alpha_*} & F_{Z_1}^{g^* \xi + i^* L_f}(X) \\ f_* \downarrow & & f_* \downarrow \\ E_{Z_2}^\xi(Y) & \xrightarrow{\alpha_*} & F_{Z_2}^\xi(Y). \end{array}$$

(2) *Suppose given a square as in (5) and $s : Y' \rightarrow Y$ such that s, f are tor-independent. Let $X' = X \times_Y Y'$, $Z'_i = Z_i \times_Y Y'$, i', g', k', f' the induced maps, and so on. Then the following square commutes*

$$\begin{array}{ccc} E_{Z_1}^{g^* \xi + i^* L_f}(X) & \xrightarrow{s^*} & E_{Z'_1}^{g'^* \xi' + i'^* L_{f'}}(X) \\ f_* \downarrow & & f'_* \downarrow \\ E_{Z_2}^\xi(Y) & \xrightarrow{s^*} & E_{Z'_2}^{\xi'}(Y'). \end{array}$$

(3) *Suppose given a commutative diagram in Sch_S as follows*

$$\begin{array}{ccc} Z_1 & \xrightarrow{i} & X \\ g \downarrow & & f \downarrow \\ Z_2 & \xrightarrow{k} & Y \\ g' \downarrow & & f' \downarrow \\ Z_3 & \xrightarrow{l} & W. \end{array}$$

Then given $\xi \in K(Z_3)$, we have

$$f'_* f_* = (f' f)_* : E_{Z_1}^{(g' g)^* \xi + i^* L_{f' f}}(X) \rightarrow E_{Z_3}^\xi(W).$$

Here we use the equivalence $L_{f' f} \simeq f'^* L_f + L_{f'} \in K(X)$, coming from the cofiber sequence $f'^* L_f \rightarrow L_{f' f} \rightarrow L_{f'}$.

Proof. (1) Since $f_* : E_{Z_1}^{g^* \xi + i^* L_f}(X) \rightarrow E_{Z_2}^\xi(Y)$ is obtained as $[1, \zeta_E]$ where ζ is a natural transformation of endofunctors of $\mathcal{SH}(S)$, this is clear.

(2) Consider the following diagram.

$$\begin{array}{ccccc}
f_* i_! \Sigma^{g^* \xi + i^* L_f} i^! f^* & \xrightarrow{\text{unit}_X} & f_* s_{X^*} s_X^* i_! \Sigma^{g^* \xi + i^* L_f} i^! f^* & \xrightarrow{\text{Ex}} & f_* s_{X^*} \Sigma^{g'^* \xi' + i'^* L_{f'}} i'^! f'^* s^* \\
\downarrow \simeq & & \downarrow \simeq & & \downarrow \simeq \\
f_* i_! \Sigma^{g^* \xi} i^! \Sigma^{L_f} f^* & \xrightarrow{\text{unit}_X} & f_* s_{X^*} s_X^* i_! \Sigma^{g^* \xi} i^! \Sigma^{L_f} f^* & \xrightarrow{\text{Ex}} & f_* s_{X^*} i'_! \Sigma^{g'^* \xi'} i'^! \Sigma^{L_{f'}} f'^* s^* \\
\downarrow p_f & & \downarrow p_f & (a) & \downarrow p_{f'} \\
f_* i_! \Sigma^{g^* \xi} i^! f^! & \xrightarrow{\text{unit}_X} & f_* s_{X^*} s_X^* i_! \Sigma^{g^* \xi} i^! f^! & \xrightarrow{\text{Ex}} & f_* s_{X^*} i'_! \Sigma^{g'^* \xi'} i'^! f'^! s^* \\
\downarrow \simeq & \searrow \text{unit} & (b) & & \downarrow \simeq \\
& & s_* s^* f_* i_! \Sigma^{g^* \xi} i^! f^! & \xrightarrow{\text{Ex}} & s_* f_* i'_! \Sigma^{g'^* \xi'} i'^! f'^! s^* \\
& & \downarrow \simeq & & \downarrow \simeq \\
k_! g_! \Sigma^{g^* \xi} g^! k^! & \xrightarrow{\text{unit}} & s_* s^* k_! g_! \Sigma^{g^* \xi} g^! k^! & \xrightarrow{\text{Ex}} & s_* k'_! g'_! \Sigma^{g'^* \xi'} g'^! k'^! s^* \\
\downarrow \simeq & & \downarrow \simeq & & \downarrow \simeq \\
k_! g_! g^! \Sigma^\xi k^! & \xrightarrow{\text{unit}} & s_* s^* k_! g_! g^! \Sigma^\xi k^! & \xrightarrow{\text{Ex}} & s_* k'_! g'_! g'^! \Sigma^{\xi'} k'^! s^* \\
\downarrow \text{counit} & & \downarrow \text{counit} & (b) & \downarrow \text{counit}' \\
k_! \Sigma^\xi k^! & \xrightarrow{\text{unit}} & s_* s^* k_! \Sigma^\xi k^! & \xrightarrow{\text{Ex}} & s_* k'_! \Sigma^{\xi'} k'^! s^*
\end{array}$$

Here $s_X : X' \rightarrow X$ is the canonical map and $\xi' = (Z'_2 \rightarrow Z_2)^* \xi$. All the unlabelled equivalences arise from moving Thom transforms through (various) pullbacks, compatibility of pullbacks and pushforwards with composition, and equivalences of the form $p_* \simeq p_!$ for p proper. All the maps labelled Ex are exchange transformations expressing compatibility of $p_*, p_!, p^!$ with base change. Denote the diagram by \mathcal{D} . Then $[1, \mathcal{DE}_Y]$ yields a diagram of abelian groups. The outer square of that diagram identifies with the square we are trying to show commutes. It hence suffices to show that \mathcal{D} commutes. All cells commute for trivial reasons, except for (a) which commutes by [EHK⁺20a, Proposition 2.2.2(ii)] and (b) which commute by stability of the counit transformations under base change.

(3) Consider the following diagram.

$$\begin{array}{ccccc}
f'_* f_* i_! \Sigma^{g'^* g'^* \xi + i^* L_f + i'^* f'^* L_{f'}} i'^! f^* f'^* & & & & \\
\downarrow \simeq & & & & \\
l_! g'_! g_! \Sigma^{g'^* g'^* \xi} i^! \Sigma^{L_f} f^* \Sigma^{L_{f'}} f'^* & \xrightarrow{\simeq} & l_!(g'g)_! \Sigma^{(g'g)^* \xi} i^! \Sigma^{L_{f'f}} (f'f)^* & & \\
\downarrow p_f & & \downarrow p_{f'} & & \downarrow p_{f'f} \\
l_! g'_! g_! \Sigma^{g'^* g'^* \xi} i^! f^! \Sigma^{L_{f'}} f'^* & (a) & & & \\
\downarrow p_{f'} & & \downarrow \simeq & & \downarrow \simeq \\
l_! g'_! g_! \Sigma^{g'^* g'^* \xi} i^! f^! f'^! & \xrightarrow{\simeq} & l_!(g'g)_! \Sigma^{(g'g)^* \xi} i^! (f'f)^! & & \\
\downarrow \text{counit}_g & \swarrow p_{f'} & \downarrow \simeq & & \downarrow \simeq \\
l_! g'_! g_! \Sigma^{g'^* \xi} k^! \Sigma^{L_{f'}} f'^* & & l_! g'_! g_! \Sigma^{g'^* \xi} k^! f'^! & & \\
\downarrow p_{f'} & \swarrow \text{counit}_g & \downarrow \simeq & & \downarrow \simeq \\
l_! g'_! \Sigma^{g'^* \xi} k^! f'^! & & l_! g'_! g_! g'^! \Sigma^\xi l^! & \xrightarrow{\simeq} & l_!(g'g)_! (g'g)^! \Sigma^\xi l^! \\
\downarrow \simeq & \swarrow \text{counit}_g & (b) & & \downarrow \simeq \\
l_! g'_! g'^! \Sigma^\xi l^! & \xrightarrow{\text{counit}_{g'}} & l_! \Sigma^\xi l^! & \xleftarrow{\text{counit}_{g'g}} &
\end{array}$$

All the unlabelled equivalences arise from moving Thom transforms through (various) pullbacks, compatibility of pullbacks and pushforwards with composition, and equivalences of the form $p_* \simeq p_!$ for p proper. Denote the diagram by \mathcal{D} . Then $[\mathbb{1}, \mathcal{D}E_W]$ yields a diagram of abelian groups. Going from the top to the bottom middle via the leftmost past, we obtain $f'_* f_*$; going instead via the rightmost path we obtain $(f'f)_*$. It hence suffices to show that \mathcal{D} commutes. All cells commute for trivial reasons, except for (a) which commutes by [EHK⁺20a, Proposition 2.2.2(i)], and (b) which commutes by compatibility of the counit transformations with composition. \square

Example 4.7. Consider a commutative square as in (5), with $X = Y$ and $f = \text{id}$, so that $g : Z_1 \hookrightarrow Z_2$ is a closed immersion. Then $f_* : E_{Z_1}^{g^* \xi}(X) \rightarrow E_{Z_2}^\xi(X)$ is the “extension of support” map. In particular taking $Z_2 = X$ as well, we obtain the map $E_{Z_1}^{i^* \xi}(X) \rightarrow E^\xi(X)$ “forgetting the support”. Lemma 4.6(3) now in particular tells us that given a proper map $f : X \rightarrow Y$, a closed immersion $i : Z \hookrightarrow X$ and $\xi \in K(Y)$, the following diagram commutes

$$\begin{array}{ccc} E_Z^{i^* f^* \xi}(X) & \longrightarrow & E^{f^* \xi}(X) \\ f_* \downarrow & & f_* \downarrow \\ E_X^\xi(X) & \xlongequal{\quad} & E^\xi(X), \end{array}$$

where the upper horizontal map forgets the support.

4.3. Orientations.

4.3.1. *Product structures.* By a ring spectrum (over S) we mean an object $E \in \mathcal{SH}(S)$ together with homotopy classes of maps $u : \mathbb{1} \rightarrow E$ and $m : E \wedge E \rightarrow E$ satisfying the evident identities. If E is a ring spectrum, then there are multiplication maps

$$E_{Z_1}^\xi(X) \times E_{Z_2}^{\xi'}(X) \rightarrow E_{Z_1 \cap Z_2}^{\xi + \xi'}(X)$$

induced by

$$(\Sigma^\xi E) \wedge (\Sigma^{\xi'} E) \simeq \Sigma^{\xi + \xi'} E \wedge E \xrightarrow{m} \Sigma^{\xi + \xi'} E$$

and the diagonal⁷

$$X/X \setminus (Z_1 \cap Z_2) \rightarrow X/X \setminus Z_1 \wedge X/X \setminus Z_2.$$

Lemma 4.8. *The multiplicative structure on E -cohomology is compatible with pullback: given $Z_1, Z_2 \subset X$, $\xi, \xi' \in K(X)$, $f : X' \rightarrow X$, the following diagram commutes*

$$\begin{array}{ccc} E_{Z_1}^\xi(X) \times E_{Z_2}^{\xi'}(X) & \longrightarrow & E_{Z_1 \cap Z_2}^{\xi + \xi'}(X) \\ f^* \times f^* \downarrow & & f^* \downarrow \\ E_{f^{-1}(Z_1)}^{f^* \xi}(X') \times E_{f^{-1}(Z_2)}^{f^* \xi'}(X') & \longrightarrow & E_{f^{-1}(Z_1) \cap f^{-1}(Z_2)}^{f^* \xi + f^* \xi'}(X') \end{array}$$

Proof. Immediate from the definitions. \square

4.3.2. *Thom spectra.* Let $G = (G_n)_n$ be a family of finitely presented S -group schemes, equipped with a morphism of associative algebras $G \rightarrow (\text{GL}_{nk, S})_n$ (for the Day convolution symmetric monoidal structure on $\text{Fun}(\mathbb{N}, \text{Grp}(\text{Sch}_S))$). Then there is a notion of (stable) vector bundle with structure group G , the associated K -theory space $K^G(X)$, and the associated *Thom spectrum* MG , which is a ring spectrum [BH17, Example 16.22].

Example 4.9. If $G_n = \text{GL}_n$, then $K^G(X) = K(X)$ and MGL is the algebraic cobordism spectrum [BH17, Theorem 16.13]. If $G_n = \text{SL}_n$ (respectively Sp_n), then $K^G(X)$ is the K -theory of oriented (respectively symplectic) vector bundles in the usual sense, and MSL (respectively $M\text{Sp}$) is the Thom spectrum as defined in [PW10a].

In order to work effectively with MG , one needs to know that it is stable under base change. This is easily seen to be true for $MGL, MSL, M\text{Sp}$ [BH17, Example 16.23]. We record the following more general result for future reference.

Proposition 4.10. *The Thom spectrum MG is stable under base change, provided that each G_n is flat and quasi-affine.*

⁷One easily checks that the diagonal $X \rightarrow X \times X$ induces a map as indicated.

Proof. We have a presheaf $K^G \in \mathcal{P}(\text{Sch}_S)$ and a map $K^G \rightarrow K$. For $f : X \rightarrow S \in \text{Sch}_S$, denote by $K_X^G \in \mathcal{P}(\text{Sm}_X)$ and $j_X : K_X^G \rightarrow K|_{\text{Sm}_X}$ the restrictions. Then by definition $\text{MG}_X = M_X(j_X)$, where $M_X : \mathcal{P}(\text{Sm}_X)_{/K} \rightarrow \mathcal{SH}(X)$ is the motivic Thom spectrum functor [BH17, §16.1]. Let $LK_S^G \in \mathcal{P}(\text{Sch}_S)$ denote the left Kan extension of K_S^G . We claim that $LK_S^G \rightarrow K^G$ is a Nisnevich equivalence. Assuming this, we deduce that $f^*K_S^G \simeq (LK_S^G)|_{\text{Sm}_X} \rightarrow K_X^G$ is a Nisnevich equivalence. Since M_X inverts Nisnevich equivalences [BH17, Proposition 16.9], this implies that $f^*\text{MG}_S \simeq \text{MG}_X$, which is the desired result.

To prove the claim, we first note that by [EHK⁺20b, Lemma 3.3.9], we may assume S affine, and it suffices to prove that the restriction of K^G to Aff_S is left Kan extended from smooth affine S -schemes. By definition $K^G = (\text{Vect}^G)^{gp}$, where $\text{Vect}^G = \coprod_{n \geq 0} BG_n$ (here the coproduct is as stacks, i.e. fppf sheaves). The desired result now follows from [EHK⁺20b, Proposition A.0.4 and Lemma A.0.5] (noting that the coproduct of stacks is the same as coproduct of Σ -presheaves, and Kan extension preserves Σ -presheaves). \square

Now let $\xi \in K^G(X)$. Then there is a canonical equivalence [BH17, Example 16.29]

$$\Sigma^\xi \text{MG}_X \simeq \Sigma^{|\xi|} \text{MG}_X.$$

We denote by $t_\xi \in \text{MG}^{\xi-|\xi|}(X)$ the class of the map

$$1 \xrightarrow{u} \text{MG}_X \simeq \Sigma^{\xi-|\xi|} \text{MG}_X.$$

4.3.3. Oriented ring spectra.

Definition 4.11. Let $E \in \mathcal{SH}(S)$ be a ring spectrum and $G = (G_n)_n$ a family of group schemes as in §4.3.2. By a *strong G -orientation* of E we mean a ring map $\text{MG} \rightarrow E$.

Example 4.12. The spectrum KO is strongly SL -oriented; see Corollary A.3.

Note that if $E \in \mathcal{SH}(S)$ is strongly G -oriented, then there is no reason a priori why E_X should be strongly G_X -oriented. This is true if MG is stable under base change, so for most reasonable G by Proposition 4.10. We will not talk about strong G -orientations unless MG is stable under base change, so assume this throughout.

Given $\xi \in K^G(X)$, the map $\text{MG} \rightarrow E$ provides us with $t_\xi = t_\xi(E) \in E^{\xi-|\xi|}(X)$.

Proposition 4.13. *Let E be strongly G -oriented and $\xi \in K^G(X)$.*

- (1) *The classes $t_\xi(E)$ are stable under base change: for $f : X' \rightarrow X$ we have $t_{f^*\xi}(E) = f^*t_\xi(E)$.*
- (2) *Multiplication by $t_\xi(E)$ induces an equivalence $\Sigma^{|\xi|}E \simeq \Sigma^\xi E$ and an isomorphism $t : E_Z^{|\xi|}(X) \simeq E_Z^\xi(X)$, called the Thom isomorphism.*
- (3) *The Thom isomorphism is compatible with base change: $f^*(t(x)) = t(f^*(x))$.*

In particular, E is G -oriented.

Proof. (1) follows from the same statement for MG , where it holds by construction. For the first half of (2), it suffices to show that $t_\xi(E)$ is a unit in the Picard-graded homotopy ring of E . This follows from the same statement for MG . The second half of (2) follows. (3) immediately follows from (1). \square

Example 4.14. Let E be strongly GL -oriented. Then for any $\xi \in K(X)$ we obtain $E^\xi(X) \simeq E^{rk(\xi)}(X)$, so E is oriented in the sense of §3.2

Definition 4.15. For $X \in \text{Sch}_S$ and \mathcal{L} a line bundle on X put

$$E_Z^n(X, \mathcal{L}) = E_Z^{n-1+\mathcal{L}}(X).$$

Example 4.16. Let E be strongly SL -oriented and $\xi \in K(X)$. Then $\xi' := \xi - (|\xi| - 1 + \det \xi) \in K(X)$ lifts canonically to $K^{\text{SL}}(X)$, whence by Proposition 4.13 we get a canonical (Thom) isomorphism $E^\xi(X) \simeq E^{|\xi|}(X, \det \xi)$. In particular, E is SL -oriented in the sense of §3.2.

Remark 4.17. If E is strongly SL -oriented, then since $\det(\mathcal{L}_1 \oplus \mathcal{L}_2) \simeq \mathcal{L}_1 \otimes \mathcal{L}_2$, by Example 4.16 the product structure on E -cohomology twisted by line bundles takes the form $E^n(X, \mathcal{L}_1) \times E^m(X, \mathcal{L}_2) \rightarrow E^{n+m}(X, \mathcal{L}_1 \otimes \mathcal{L}_2)$.

Remark 4.18. Strong G -orientations have better permanence properties than ordinary ones (provided that MG is stable under base change): they are stable under base change and taking (very) effective covers, for example.

4.4. **SL^c-orientations.** A. Ananyevskiy has done important work on SL and SL^c orientations. We shall make use of the following result; see e.g. [Ana20, Theorem 1.1].

Proposition 4.19 (Ananyevskiy). *Let $E \in \mathcal{SH}(S)$ be SL-oriented and $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3$ be line bundles on X . Suppose given an isomorphism $\mathcal{L}_1 \simeq \mathcal{L}_2 \otimes \mathcal{L}_3^{\otimes 2}$.*

(1) *There is a canonical equivalence $\Sigma^{\mathcal{L}_1} E \simeq \Sigma^{\mathcal{L}_2} E$, compatible with base change.*

(2) *There is a canonical isomorphism $E_{\mathbb{Z}}^n(X, \mathcal{L}_1) \simeq E_{\mathbb{Z}}^n(X, \mathcal{L}_2)$, compatible with base change.*

In particular, the cohomology theory represented by E is SL^c-oriented.

Proof. Note that (2) follows from (1). Let $\mathcal{L} = \mathcal{L}_3$. It suffices to exhibit a canonical equivalence $\Sigma^{\mathcal{L}^{\otimes 2}} E \simeq \Sigma^{\mathcal{O}} E$. We have canonical equivalences

$$\Sigma^{\mathcal{L}+\mathcal{L}} E \simeq \Sigma^{\mathcal{O}+\mathcal{L}^{\otimes 2}} E, \quad \Sigma^{\mathcal{L}+\mathcal{L}^*} E \simeq \Sigma^{\mathcal{O}^2} E, \quad \Sigma^{\mathcal{L}+\mathcal{L}} \simeq \Sigma^{\mathcal{L}+\mathcal{L}^*},$$

by [Ana20, Corollary 3.9, Lemma 4.1]. Consequently $\Sigma^{\mathcal{O}+\mathcal{L}^{\otimes 2}} E \simeq \Sigma^{\mathcal{O}^2} E$, whence the claim. \square

5. EULER CLASSES FOR REPRESENTABLE THEORIES

5.1. **Tautological Euler class.** Let $E \in \mathcal{SH}(S)$ be a ring spectrum, $X \in \text{Sch}_S$ and V a vector bundle on X .

Definition 5.1. We denote by $e(V) = e(V, E) \in E^{V^*}(X)$ the *tautological Euler class* of V , defined as the composite

$$\mathbb{1}_X \simeq \Sigma_+^\infty V \rightarrow \Sigma^\infty V / (V \setminus 0) \simeq \Sigma^{V^*} \mathbb{1} \xrightarrow{u} \Sigma^{V^*} E|_X \in \mathcal{SH}(X).$$

Lemma 5.2. *Let $f : X' \rightarrow X \in \text{Sch}_S$. Then*

$$f^* e(V, E) = e(f^* V, E) \in E^{f^* V}(X').$$

Proof. Immediate. \square

If E is strongly SL-oriented in the sense of §4.3.3, and hence SL^c-oriented in the sense of §3.2, then for any relatively oriented vector bundle V over a smooth and proper scheme X/S we obtain an Euler number $n(V, \rho, E) \in E^0(S)$. See §3.4.

5.2. Integrally defined Euler numbers.

Corollary 5.3 (Euler numbers are stable under base change). *Let E be a strongly SL-oriented cohomology theory and let V be vector bundle V over a smooth and proper scheme X/S , relatively oriented by ρ . Let $f : S' \rightarrow S$ be a morphism of schemes. Then*

$$f^* n(V, \rho, E) = n(f^* V, f^* \rho, f^* E) \in E^0(S').$$

Proof. This holds since all our constructions are stable under base change. See in particular Lemma 4.6(2) (for compatibility of Gysin maps with pullback, which applies since $X \rightarrow S$ is smooth), Proposition 4.19 and Proposition 4.13(3) (ensuring that the identification $E^{V^*}(X) \simeq E^{L_\pi}(X)$ is compatible with base change) and Lemma 5.2 (for compatibility of Euler classes with base change). \square

Proposition 5.4. *Let d be even or $d = 1$, $X/\mathbb{Z}[1/d]$ smooth and proper and V/X a relatively oriented vector bundle. Then for any field k with $2d \in k^\times$ we have*

$$n(V_k, \rho, \mathbb{H}\tilde{\mathbb{Z}}) \in \mathbb{Z}[\langle -1 \rangle, \langle 2 \rangle, \dots, \langle d \rangle] \subset \text{GW}(k).$$

In fact there is a formula

$$n(V_k, \rho, \mathbb{H}\tilde{\mathbb{Z}}) = \sum_{a \in \mathbb{Z}[1/d]^\times} n_a \langle a \rangle$$

which holds over any such field, with the coefficients $n_a \in \mathbb{Z}$ independent of k (and zero for all but finitely many a).

Remark 5.5. If $d = 1$, Proposition 5.4 relies on the novel results about Hermitian K -theory of the integers from [CDH⁺20]. In the below proof, this is manifested in the dependence of [BH20, Lemma 3.38(2)] on these results. We will later use Proposition 5.4 for the $d = 1$ case of Theorem 5.11, whence this result is also using [CDH⁺20] in an essential way. For $d \geq 2$, the proof is independent of [CDH⁺20].

Note that here the assumption that the rank of V equals the dimension of X is included in the hypothesis that V/X a relatively oriented vector bundle (see Definition 1.5).

Proof. Recall the very effective cover functor \tilde{f}_0 and the truncation in the effective homotopy t -structure π_0^{eff} , for example from [Bac17, §§3.4]. We have a diagram of spectra

$$\mathrm{KO}_k \leftarrow \tilde{f}_0 \mathrm{KO}_k \rightarrow \pi_0^{\text{eff}} \mathrm{KO}_k \leftarrow \pi_0^{\text{eff}} \mathrm{MSL}_k \simeq H\tilde{\mathbb{Z}};$$

see [BH17, Example 16.34] for the last equivalence. The functors \tilde{f}_0 and π_0^{eff} are lax monoidal in an appropriate sense, so this is a diagram of ring spectra. Moreover all of the ring spectra are strongly SL-oriented (via the ring map $\mathrm{MSL}_k \rightarrow \pi_0^{\text{eff}} \mathrm{MSL}_k$); see also Remark 4.18. Finally all the maps induce isomorphisms on $[1, -]$, essentially by construction. It follows that $n(V_k, \rho, \mathrm{KO}) = n(V_k, \rho, H\tilde{\mathbb{Z}}) \in \mathrm{GW}(k)$. We may thus as well prove the result for $n(V_k, \rho, \mathrm{KO})$ instead.

If d is even, then we have $\mathrm{KO} \in \mathcal{SH}(\mathbb{Z}[1/d])$, and by Corollary 5.3 we see that $n(V_k, \rho, \mathrm{KO}) \in \mathrm{im}(\mathrm{GW}(\mathbb{Z}[1/d]) \rightarrow \mathrm{GW}(k))$. The result thus follows from Lemma 5.6 below. If $d = 1$, we employ the SL-oriented ring spectrum $\mathrm{KO}' \in \mathcal{SH}(\mathbb{Z})$ from [BH20, §3.8.3]. We find that $n(V_k, \rho, \mathrm{KO})$ is the image of $n(V, \rho, \mathrm{KO}') \in \mathrm{KO}'^0(\mathbb{Z})$. This latter group is isomorphic to $\mathrm{GW}(\mathbb{Z})$ by [BH20, Lemma 3.38(2)], whence the result. \square

Lemma 5.6. *Let d be even or $d = 1$. As a ring, $\mathrm{GW}(\mathbb{Z}[1/d])$ is generated by $\langle -1 \rangle$ and $\langle p \rangle$, for the primes $p \mid d$.*

Proof. Let $A = \mathbb{Z}[1/d]$ and $U = \mathrm{Spec} A$. If $d = 1$, this is well-known [MH73, Theorem II.4.3]. Hence from now on $1/2 \in A$. Using Lemma B.2, it suffices to prove the analogous statement for $W(A)$. Let $I \subset W$ be the fundamental ideal and denote by I^* its powers; we view these as presheaves on U . Consider the commutative graded ring $k_* = H_{\mathrm{Nis}}^*(U, I^*/I^{*+1})$. Of course $k_0 = \mathbb{Z}/2$. It follows from [BØ21, Corollary 4.9], [Bac20, Theorem 2.1, Lemma 2.7] and [Spi18, Theorem 3.9] that $k_1 \simeq \mathcal{O}^\times(U)/2$ and $k_n \simeq \mathbb{Z}/2\{(-1)^n\}$ for $n \geq 2$. We have the classes $\langle a \rangle - 1 \in I(U)$ (for $a \in \mathcal{O}^\times(U)$) showing that $H_{\mathrm{Nis}}^0(U, I^n) \rightarrow k_n$ is surjective. The short exact sequences

$$0 \rightarrow H_{\mathrm{Nis}}^0(U, I^{n+1}) \rightarrow H_{\mathrm{Nis}}^0(U, I^n) \rightarrow k_n$$

thus show that $H_{\mathrm{Nis}}^0(U, W)$ is generated by the $\langle a \rangle$ together with $H_{\mathrm{Nis}}^0(U, I^n)$, for any n . For n sufficiently large, $H_{\mathrm{Nis}}^0(U, I^n) \simeq 2^n H_{\mathrm{ret}}^0(U, \mathbb{Z}) \simeq 2^n \mathbb{Z}$ [Bac20, Proposition 2.3] is generated by $\langle (-1) - 1 \rangle^n$; thus $H_{\mathrm{Nis}}^0(U, W)$ is generated by the $\langle a \rangle$. It remains to observe that $H_{\mathrm{Nis}}^0(U, W) = W(U)$. This follows from the descent spectral sequence for computing $W(U) = [U, \mathrm{KW}]$, using that the motivic spectrum KW has $\pi_* \mathrm{KW} = a_{\mathrm{Nis}} W$ for $* \equiv 0 \pmod{4}$ and $\pi_* \mathrm{KW} = 0$. \square

Notation 5.7. Let $A \hookrightarrow \mathbb{R}$ and V be a relatively oriented, rank n vector bundle on a smooth proper n -dimensional scheme X over A . We have $\mathrm{GW}(\mathbb{R}) \simeq \mathbb{Z} \oplus \mathbb{Z}\langle -1 \rangle$. There are thus unique integers $n_{\mathbb{R}}, n_{\mathbb{C}} \in \mathbb{Z}$ such that

$$n(V_{\mathbb{R}}, \rho, H\tilde{\mathbb{Z}}) = \frac{n_{\mathbb{C}} + n_{\mathbb{R}}}{2} + \frac{n_{\mathbb{C}} - n_{\mathbb{R}}}{2} \langle -1 \rangle.$$

The integers $n_{\mathbb{R}}$ and $n_{\mathbb{C}}$ are the Euler numbers of the corresponding real and complex topological vector bundles respectively, at least when X is projective, justifying the notation. To show this, consider the cycle class map $\mathrm{CH}^*(X) \rightarrow H^*(X(\mathbb{C}), \mathbb{Z})$ from the Chow ring of a smooth \mathbb{C} -scheme X to the singular cohomology of the complex manifold $X(\mathbb{C})$. See [Ful84, Chapter 19]. Furthermore, there are real cycle class maps from oriented Chow of \mathbb{R} -smooth schemes X to the singular cohomology of the real manifold $X(\mathbb{R})$, discussed in [Jac17] and [HWXZ19] (as well as more refined real cycle class maps defined in [BW20] and [HWXZ19]): For a smooth \mathbb{R} -scheme X and a line bundle $\mathcal{L} \rightarrow X$, consider the real cycle class map

$$\widetilde{\mathrm{CH}}^*(X, \mathcal{L}) \cong H^n(X, \mathrm{K}_n^{\mathrm{MW}}(\omega_{X/k})) \cong H\tilde{\mathbb{Z}}^{L_\pi}(X) \rightarrow H^*(X(\mathbb{C}), \mathbb{Z}(\mathcal{L}))$$

from the oriented Chow groups twisted by \mathcal{L} to the singular homology of the associated local system on $X(\mathbb{R})$. We use results on the real cycle class map due to Hornbostel, Wendt, Xie, and Zibrowius [HWXZ19], including compatibility with pushforwards. Since they only had need of this compatibility in the case of the pushforward by a closed immersion, we first extend this slightly.

Lemma 5.8. *Let $\pi : X \rightarrow \mathrm{Spec} k$ be the structure map of a smooth, projective scheme X of dimension n over the real numbers \mathbb{R} . Then the following square commutes,*

$$\begin{array}{ccc} H\tilde{\mathbb{Z}}^{L_\pi}(X) & \longrightarrow & H^n(X(\mathbb{R}), \mathbb{Z}(\omega_{X/k})) \\ \pi_* \downarrow & & \downarrow \\ \mathrm{GW}(\mathbb{R}) & \longrightarrow & H^0(*, \mathbb{Z}). \end{array}$$

where the vertical maps are the canonical pushforwards and the horizontal maps are the real cycle class maps.

Proof. π is the composition of a closed immersion $i : X \hookrightarrow \mathbb{P}_k^n$ and the structure map $p : \mathbb{P}_k^n \rightarrow \text{Spec } k$. The algebraic pushforward π_* is the composition $p_* \circ i_*$ and the analogous statement holds for the topological pushforward by classical algebraic topology. By [HWXZ19, Theorem 4.7], the pushforward i_* commutes with real realization. We may thus reduce to the case where $X = \mathbb{P}_k^n$ is projective n -space over a field k . In this case, π_* is an isomorphism by [Fas13, Prop 6.3, Theorem 11.7]. Let $s : \text{Spec } K \rightarrow \mathbb{P}_k^n$ be the closed immersion given by the origin. Since $\pi s = 1$, and the real realization maps commute with the algebraic and topological pushforwards of s by [HWXZ19, Theorem 4.7], the real realization maps also commute with the algebraic and topological pushforwards of π . \square

Proposition 5.9. *Let $A \hookrightarrow \mathbb{R}$ and V be a relatively oriented, rank n vector bundle on a smooth projective n -dimensional scheme X over A . Then $n_{\mathbb{R}}$ and $n_{\mathbb{C}}$ are the Euler numbers of the corresponding real and complex topological vector bundles respectively, i.e., $n_{\mathbb{R}}$ is the topological Euler number of the relatively oriented topological \mathbb{R}^n -bundle $V(\mathbb{R})$ associated to the real points of V on the real n -manifold $X(\mathbb{R})$, and $n_{\mathbb{C}}$ is the analogous topological Euler number on $V(\mathbb{C}) \rightarrow X(\mathbb{C})$.*

Proof. By [HWXZ19, Proposition 6.1], the \mathbb{A}^1 -Euler class $e(V, H\tilde{\mathbb{Z}})$ of $V_{\mathbb{R}}$ in the oriented Chow group $H\tilde{\mathbb{Z}}^{V^*}(X_{\mathbb{R}})$ maps to the topological Euler class of $V(\mathbb{R})$ under the real cycle class map. By Lemma 5.8, it follows that the image of the Euler number $n(V, H\tilde{\mathbb{Z}})$ under the real cycle class map is the topological Euler number of $V(\mathbb{R})$. Under the canonical isomorphism $H^0(*, \mathbb{Z}) \cong \mathbb{Z}$, the real cycle class map $\text{GW}(\mathbb{R}) \rightarrow H^0(*, \mathbb{Z}) \cong \mathbb{Z}$ is the map taking a bilinear form over \mathbb{R} to its signature. It follows that $n_{\mathbb{R}}$ is the topological Euler number of $V(\mathbb{R})$.

Let $\gamma : \widetilde{\text{CH}}^*(X, \det V^*) \cong H\tilde{\mathbb{Z}}^{V^*}(X_{\mathbb{R}}) \rightarrow \text{CH}^*(X) \rightarrow \text{CH}^*(X_{\mathbb{C}}) \rightarrow H^*(X(\mathbb{C}), \mathbb{Z})$ denote the composition of the canonical map to Chow followed by the (usual) cycle class map, and we similarly have

$$\gamma : \text{GW}(\mathbb{R}) \cong \widetilde{\text{CH}}^0(\text{Spec } \mathbb{R}) \rightarrow H^0(\text{Spec } \mathbb{R}(\mathbb{C}), \mathbb{Z}) \cong H^0(*, \mathbb{Z}) \cong \mathbb{Z},$$

which sends the class of a bilinear form in $\text{GW}(\mathbb{R})$ to its rank. The cycle class map is compatible with Chern classes ([Ful84, Proposition 19.1.2]), whence the image of $e(V, H\tilde{\mathbb{Z}})$ under γ is the topological Euler class of $V(\mathbb{C})$. The cycle class map commutes with the relevant pushforwards and pullbacks [Ful84, Corollary 19.2, Example 19.2.1], so we have that $\gamma(n(V, H\tilde{\mathbb{Z}})) = n_{\mathbb{C}}$ is the topological Euler number of $V(\mathbb{C})$. \square

Remark 5.10. If proper pushforwards in algebra and topology commute with real realization, as predicted by [HWXZ19, 4.5 Remark], then Proposition 5.9 holds more generally for smooth, proper schemes. It seems that proving this would take us too far afield, however. Alternatively, if $V \otimes \mathbb{R}$ admits a non-degenerate section, then the results of §8 (showing that the Euler number can be computed in terms of Scheja-Storch forms) imply that Proposition 5.9 holds for smooth, proper schemes, arguing as in [SW21, Lemma 5].

Theorem 5.11. *Suppose X is smooth and proper over $\mathbb{Z}[1/2]$. Let V be a relatively oriented vector bundle on X and let V_k denote the base change of V to k for any field k . Then either*

$$(6) \quad n^{\text{GS}}(V_k, \rho) = \frac{n_{\mathbb{C}} + n_{\mathbb{R}}}{2} + \frac{n_{\mathbb{C}} - n_{\mathbb{R}}}{2} \langle -1 \rangle$$

or

$$(7) \quad n^{\text{GS}}(V_k, \rho) = \frac{n_{\mathbb{C}} + n_{\mathbb{R}}}{2} + \frac{n_{\mathbb{C}} - n_{\mathbb{R}}}{2} \langle -1 \rangle + \langle 2 \rangle - 1,$$

where the same formula holds for all fields k of characteristic $\neq 2$.

If instead X is smooth proper over \mathbb{Z} , then (6) holds for any field k (including fields k of characteristic two).

Recall that for the last claim regarding X smooth and proper over \mathbb{Z} , we rely on [CDH⁺20]. See Remark 5.5.

Proof. First assume that $\text{char}(k) \neq 2$. By Corollary 1.4 we have $n^{\text{GS}}(V_k, \rho) = n(V_k, \rho, H\tilde{\mathbb{Z}})$. If the base is \mathbb{Z} , then we learn from Proposition 5.4 that there exist $a, b \in \mathbb{Z}$ (independent of $k!$) such that

$$n(V_k, \rho, H\tilde{\mathbb{Z}}) = a + b \langle -1 \rangle.$$

In $\text{GW}(\mathbb{Z}[1/2]) \hookrightarrow \text{GW}(\mathbb{Q})$ we have the relations

$$\langle -2 \rangle = 1 + \langle -1 \rangle - \langle 2 \rangle \text{ and } 2\langle 2 \rangle = 2$$

(the former because $\langle a \rangle + \langle -a \rangle = \langle 1 \rangle + \langle -1 \rangle$ for any a , and see e.g. [Bac18, Lemma 42] for the latter). Hence if the base is $\mathbb{Z}[1/2]$, there exist $a, b \in \mathbb{Z}, c \in \{0, 1\}$ (independent of the choice of field k !) such that

$$n(V_k, \rho, H\tilde{Z}) = a + b\langle -1 \rangle + c\langle 2 \rangle.$$

If the base is \mathbb{Z} , let us put $c = 0$. By construction we have

$$n_{\mathbb{R}} = \text{sign } n(V_{\mathbb{R}}, \rho, H\tilde{Z}) = (a + c) - b$$

and

$$n_{\mathbb{C}} = \text{rank } n(V_{\mathbb{C}}, \rho, H\tilde{Z}) = (a + c) + b,$$

which determines $a + c$ and b , so that there only remain at most two possible values for $n(V_k, \rho, H\tilde{Z})$.

Now suppose that $\text{char}(k) = 2$ (so that in particular the base is \mathbb{Z}). Since $\langle 1 \rangle = \langle -1 \rangle$ over fields of characteristic 2, we need to show that $n^{\text{GS}}(V_k, \rho) = n_{\mathbb{C}}$. We may as well assume that $k = \mathbb{F}_2$. The rank induces an isomorphism $\text{GW}(\mathbb{F}_2) \cong \mathbb{Z}$ (see e.g. Corollary B.4). Considering the canonical maps

$$\text{GW}(\mathbb{F}_2) \rightarrow \text{KGL}^0(\mathbb{F}_2) \leftarrow \text{KGL}^0(\mathbb{Z}) \rightarrow \text{KGL}^0(\mathbb{Z}[1/2]) \leftarrow \text{GW}(\mathbb{Z}[1/2])$$

in which all but the right-most one are isomorphisms, we get the string of equalities

$$n(V_{\mathbb{F}_2}) = n(V_{\mathbb{F}_2}, \text{KGL}) = n(V_{\mathbb{Z}}, \text{KGL}) = n(V_{\mathbb{Z}[1/2]}, \text{KGL}) = rk(n(V_{\mathbb{Z}[1/2]}, \text{KO})).$$

The result follows. \square

Remark 5.12. The difference of the two values given in Theorem 5.11 is $\langle 2 \rangle - 1$, so the two possibilities can be distinguished by the value of $\text{disc } n(V_k)$ in $k^*/(k^*)^2$ for any field k in which 2 is not a square, such as \mathbb{F}_3 or \mathbb{F}_5 . Such discriminants can be evaluated by a computer, as shown to the second named author by Anton Leyton and Sabrina Pauli.

5.3. Refined Euler classes and numbers.

Definition 5.13 (refined Euler class). Let $E \in \mathcal{SH}(S)$ be a homotopy ring spectrum, $X \in \text{Sch}_S$, $V \rightarrow X$ a vector bundle and $\sigma : X \rightarrow V$ a section with zero scheme $Z = Z(\sigma)$. We denote by $e(V, \sigma) = e(V, \sigma, E) \in E_{\mathbb{Z}}^{V^*}(X)$ the class corresponding to the composite

$$X/X \setminus Z \xrightarrow{\sigma} V/V \setminus 0 \simeq \Sigma^{V^*} \mathbb{1} \xrightarrow{u} \Sigma^{V^*} E|_X \in \mathcal{SH}(X);$$

see Example 4.2.

Remark 5.14. It is clear by construction that refined Euler classes are stable under base change.

Remark 5.15. In [LR20, Definition 3.9] the authors define for an SL-oriented ring spectrum E and a vector bundle $p : V \rightarrow X$ the *canonical Thom class* $th(V) \in E_0^{p^*V}(V)$, which one checks coincides with $e(p^*V, \sigma_0)$, where σ_0 is the tautological section of p^*V . Since $e(V, \sigma) = \sigma^*e(p^*V, \sigma_0)$ we deduce that

$$e(V, \sigma) = \sigma^*th(V).$$

Lemma 5.16. *The “forgetting support” map $E_{\mathbb{Z}}^{V^*}(X) \rightarrow E^{V^*}(X)$ sends $e(V, \sigma)$ to $e(V)$.*

Proof. This follows from the commutative diagram

$$\begin{array}{ccccc} V & \longrightarrow & V/V \setminus 0 & \longrightarrow & \Sigma^{V^*} E \\ \sigma \uparrow & & \sigma \uparrow & & \\ X & \longrightarrow & X/X \setminus Z & & \end{array}$$

Indeed the composite from the bottom left to the top right along the top left represents $e(V)$, by Definition 5.1 (note that σ is a homotopy inverse to the projection $V \rightarrow X$), whereas the composite along the bottom right represents $e(V, \sigma)$ with support forgotten, by Definition 5.13 and Example 4.2. \square

Definition 5.17 (refined Euler number). Suppose that $\pi : X \rightarrow S$ is smoothable lci, $Z \rightarrow S$ is finite and V is relatively oriented (so in particular $rk(V) = rk(L_{\pi})$). Suppose further that E is SL-oriented. Then we put

$$n(V, \sigma, \rho) = n(V, \sigma, \rho, E) = \pi_* e(V, \sigma) \in E^0(S).$$

Corollary 5.18. *Suppose that additionally $\pi : X \rightarrow S$ is smooth and proper. Then $n(V, \sigma, \rho) = n(V, \rho) \in E^0(S)$.*

Proof. Combine Lemma 5.16 and Example 4.7. \square

5.4. Refined Euler classes and the six functors formalism. We now relate our Euler classes to the six functors formalism. The following result shows that our refined Euler class coincides with the one defined by Déglise–Jin–Khan [DJK18, Remark 3.2.10].

Proposition 5.19. *Let $E \in \mathcal{SH}(S)$ be a homotopy ring spectrum, $X \in \text{Sm}_S$, V a vector bundle over X and σ a section of V . Then*

$$e(V, \sigma, E) = \sigma^* z_*(1),$$

where $z : X \rightarrow V$ is the zero section and we use the canonical isomorphism $N_z \simeq V$ to form the pushforward z_* .

Proof. Let $p : V \rightarrow X$ be the projection and $s_0 : V \rightarrow p^*V$ the canonical section. Then $\sigma^*(p^*V, \sigma_0) = (V, s)$ and hence, using Remark 5.14, it suffices to show that $z_*(1) = e(p^*V, \sigma_0, E)$. By [EHK⁺20a, last sentence of §2.1.1] we know that

$$z_* : E^0(X) \simeq E^{z^*p^*V^*+L_z} \rightarrow E_X^{p^*V^*}(V)$$

is the purity equivalence. In the case of the zero section of a vector bundle, it just takes the tautological form

$$[\mathbb{1}, E]_X \simeq [V/V \setminus 0, V/V \setminus 0 \wedge E]_X \simeq [V/V \setminus 0, p^*V/p^*V \setminus 0 \wedge E_V]_V,$$

and hence indeed sends 1 to $e(p^*V, \sigma_0)$. □

It follows from the above that Euler classes of vector bundles are determined by Euler classes of vector bundles with non-degenerate sections.

Example 5.20. Let $V \rightarrow X$ be a vector bundle and σ a section. Suppose $1/2 \in S$. The Euler Class of V in $\text{KW} = \text{KO}[\eta^{-1}]$ is given by the Koszul complex with its canonical symmetric bilinear form. Indeed we may assume that V has a regular section, in which case this follows from the existence of the morphism of cohomology theories $\text{BL}^{\text{naive}} \rightarrow \text{KW}$ and Proposition 2.11.

We deduce that the “Meta-Theorem” of §3.4 holds in this setting, even in a slightly stronger form with supports:

Corollary 5.21. *If σ is a non-degenerate section (i.e. locally given by a regular sequence), then $i_*(1) = e(V, \sigma) \in E_Z^{V^*}(X)$. In particular, forgetting supports (taking the image along $E_Z^{V^*}(X) \rightarrow E^{V^*}(X)$) we have $i_*(1) = e(V) \in E^{V^*}(X)$, i.e. Theorem 3.9 holds in this situation.*

Proof. The second statement follows from the first and Lemma 5.16; hence we shall prove the first. Consider the following cartesian square

$$\begin{array}{ccc} Z & \xrightarrow{i} & X \\ i \downarrow & & z \downarrow \\ X & \xrightarrow{\sigma} & V. \end{array}$$

A well-known consequence of regularity of σ is that this square is tor-independent.⁸ Since i_* is compatible with tor-independent base change (Lemma 4.6(2)), we deduce from Proposition 5.19 that

$$e(V, \sigma, E) = \sigma^* z_*(1) = i_* i^*(1) = i_*(1).$$

This was to be shown. □

It follows that, as explained in §3.4, both the ordinary and refined Euler numbers in E -cohomology can be computed as sums of local indices. The remainder of this paper is mainly concerned with determining these indices, for certain examples of E .

Remark 5.22. These results can be generalized slightly. Fix a scheme S and an SL-oriented ring spectrum $E \in \mathcal{SH}(S)$.

- (1) Let X/S smoothable lci, V/X a vector bundle, relatively oriented in the sense of Definition 1.5. If X/S is in addition proper, then we can transfer the Euler class along the structure morphism (see §4.2.4) as before to obtain an Euler number $n(V, \rho, E) \in E^0(S)$. More generally, without assuming X/S proper, given a section σ with zero scheme Z proper over S , we obtain the refined Euler number $n(V, \sigma, \rho, E)$.

⁸To see this note that $z_*\mathcal{O}_X$ can be resolved by the Koszul complex $K(p^*V, \sigma_0)$ for the tautological section σ_0 of the pullback p^*V of V along $p : V \rightarrow X$. It follows now from the projection formula that $z_*\mathcal{O}_X \otimes^L \sigma_*\mathcal{O}_X \simeq \sigma_*\sigma^*z_*\mathcal{O}_X \simeq \sigma_*\sigma^*K(p^*V, \sigma_0) \simeq \sigma_*K(V, \sigma)$. Since by definition σ locally corresponds to a regular sequence (and σ is affine), the claim follows.

- (2) Let X/S be arbitrary, V a vector bundle, σ a section of V with zero scheme $i : Z \rightarrow X$, and suppose that i is a regular immersion (but σ need not be a non-degenerate section, i.e. Z could have higher than expected dimension). In this case there is an *excess bundle* $\mathcal{E} = \text{cok}(N_{Z/X} \rightarrow V|_Z)$.⁹ A straightforward adaptation of the proof of Corollary 5.21, using the excess intersection formula [DJK18, Proposition 3.3.4], shows that

$$e(V, \sigma, E) = i_*(e(\mathcal{E}, E)).$$

- (3) Putting everything together, let X/S be proper smoothable lci, V a relatively oriented vector bundle and σ a section with zero scheme Z regularly immersed in X . Then

$$n(V, \rho, E) = \sum_{Z' \subset Z} n(\mathcal{E}|_{Z'}, \rho', E).$$

Here the sum is over clopen components Z' of Z and ρ' denotes the induced relative orientation of \mathcal{E} . Note that if σ is non-degenerate on Z' , i.e. $\mathcal{E}|_{Z'} = 0$, then $e(\mathcal{E}|_{Z'}, E) = 1 \in E^0(Z')$ and $n(\mathcal{E}|_{Z'}, \rho', E) = \text{ind}_{Z'}(\sigma)$ as before.

6. d -DIMENSIONAL PLANES ON COMPLETE INTERSECTIONS IN PROJECTIVE SPACE

6.1. Some Euler numbers of symmetric powers on Grassmannians. Grassmannians and flag varieties are smooth and proper over \mathbb{Z} , and the Euler classes of many of their vector bundles have interesting interpretations in enumerative geometry. Computations over \mathbb{R} and \mathbb{C} are available in the literature in connection with enumerative results or accessible with localization techniques in equivariant cohomology [AB84] [GP99]. Integrality of Euler classes, as in Theorem 5.11 and Proposition 5.4, can leverage such results to all fields. We do this now using the $\mathbb{Z}[1/2]$ case of Theorem 5.11 and a characteristic class argument. This is independent of the recent work [CDH⁺20] on Hermitian K-theory over \mathbb{Z} and the characteristic class argument may be of some independent interest. One can alternatively deduce Corollary 6.3 from the \mathbb{Z} case of Theorem 5.11 and [CDH⁺20].

Remark 6.1. Suppose X is a smooth, proper \mathbb{Z} -scheme with geometrically connected fibers and $\text{Pic}X$ torsion free, for example, X a Grassmannian or projective space. For a relatively orientable vector bundle $V \rightarrow X$ defined over \mathbb{Z} , there are at most two isomorphism classes of relative orientations. Namely, by assumption, there is a line bundle $L \rightarrow X$ and isomorphism $\rho : L^{\otimes 2} \xrightarrow{\cong} \omega_{X/\mathbb{Z}} \otimes \det V$. Since $\text{Pic}X$ is torsion free, any relative orientation is an isomorphism $L^{\otimes 2} \xrightarrow{\cong} \omega_{X/\mathbb{Z}} \otimes \det V$, whence two such differ by a global section of $\text{Hom}(L^{\otimes 2}, L^{\otimes 2}) \cong \mathcal{O}(X)^*$. By hypothesis on X , the fibers of the pushforward of \mathcal{O}_X all have rank 1, whence this pushforward is $\mathcal{O}_{\mathbb{Z}}$ and $\mathcal{O}(X)^* \cong \mathbb{Z}^* = \{\pm 1\}$. Thus any relative orientation is isomorphic to ρ or $-\rho$. We then have $n(V, \rho) = \langle -1 \rangle n(V, -\rho)$.

Consequently we suppress the choice of orientation and just write $n(V)$. Beware that this does not mean that every vector bundle is relatively orientable, though!

Remark 6.2. There is a canonical relative orientation for certain classes of vector bundles on Grassmannians $X = \text{Gr}(d, n)$: a point p of X with residue field L corresponds to a dimension $d + 1$ subspace of L^{n+1} . A choice of basis $\{e_0, \dots, e_n\}$ of L^{n+1} such that the span of $\{e_0, \dots, e_d\}$ is p defines canonical local coordinates for an affine chart isomorphic to $\mathbb{A}^{(n-d)(d+1)}$ with p as the origin (see for example [KW21, Definition 42]). This defines local trivializations of the tautological and quotient bundles on X , and therefore also of their tensor, symmetric, exterior powers and their duals. A vector bundle V formed from such operations on the tautological and quotient bundles and which is relatively orientable on X inherits a canonical relative orientation ρ such that the local coordinates and trivializations just described are compatible with ρ in the sense of [KW21, Definition 21]. This is described in [KW21, Proposition 45] in a special case, but the argument holds in the stated generality. (One only needs the determinants of the clutching functions to be squares which follows from the relative orientability of V . Together with the explicit coordinates this gives the relative orientation.) This relative orientation has the property that it is defined over \mathbb{Z} and for any very non-degenerate section σ , the data just described gives a system of coordinates in the sense of Definition 2.28.

Corollary 6.3. *Let $d \leq n$ be positive integers, and let $X = \text{Gr}(d, n)$ be the Grassmannian of d -planes \mathbb{P}^d in \mathbb{P}^n . Let $V = \bigoplus_{i=1}^j \text{Sym}^{n_i} S^*$, where S denotes the tautological bundle on X , and n_1, \dots, n_j are*

⁹The map $N_{Z/X} \rightarrow V|_Z$ is always injective, and is an isomorphism precisely if the section is non-degenerate.

positive integers such that $\text{rank } V = \dim X$ and V is relatively orientable, i.e. such that $\sum_{i=1}^j \binom{n_i+d}{d} = (d+1)(n-d)$ and $\sum_{i=1}^j \frac{n_i}{d+1} \binom{n_i+d}{d} + n + 1$ is even. Then

$$n(V) = \frac{n_{\mathbb{C}} + n_{\mathbb{R}}}{2} + \frac{n_{\mathbb{C}} - n_{\mathbb{R}}}{2} \langle -1 \rangle$$

over any ring in which 2 is invertible (where we interpret $n(V)$ as $n(V, KO)$), or any field (where we interpret $n(V)$ as $n^{\text{GS}}(V)$).

Proof. Let $\mathcal{O}(1)$ denote the generator of $\text{Pic } X$ given by the pullback of the tautological bundle under the Plücker embedding. Then $\omega_{X/\mathbb{Z}} \cong \mathcal{O}(-n-1)$ and $\det \text{Sym}^{n_i} \mathcal{S}^* \cong \mathcal{O}(\frac{n_i}{d+1} \binom{n_i+d}{d})$. Thus V is relatively orientable as a bundle over \mathbb{Z} and Remarks 6.1 and 6.2 apply.

By the $\mathbb{Z}[1/2]$ case of Theorem 5.11, it is enough to show that the discriminant of $n(V_{\mathbb{F}_p})$ is trivial for some prime p congruent to 1 mod 4, and such that 2 is not a square. See Remark 5.12. Let $\text{EM}(\mathbb{W})$ denote the Eilenberg–MacLane spectrum of the η -inverted Milnor–Witt sheaves $K_*^{\text{MW}}[\eta^{-1}]$, cf. [Lev19, Remark 3.1], and consider the associated Euler class $e(V, \text{EM}(\mathbb{W}))$. Then $n(V, \text{EM}(\mathbb{W}))$ determines the Witt class of $n(V)$, and hence the discriminant of $n(V)$ as well.

d is even: Suppose that d is even. Let π denote the structure map of X . Since $n(V, \text{EM}(\mathbb{W})) = \pi_* \prod_{i=1}^j e(\text{Sym}^{n_i} \mathcal{S}^*, \text{EM}(\mathbb{W}))$, it is enough to show that $e(\text{Sym}^{n_1} \mathcal{S}^*, \text{EM}(\mathbb{W})) = 0$. By the Jouanolou device, we may assume any vector bundle is pulled back from the universal bundle. It is therefore enough to show the same for the dual tautological bundle on the universal Grassmannian BGL_{d+1} , i.e., let \mathcal{S}_{d+1}^* denote the dual of the tautological bundle on BGL_{d+1} ; we show that $e(\text{Sym}^{n_1} \mathcal{S}_{d+1}^*, \text{EM}(\mathbb{W})) = 0$. By Ananyevskiy’s splitting principle [Ana15, Theorem 6] and its extension due to M. Levine [Lev19, Theorem 4.1], we may show the vanishing of the $\text{EM}(\mathbb{W})$ -Euler class of $\text{Sym}^{n_1} \mathcal{S}_{d+1}^*$ after pullback to

$$BSL_2 \times BSL_2 \times \dots \times BSL_2 \times BSL_1$$

via the map classifying the external Whitney sum of the tautological bundles. Here we use that $d+1$ is odd. This pullback of $\text{Sym}^{n_1} \mathcal{S}_{d+1}^*$ contains the odd-rank summand $\text{Sym}^{n_1} \mathcal{S}_1^*$, and therefore its $\text{EM}(\mathbb{W})$ -Euler class is 0 as desired [Ana15, Lemma 3] [Lev19, Lemma 4.3].

d is odd: Let k be a finite field whose order is prime to $2 \prod_{i=1}^j (n_i)!$, congruent to 1 mod 4 (so -1 is a square), and such that 2 is not a square. By Theorem 5.11, it suffices to show that the discriminant of $n(V, \text{Gr}(d, n)) \in \mathbb{W}(k) \cong \text{GW}(k)/\mathbb{Z}h$ is trivial, cf. Remark 5.12.

Define r in \mathbb{Z} so that $d = 2r + 1$. Let \mathcal{S}_{d+1}^* denote the dual tautological bundle on BSL_{d+1} and let p_1, \dots, p_r, p_{r+1} and e in $\text{EM}(\mathbb{W})^*(BSL_{d+1})$ denote its Pontryagin and Euler classes respectively. (Often one would let p_i be the Pontryagin classes of the tautological bundle, not its dual, but this is more convenient here.) By Lemma 6.5, $e(\text{Sym}^{n_1} \mathcal{S}_{d+1}^*, \text{EM}(\mathbb{W}))$ is in the image of $\mathbb{Z}[p_1, \dots, p_r, e] \rightarrow \text{EM}(\mathbb{W})^*(BSL_{d+1})$. (Note that we have omitted p_{r+1} as $p_{r+1} = e^2$ [Ana15, Corollary 3].) Therefore $e(V, \text{Gr}(d, n))$ can be expressed as a polynomial with integer coefficients in the Pontryagin classes and Euler class of the dual tautological bundle \mathcal{S}^* on $\text{Gr}(d, n)$. By Lemma 6.7, it follows that $\text{disc } n(V, \text{Gr}(d, n)) = 1$ in $k^*/(k^*)^2$ as desired. \square

M. Levine [Lev19] uses the normalizer N of the standard torus of SL_2

$$1 \rightarrow \left\{ \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \right\} \rightarrow N \rightarrow \left\{ \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\} \rightarrow 1$$

and bundles $\tilde{O}(a)$ and $\tilde{O}^-(a)$ for a in \mathbb{Z} corresponding to the representations

$$\begin{aligned} \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} &\mapsto \begin{pmatrix} t^a & 0 \\ 0 & t^{-a} \end{pmatrix} \\ \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} &\mapsto \begin{pmatrix} 0 & 1 \\ (-1)^a & 0 \end{pmatrix} \end{aligned}$$

and

$$\begin{aligned} \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} &\mapsto \begin{pmatrix} t^a & 0 \\ 0 & t^{-a} \end{pmatrix} \\ \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} &\mapsto \begin{pmatrix} 0 & -1 \\ (-1)^{a+1} & 0 \end{pmatrix}, \end{aligned}$$

respectively, to compute characteristic classes, and we use his technique. We will use the notation $\tilde{O}^{(-)}(a)$ to mean either $\tilde{O}(a)$ and $\tilde{O}^-(a)$ when a claim holds for both possibilities. We likewise use the $\text{EM}(\mathbb{W})$ -Pontryagin (or Borel) classes of a vector bundle with trivialized determinant of Panin and Walter

[PW10b]. See [Ana15, Introduction, Section 3] or [Lev19, Section 3] [Wen20, Section 2] for background on these classes.

Let $e_i \in \text{EM}(\mathbb{W})^*(N^{r+1})$ denote the pullback of $e(S_2^*, \text{EM}(\mathbb{W}))$ under the i th projection $BN^{r+1} \rightarrow BN$ composed with the canonical map $BN \rightarrow \text{BSL}_2$.

Given vector bundles V and E on schemes X and Y , respectively, let $V \boxtimes E$ denote the vector bundle on $X \times Y$ given by the tensor product of the pullbacks of V and E .

Lemma 6.4. *Suppose $r \geq 1$ and a_1, \dots, a_{r+1} are integers, and that our base scheme is a field k with characteristic not dividing $2 \prod_{i=1}^{r+1} a_i$. The Euler class $e(\boxtimes_{i=1}^{r+1} \tilde{O}^{(-)}(a_i), \text{EM}(\mathbb{W}))$ and $\text{EM}(\mathbb{W})$ -Pontryagin classes $p_j(\boxtimes_{i=1}^{r+1} \tilde{O}^{(-)}(a_i), \text{EM}(\mathbb{W}))$ for $j = 1, \dots, 2^r$ are in the image of the map*

$$\mathbb{Z}[e_1^2, \dots, e_{r+1}^2] \rightarrow \text{EM}(\mathbb{W})^*(BN^{r+1}).$$

Proof. Proceed by induction on r . By Ananyevskiy's splitting principle [Ana15, Theorem 6], there exists a map $\pi : Y \rightarrow BN^r$ such that the pullback of $\boxtimes_{i=2}^{r+1} \tilde{O}^{(-)}(a_i)$ is a direct sum of rank 2 bundles V_i on Y ,

$$\pi^* \boxtimes_{i=2}^{r+1} \tilde{O}^{(-)}(a_i) \cong \bigoplus_{i=1}^{2^{r-1}} V_i,$$

and the map

$$\text{EM}(\mathbb{W})^*(BN \times BN^r) \rightarrow \text{EM}(\mathbb{W})^*(BN \times Y)$$

is injective. We pull back the vector bundle $\boxtimes_{i=1}^{r+1} \tilde{O}^{(-)}(a_i)$ along the map $1_{BN} \times \pi : BN \times Y \rightarrow BN^{r+1}$ and obtain an isomorphism

$$(8) \quad (1_{BN} \times \pi)^* \boxtimes_{i=1}^{r+1} \tilde{O}^{(-)}(a_i) \cong \bigoplus_{i=1}^{2^{r-1}} (\tilde{O}^{(-)}(a_1) \boxtimes V_i).$$

By [Lev19, Theorem 7.1], and the equality $e(S_2, \text{EM}(\mathbb{W}))^2 = e(S_2^*, \text{EM}(\mathbb{W}))^2$ which follows from [Lev20, Theorem 11.1],

$$(9) \quad e(\tilde{O}^{(-)}(a_1), \text{EM}(\mathbb{W}))^2 = a_1^2 e(S_2^*, \text{EM}(\mathbb{W}))^2.$$

Since $\tilde{O}^{(-)}(a_1)$ and V_i both have rank 2, [Lev19, Proposition 9.1] and Equation (9) imply that

$$(10) \quad e(\tilde{O}^{(-)}(a_1) \boxtimes V_i, \text{EM}(\mathbb{W})) = a_1^2 e_1^2 - e(V_i, \text{EM}(\mathbb{W}))^2$$

and

$$(11) \quad p_1(\tilde{O}^{(-)}(a_1) \boxtimes V_i, \text{EM}(\mathbb{W})) = 2(a_1^2 e_1^2 + e(V_i, \text{EM}(\mathbb{W}))^2).$$

This establishes the claim when $r = 1$, because $p_2 = e^2$.

We now assume the claim holds for $r - 1$. By Equation 8,

$$(12) \quad e((1_{BN} \times \pi)^* \boxtimes_{i=1}^{r+1} \tilde{O}^{(-)}(a_i), \text{EM}(\mathbb{W})) = \prod_{i=1}^{2^{r-1}} e(\tilde{O}^{(-)}(a_1) \boxtimes V_i, \text{EM}(\mathbb{W}))$$

and

$$(13) \quad p((1_{BN} \times \pi)^* \boxtimes_{i=1}^{r+1} \tilde{O}^{(-)}(a_i), \text{EM}(\mathbb{W})) = \prod_{i=1}^{2^{r-1}} p(\tilde{O}^{(-)}(a_1) \boxtimes V_i, \text{EM}(\mathbb{W})),$$

where p denotes the total Pontryagin class.

Combining Equations (10) and (12) shows that

$$e((1_{BN} \times \pi)^* \boxtimes_{i=1}^{r+1} \tilde{O}^{(-)}(a_i), \text{EM}(\mathbb{W})) = \sum_{i=0}^{2^{r-1}} (a_1 e_1)^{2i} \sigma_{2^{r-1}-i}(e(V_1, \text{EM}(\mathbb{W}))^2, \dots, e(V_{2^{r-1}}, \text{EM}(\mathbb{W}))^2),$$

where σ_i denotes the i th elementary symmetric function. Since the V_i have rank 2,

$$e(V_1, \text{EM}(\mathbb{W}))^2 = p_1(V_1, \text{EM}(\mathbb{W})),$$

and the Whitney sum formula for Pontryagin classes implies that

$$p_i(\bigoplus_{j=1}^{2^{r-1}} V_j, \text{EM}(\mathbb{W})) = \sigma_i(e(V_1, \text{EM}(\mathbb{W}))^2, \dots, e(V_{2^{r-1}}, \text{EM}(\mathbb{W}))^2).$$

Since $\bigoplus_{j=1}^{2^{r-1}} V_j \cong \pi^* \boxtimes_{i=2}^{r+1} \tilde{O}^{(-)}(a_i)$, it follows by induction that $e(\boxtimes_{i=1}^{r+1} \tilde{O}^{(-)}(a_i), \text{EM}(\mathbb{W}))$ is in the image of $\mathbb{Z}[e_1^2, \dots, e_{r+1}^2]$.

Combining equations (11) (10) and (13), we have

$$p((1_{BN} \times \pi)^* \boxtimes_{i=1}^{r+1} \tilde{O}^{(-)}(a_i), \text{EM}(\mathbb{W})) = \prod_{i=1}^{2^{r-1}} (1 + 2(a_1^2 e_1^2 + e(V_i, \text{EM}(\mathbb{W}))^2) + (a_1^2 e_1^2 - e(V_i, \text{EM}(\mathbb{W}))^2)^2).$$

Because the elementary symmetric polynomials generate all symmetric polynomials, it follows that $p((1_{BN} \times \pi)^* \boxtimes_{i=1}^{r+1} \tilde{O}^{(-)}(a_i), \text{EM}(\mathbb{W}))$ is in the image of

$$\mathbb{Z}[e_1^2, \sigma_i(e(V_1, \text{EM}(\mathbb{W}))^2), \dots, e(V_{2^{r-1}}, \text{EM}(\mathbb{W}))^2) : i = 1, \dots, 2^{r-1}].$$

As above, these elementary symmetric functions are the Pontryagin classes of the pullback of $\boxtimes_{i=2}^{r+1} \tilde{O}^{(-)}(a_i)$, finishing the proof by induction. \square

Let p_1, \dots, p_r and e in $\text{EM}(\mathbb{W})^*(B\text{SL}_{d+1})$ denote the Pontryagin and Euler classes respectively of the dual tautological bundle S_{d+1}^* on $B\text{SL}_{d+1}$.

Lemma 6.5. *Let $d = 2r+1$ be an odd integer and let n be a positive integer. Let k be a field of characteristic not dividing $2n!$. Then $e(\text{Sym}^n S_{d+1}^*, \text{EM}(\mathbb{W}))$ is in the image of $\mathbb{Z}[p_1, \dots, p_r, e] \rightarrow \text{EM}(\mathbb{W})^*(B\text{SL}_{d+1})$.*

Proof. Let f denote the composite

$$BN^{r+1} \rightarrow B\text{SL}_2^{r+1} \rightarrow B\text{SL}_{d+1}$$

of the $(r+1)$ -fold product of the canonical map $BN \rightarrow B\text{SL}_2$ with the map $B\text{SL}_2^{r+1} \rightarrow B\text{SL}_{d+1}$ classifying the external direct sum $\oplus_{i=1}^{r+1} S_2$. There is an isomorphism

$$(14) \quad f^* \text{Sym}^n S_{d+1}^* \cong \bigoplus_{\substack{(a_1, \dots, a_{r+1}) \in \mathbb{Z}_{\geq 0}^{r+1} \\ \sum_i a_i = n}} \boxtimes_{i=1}^r \text{Sym}^{a_i} \tilde{O}(1).$$

By inspection, the symmetric powers of the tautological bundle $\tilde{O}(1)$ on BN split into a sum of bundles of rank ≤ 2 , cf. [Lev19, p. 38]:

$$(15) \quad \text{Sym}^a \tilde{O}(1) \cong \begin{cases} \bigoplus_{l=0}^b \tilde{O}^{(-1)^l}(i-2l), & \text{when } a = 2b+1 \text{ is odd} \\ \bigoplus_{l=0}^{b-1} \tilde{O}^{(-1)^l}(i-2l) \oplus \mathcal{O}, & \text{when } a = 2b \text{ is even and } b \text{ is even} \\ \bigoplus_{l=0}^{b-1} \tilde{O}^{(-1)^l}(i-2l) \oplus \gamma, & \text{when } a = 2b \text{ is even and } b \text{ is odd,} \end{cases}$$

where γ is the line bundle corresponding to the representation $N \rightarrow \text{GL}_1$ sending the torus to 1 and $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ to -1 .

Combining Equations (14) and (15), we can decompose $f^* \text{Sym}^n S_{d+1}^*$ into a direct sum with summands which have various numbers of factors of rank 2. Separate these summands into those with at least two rank 2 factors and those with only one rank 2 factor, if any of the latter sort appear. (This occurs when we can take all but one a_i to be even.) The direct sum of the latter such terms can alternatively be expressed as a sum of pullbacks of $\text{Sym}^{a_i} \tilde{O}(1)$ under some projection $N^{r+1} \rightarrow N$ tensored with some γ 's pulled back from other projections. We may ignore the factors of γ by [Lev20, §10 p. 78 (2)] because $e(\gamma) = 0$ as γ is a bundle of odd rank. Since \tilde{O} is rank 2, and the characteristic of k does not divide $2a_i$, we may apply [Lev19, Theorem 8.1] and conclude that $e(\text{Sym}^{a_i} \tilde{O}(1))$ is an integer multiple of a power of $e(\tilde{O}(1))$. Since the summands are symmetric under the permutation action of the symmetric group on $r+1$ letters on BN^{r+1} , it follows that the Euler class of these summands is an integer multiple of a power of e .

We now consider the Euler class of the rest of the summands. Namely, it suffices to show that the Euler class ϵ_1 of the summands with at least two rank 2 factors is also in the image of $\mathbb{Z}[p_1, \dots, p_r, e]$. We may again ignore the factors of γ , as these do not change the Euler class. By Lemma 6.4, ϵ_1 is the image of an element of $\mathbb{Z}[e_1^2, \dots, e_{r+1}^2]$. Moreover, because each tuple (a_1, \dots, a_{r+1}) of the direct sum occurs in every permutation, we may choose an element of $\mathbb{Z}[e_1^2, \dots, e_{r+1}^2]$ which is invariant under the permutation action of the symmetric group on $r+1$ letters and which maps to ϵ_1 . Thus, ϵ_1 is in the image of the map

$$\mathbb{Z}[\sigma_1(e_1^2, \dots, e_{r+1}^2), \dots, \sigma_{r+1}(e_1^2, \dots, e_{r+1}^2)] \rightarrow \text{EM}(\mathbb{W})^*(BN^{r+1}),$$

where σ_i denotes the i th elementary symmetric polynomial. Since $\sigma_i((e_1^2, \dots, e_{r+1}^2))$ is the pullback to BN of $p_i(S_{d+1}^* \rightarrow B\text{SL}_{d+1}, \text{EM}(\mathbb{W}))$, we have that ϵ_1 is in the image of $\mathbb{Z}[p_1, \dots, p_r, e]$ as desired. \square

The Pontryagin and Euler classes of $S_{d+1}^* \rightarrow \text{Gr}(d, n)$ are pulled back from those of $S_{d+1}^* \rightarrow B\text{GL}_{d+1}$. The $\text{EM}(\mathbb{W})^*$ -cohomology and twisted cohomology of $B\text{GL}_{d+1}$ injects into that of $B\text{SL}_{d+1}$,

$$\text{EM}(\mathbb{W})^*(B\text{GL}_{d+1}) \oplus \text{EM}(\mathbb{W})^*(B\text{GL}_{d+1}, \det \mathcal{S}) \subseteq \text{EM}(\mathbb{W})^*(B\text{SL}_{d+1}),$$

by [Lev19, Theorem 4.1]. Under this injection, the Pontryagin and Euler classes of $S_{d+1}^* \rightarrow B\text{GL}_{d+1}$ are sent to p_1, \dots, p_r, p_{r+1} and e , respectively, so we will let p_i and e denote the corresponding characteristic

classes of $\mathcal{S}^* \rightarrow \text{BGL}_{d+1}$ and $\mathcal{S}_{d+1}^* \rightarrow \text{Gr}(d, n)$ as well. Let $\pi : \text{Gr}(d, n) \rightarrow \text{Spec } k$ denote the structure map, and

$$\pi_* : \text{EM}(\mathbb{W})^*(\text{Gr}(d, n), (\det \mathcal{S}_{d+1}^*)^{\otimes -(n+1)}) \rightarrow \mathbb{W}(k)$$

the induced push forward on $\text{EM}(\mathbb{W})^*$ -cohomology.

Lemma 6.6. *Let $d = 2r+1$ be odd. For any non-negative integers a_1, \dots, a_{r+1}, b such that $\sum (4i)a_i + b(d+1) = (d+1)(n-d)$ and $b \equiv n+1 \pmod{2}$, the monomial $e^b \prod_{i=1}^{r+1} p_i^{a_i}$ in $\text{EM}(\mathbb{W})^*(\text{Gr}(d, n), (\det \mathcal{S}_{d+1}^*)^{\otimes b})$ is in the image of $\mathbb{Z}[e]$, or in other words, there exists c in \mathbb{Z} such that $e^b \prod_{i=1}^{r+1} p_i^{a_i} = ce^{n-d}$.*

Proof. Let Q denote the quotient bundle on $\text{Gr}(d, n)$, defined by the short exact sequence

$$0 \rightarrow \mathcal{S}_{d+1} \rightarrow \mathcal{O}^{n+1} \rightarrow Q \rightarrow 0.$$

In particular, the rank of Q is $n-d$. The non-vanishing Pontryagin classes of \mathcal{S}_{d+1} are p_1, \dots, p_r, p_{r+1} with $e^2 = p_{r+1}$. Define s so that $n-d = 2s$ or $n-d = 2s+1$ depending on whether $n-d$ is odd or even. Let $p_1^\perp, \dots, p_s^\perp$ denote the non-vanishing Pontryagin classes of the dual to the quotient bundle Q^* on $\text{Gr}(d, n)$. By [Ana15, Lemma 15],

$$(1 + p_1 + \dots + p_{r+1})(1 + p_1^\perp + \dots + p_s^\perp) = 1$$

in $\mathbb{W}^*(\text{Gr}(d, n))$. Setting the notation A for the ring

$$A = \mathbb{Z}[p_1, \dots, p_r, p_{r+1}, p_1^\perp, \dots, p_s^\perp] / \langle (1 + p_1 + \dots + p_r + p_{r+1})(1 + p_1^\perp + \dots + p_s^\perp) - 1 \rangle,$$

we therefore have a homomorphism $\tau : A \rightarrow \text{EM}(\mathbb{W})^*(\text{Gr}(d, n))$. There is a canonical isomorphism

$$\mathbb{H}^*(\mathbb{C} \text{Gr}(r, r+s); \mathbb{Z}) \cong A,$$

where $\mathbb{H}^*(\mathbb{C} \text{Gr}(r, r+s); \mathbb{Z})$ denotes the singular cohomology of the \mathbb{C} -manifold associated to the \mathbb{C} -points of the Grassmannian $\text{Gr}(r, r+s)$, sending the i th Chern class of the dual tautological bundle to p_i . The top dimensional singular cohomology $\mathbb{H}^{(r+1)s}(\mathbb{C} \text{Gr}(r, r+s); \mathbb{Z})$ is isomorphic to \mathbb{Z} by Poincaré duality. Under our chosen isomorphism, the monomial p_{r+1}^s corresponds to the top Chern class $c_{(r+1)s}(\mathcal{S}_{r+1}^{\oplus s} \rightarrow \mathbb{C} \text{Gr}(r, r+s))$ of the direct sum of s -copies of the dual tautological bundle, which is a generator (with the usual \mathbb{C} -orientations, $c_{(r+1)s}(\mathcal{S}_{r+1}^{\oplus s} \rightarrow \mathbb{C} \text{Gr}(r, r+s))$ counts the number of linear subspaces of dimension r in a complete intersection of s linear hypersurfaces in $\mathbb{C}\mathbb{P}^{r+s}$, and this number is 1, cf. Remark 6.9 and Lemma 6.7). Therefore, for any monomial $\prod_{i=1}^{r+1} p_i^{a_i}$ with $\sum_{i=1}^{r+1} ia_i = (r+1)s$, there is an integer c' such that

$$(16) \quad \prod_{i=1}^{r+1} p_i^{a_i} = c' p_{r+1}^s$$

in $\mathbb{H}^*(\mathbb{C} \text{Gr}(r, r+s); \mathbb{Z})$.

Since d is odd, $n+1 \equiv n-d \pmod{2}$, and therefore $b \equiv n-d \pmod{2}$. Note that if $n-d$ is odd, $b \geq 0$. We may then define a non-negative integer b' by the rule

$$b' = \begin{cases} b/2, & \text{if } n-d \equiv 0 \pmod{2} \\ (b-1)/2, & \text{if } n-d \equiv 1 \pmod{2}. \end{cases}$$

With this notation, $n-d-b = 2(s-b')$. Thus

$$\sum (4i)a_i = (d+1)(n-d-b) = (d+1)2(s-b') = 4(r+1)(s-b'),$$

whence

$$\sum ia_i = (r+1)(s-b').$$

By Equation (16), there is an integer c so that we have the equality

$$p_{r+1}^{b'} \prod_{i=1}^{r+1} p_i^{a_i} = cp_{r+1}^s$$

in $\mathbb{H}^*(\mathbb{C} \text{Gr}(r, r+s); \mathbb{Z})$. Applying τ , we see that

$$e^{2b'} \prod_{i=1}^{r+1} p_i^{a_i} = c' e^{2s},$$

which implies the claim, either immediately in the case that $n-d$ is even, or by multiplying by e if $n-d$ is odd. \square

Lemma 6.7. *Let d be odd. Suppose k is a finite field such that -1 is a square. For any non-negative integers a_1, \dots, a_{r+1}, b such that $b(d+1) + \sum_{i=1}^{r+1} 4ia_i = (d+1)(n-d)$ and $b \equiv n+1 \pmod{2}$, the pushforward $\pi_*(e^b \prod_{i=1}^{r+1} p_i^{a_i})$ has trivial discriminant.*

Remark 6.8. The condition $b \equiv n+1 \pmod{2}$ ensures that $e^b \prod_{i=1}^{r+1} p_i^{a_i}$ lies in the appropriate twist of the Witt-cohomology of $\mathrm{Gr}(d, n)$, i.e. $e^b \prod_{i=1}^{r+1} p_i^{a_i}$ is in $\mathrm{EM}(\mathbb{W})^*(\mathrm{Gr}(d, n), \omega_{\mathrm{Gr}(d, n)}/k)$, as opposed to $\mathrm{EM}(\mathbb{W})^*(\mathrm{Gr}(d, n))$, so that we may apply π_* . The condition on the sum $b(d+1) + \sum_{i=1}^{r+1} 4ia_i$ ensures that $e^b \prod_{i=1}^{r+1} p_i^{a_i}$ lies in the $(d+1)(n-d)$ -degree $\mathrm{EM}(\mathbb{W})^*$ -cohomology of $\mathrm{Gr}(d, n)$, so the codomain of π_* is $\mathbb{W}(k)$.

Proof. By Lemma 6.6, it suffices to show that $\mathrm{disc} \pi_* e^{n-d} = 1$. We may identify $\pi_* e^{n-d}$ with the Euler number of $\bigoplus_{j=1}^{n-d} \mathcal{S}^*$,

$$\pi_* e^{n-d} = n(\bigoplus_{j=1}^{n-d} \mathcal{S}^*, \mathrm{EM}(\mathbb{W})).$$

Let x_0, \dots, x_n be coordinates on projective space $\mathbb{P}_k^n = \mathrm{Proj} k[x_0, \dots, x_n]$. The Euler number $n(\bigoplus_{j=1}^{n-d} \mathcal{S}^*, \mathrm{EM}(\mathbb{W}))$ can be calculated with the section $\sigma = \bigoplus_{i=d+1}^n x_i$ as in §2.4. There is an analogous section $\sigma_{\mathbb{Z}} = \bigoplus_{i=d+1}^n x_i$ of $\bigoplus_{j=1}^{n-d} \mathcal{S}_{\mathbb{Z}}^* \rightarrow \mathrm{Gr}(d, n)_{\mathbb{Z}}$ defined over \mathbb{Z} . The zero locus $\sigma_{\mathbb{Z}} = 0$ is the single \mathbb{Z} -point of the Grassmannian associated to the linear subspace of \mathbb{P}_k^n given by $x_{d+1} = x_{d+2} = \dots = x_n = 0$.

The vanishing locus of $\sigma_{\mathbb{Z}}$ is the origin of the affine space $\mathbb{A}_{\mathbb{Z}}^{(d+1)(n-d)} = \mathrm{Spec} \mathbb{Z}[a_{ij} : i = 0, \dots, d, j = d+1, \dots, n] \hookrightarrow \mathrm{Gr}(d, n)$ whose point (a_{ij}) corresponds to the row space of

$$(17) \quad \begin{pmatrix} 1 & 0 & \cdots & 0 & a_{0,d+1} & a_{0,d+2} & \cdots & a_{0,n} \\ 0 & 1 & \cdots & 0 & a_{1,d+1} & a_{1,d+2} & \cdots & a_{1,n} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 & a_{d,d+1} & a_{d,d+2} & \cdots & a_{d,n} \end{pmatrix}.$$

See for example, [EH16, Section 3.2], for a description of these coordinates on this open affine of the Grassmannian. Let e_0, \dots, e_n denote the standard basis of the free module of rank $n+1$. Let $\tilde{e}_0, \dots, \tilde{e}_n$ denote the basis consisting of the row space of (17) followed by $e_{d+1}, e_{d+1}, \dots, e_n$. Let $\tilde{x}_0, \dots, \tilde{x}_n$ denote the dual basis to $\tilde{e}_0, \dots, \tilde{e}_n$. Over this $\mathbb{A}_{\mathbb{Z}}^{(d+1)(n-d)}$, the vector bundle $\mathcal{S}_{\mathbb{Z}}^*$ is trivialized by the basis of sections $\{\tilde{x}_0, \dots, \tilde{x}_d\}$. Then we may interpret σ as a function $\mathbb{A}_{\mathbb{Z}}^{(d+1)(n-d)} \rightarrow \mathbb{A}_{\mathbb{Z}}^{(d+1)(n-d)}$. Namely,

$$x_u((a_{ij})) = \sum_{l=0}^d (x_u(a_{ij}))(\tilde{e}_l) \tilde{x}_l$$

for

$$u = d+1, \dots, n.$$

As a subscheme, $\sigma_{\mathbb{Z}} = 0$ is therefore the zero locus of

$$(x_u(a_{ij}))(\tilde{e}_l) = a_{l,u}$$

for $l = 0, \dots, d$ and $u = d+1, \dots, n$. Thus the subscheme of $\mathrm{Gr}(d, n)_{\mathbb{Z}}$ given by $\{\sigma_{\mathbb{Z}} = 0\}$ is a section of the structure map $\mathrm{Gr}(d, n)_{\mathbb{Z}} \rightarrow \mathrm{Spec} \mathbb{Z}$. In particular, it is finite and étale of rank 1. It follows that the Jacobian of σ (which is described further at the beginning of Section 6.2)

$$\mathrm{Jac} \sigma \in \mathrm{Hom}(\det T \mathrm{Gr}(d, n)|_{\{\sigma_{\mathbb{Z}}=0\}}, \det \bigoplus_{j=1}^{n-d} \mathcal{S}_{\mathbb{Z}}^*|_{\{\sigma_{\mathbb{Z}}=0\}})$$

is nowhere vanishing. Thus under the relative orientation

$$\mathrm{Hom}(\det T \mathrm{Gr}(d, n), \det \bigoplus_{j=1}^{n-d} \mathcal{S}_{\mathbb{Z}}^*) \cong L^{\otimes 2},$$

we have that $\mathrm{Jac} \sigma$ is a nowhere vanishing section of the restriction of $L^{\otimes 2}$. Thus $\langle \mathrm{Jac} \sigma_k \rangle$ is either $\langle -1 \rangle$ or $\langle 1 \rangle$; but $\langle -1 \rangle = 1$ by assumption. Since $n(\bigoplus_{j=1}^{n-d} \mathcal{S}_k^*, \mathrm{EM}(\mathbb{W})) = \langle \mathrm{Jac} \sigma_k \rangle$ (cf. Example 2.21), this proves the claim. \square

6.2. An arithmetic count of the d -planes on a complete intersection in projective space. A complete intersection of hypersurfaces

$$X = \{F_1 = F_2 = \dots = F_j = 0\} \subset \mathbb{P}^n$$

with F_i of degree n_i gives rise to a section σ of $V = \bigoplus_{i=1}^j \mathrm{Sym}^{n_i} \mathcal{S}^*$, defined by $\sigma(\mathbb{P}L) = \bigoplus_{i=1}^j F_i|_L$, where L is any $d+1$ dimensional linear subspace of \mathbb{A}^{n+1} containing the origin, and $\mathbb{P}L$ denotes the corresponding point of $\mathrm{Gr}(d, n)$. The zeros of σ are then precisely the d -planes in X . See for example [DM98]. By [DM98, Théorème 2.1], the closed subscheme $\{\sigma = 0\}$ of $\mathrm{Gr}(d, n)$ is smooth for general X

and either of the expected dimension $(d+1)(n-d) - \text{rank } V = (d+1)(n-d) - \sum_{i=1}^j \binom{n_i+d}{d}$ or empty, with the empty case occurring exactly when one or both of $(d+1)(n-d) - \text{rank } V$ and $n-2r-j$ is less than 0. In particular, when $(d+1)(n-d) - \text{rank } V = 0$, the zeros of σ are isolated and étale over k for a general complete intersection X . The canonical relative orientation (Remark 6.2) of V determines an isomorphism $\text{Hom}(\det T \text{Gr}(d, n), \det V) \cong L^{\otimes 2}$ for a line bundle L on $\text{Gr}(d, n)$. The Jacobian determinant $\text{Jac} \sigma$ at a zero p of σ is an element of the fiber of the vector bundle $\text{Hom}(\det T \text{Gr}(d, n), \det V)$ at p . Choosing any local trivialization of L , we have a well-defined element $\text{Jac} \Sigma(p)$ in $k(p)/(k(p)^*)^2$, which can also be computed by choosing a local trivialization of V and local coordinates of $\text{Gr}(d, n)$ compatible with the relative orientation and computing $\text{Jac} \Sigma(p) = \det \left(\frac{\partial \sigma_k}{\partial x_i} \right)$.

Corollary 6.9. *Let $X = \{F_1 = F_2 = \dots = F_j = 0\} \subset \mathbb{P}^n$ be a general complete intersection of hypersurfaces $F_i = 0$ of degree n_i in \mathbb{P}_k^n projective space over a field k . Suppose that $\sum_{i=1}^j \binom{n_i+d}{d} = (d+1)(n-d)$ and $\sum_{i=1}^j \frac{n_i}{d+1} \binom{n_i+d}{d} + n+1$ is even. Then*

$$\sum_{d\text{-planes } \mathbb{P}L \text{ in } X} \text{tr}_{k(\mathbb{P}L)/k} \langle \text{Jac } \sigma(\mathbb{P}L) \rangle = \frac{n_{\mathbb{C}} + n_{\mathbb{R}}}{2} + \frac{n_{\mathbb{C}} + n_{\mathbb{R}}}{2} \langle -1 \rangle$$

where

- $k(\mathbb{P}L)$ denotes the residue field of $\mathbb{P}L$ viewed as a point on the Grassmannian
- $n_{\mathbb{C}}$ (respectively $n_{\mathbb{R}}$) is the topological Euler number of the complex (respectively real) vector bundle associated to the algebraic vector bundle $V = \bigoplus_{i=1}^j \text{Sym}^{n_i} \mathcal{S}^*$ given the canonical relative orientation (6.2)
- and $\text{Jac} \sigma$ is the Jacobian determinant.

Proof. By [DM98, Théorème 2.1], the zeros of σ are isolated and étale over k . It follows [KW21, p.18, Proposition 34] that for a zero of σ corresponding to the d -plane $\mathbb{P}L$, the local index is computed $\text{ind}_{\mathbb{P}L}^{\text{PH}} \sigma = \text{tr}_{k(L)/k} \langle \text{Jac } \sigma(\mathbb{P}L) \rangle$. See Section 2.4 for the definition of the notation ind^{PH} . Thus $n^{\text{PH}}(V, \sigma) = \sum_{d\text{-planes } \mathbb{P}L \text{ in } X} \text{tr}_{k(L)/k} \langle \text{Jac } \sigma(\mathbb{P}L) \rangle$ by Definition 2.24. Corollary 6.3 computes $n^{\text{PH}}(V, \sigma)$. Proposition 5.9 shows that $n_{\mathbb{R}}$ and $n_{\mathbb{C}}$ are the claimed topological Euler numbers. \square

Remark 6.10. Note that Corollary 6.9 is a weighted count of the dimension d hyperplanes on the complete intersection X , depending only on n_1, \dots, n_j and not on the choice of polynomials F_1, \dots, F_j as long as these are chosen generally.

Example 6.11. Examples where Corollary 6.9 applies include: i) lines on a degree $2n-1$ hypersurface of dimension n , ii) 3-planes on a degree d hypersurface of dimension $2 + \frac{1}{3} \binom{d+3}{3}$, when this is an integer. iii) Lines on a complete intersection of two degree $n-2$ polynomials in \mathbb{P}^n for n odd.

Matthias Wendt's oriented Schubert calculus shows that enriched intersections of Schubert varieties are determined by the \mathbb{R} and \mathbb{C} realizations in the same manner [Wen20, Theorem 8.6], as well as giving enumerative applications [Wen20, Section 9].

For $d=j=1$ and $n_1=3$, Corollary 6.3 is work of Kass and the second named author [KW21] over a general field. For $d=1$ and general j and n_i , it is work of M. Levine [Lev19] over a perfect field either of characteristic 0 or of characteristic prime to 2 and the odd n_i s. Our result eliminates the assumption on the characteristic, and generalizes to arbitrary relatively orientable d, n, n_1, \dots, n_j .

In order to obtain an enumerative theorem whose statement is independent of \mathbb{A}^1 -homotopy theory, one needs an arithmetic-geometric interpretation of the local indices:

Question 6.12. Can the local indices $\text{ind}_{\mathbb{P}L}^{\text{PH}} \sigma = \text{tr}_{k(L)/k} \langle \text{Jac } \sigma(\mathbb{P}L) \rangle$ be expressed in terms of the arithmetic-geometry of the d -plane $\mathbb{P}L$ on X ?

Such expressions are available over \mathbb{R} for $d=j=1$ [FK21], and over a field k of characteristic not 2, for lines on a cubic surface [KW21], lines on a quintic 3-fold [Pau20], and points on a complete intersection of hypersurfaces [McK21]. S. Pauli has interesting observations on such results for lines on the complete intersection of two cubics in \mathbb{P}^5 . Dropping the assumption that the zeros of σ are isolated, she can compute contributions from infinite families of lines on a quintic 3-fold in some cases [Pau20]. An alternative point of view in terms of (S)pin structures for $d=j=1$ and $n_1=5$, as well as computations of the real Euler number is discussed [Sol06, Example 1.6, Theorem 8.8].

Example 6.13. The computations of the Euler classes of \mathbb{C} and \mathbb{R} -points in Finashin and Kharlamov's paper [FK15, p. 190] imply the following enriched counts of 3-planes on hypersurfaces over any field k .

$n(\mathrm{Sym}^3 \mathcal{S}^* \rightarrow \mathrm{Gr}(3, 8)) = 160839\langle 1 \rangle + 160650\langle -1 \rangle$ corresponds to an enriched count of 3-planes in a 7-dimensional cubic hypersurface. Namely, for a general degree 3 polynomial F in 9 variables, the corresponding cubic hypersurface $X \subset \mathbb{P}^8$ contains finitely many 3-planes as discussed above and

$$\sum_{3\text{-planes } P \subset X} \mathrm{tr}_{k(L)/k} \langle \mathrm{Jac} \sigma_F(P) \rangle = 160839\langle 1 \rangle + 160650\langle -1 \rangle,$$

where σ_F is the section of $\mathrm{Sym}^3 \mathcal{S}^*$ defined by $\sigma_F[P] = F|_P$.

Similarly, $n(\mathrm{Sym}^5 \mathcal{S}^* \rightarrow \mathrm{Gr}(3, 17)) =$

$$32063862647475902965720976420325\langle 1 \rangle + 32063862647475902965683320692800\langle -1 \rangle$$

corresponds to an enriched count of 3-planes in a 16-dimensional degree 5-hypersurface.

7. INDICES OF SECTIONS OF VECTOR BUNDLES AND \mathbb{A}^1 -DEGREES

7.1. \mathbb{A}^1 -degrees. Recall the following.

Definition 7.1 (local \mathbb{A}^1 -degree). Let S be a scheme, $X \in \mathrm{Sch}_S$ and $F : X \rightarrow \mathbb{A}_S^n$ be a morphism. We say that F has an isolated zero at $Z \subset X$ if Z is a clopen subscheme of $Z(F)$ such that Z/S is finite.

Now let $X \subset \mathbb{A}_S^n$ be open and suppose that F has an isolated zero at $Z \subset X$. We define the *local \mathbb{A}^1 -degree of F at Z*

$$\mathrm{deg}_Z(F) \in [\mathbb{P}^n/\mathbb{P}^{n-1}, \mathbb{P}^n/\mathbb{P}^{n-1}]_{\mathcal{S}\mathcal{H}(S)} \simeq [\mathbb{1}, \mathbb{1}]_{\mathcal{S}\mathcal{H}(S)}$$

as the morphism corresponding to the unstable map

$$\mathbb{P}^n/\mathbb{P}^{n-1} \rightarrow \mathbb{P}^n/\mathbb{P}^n \setminus Z \simeq X/X \setminus Z \hookrightarrow X/X \setminus Z(F) \xrightarrow{F} \mathbb{A}^n/\mathbb{A}^n \setminus 0 \simeq \mathbb{P}^n/\mathbb{P}^{n-1}.$$

Here we use that by assumption $Z(F) \simeq Z \amalg Z'$, and hence $X/X \setminus Z(F) \simeq X/X \setminus Z \amalg X/X \setminus Z'$.

Example 7.2. If $S = \mathrm{Spec}(k)$ is the spectrum of a field, then an isolated zero $z \in \mathbb{A}_k^n$ of $F : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$ in the usual sense is also an isolated zero $\{z\} \subset \mathbb{A}_k^n$ in the above sense, and $\mathrm{deg}_z(F) \in \mathrm{GW}(k)$ is the usual local \mathbb{A}^1 -degree of [KW19, Definition 11].

Lemma 7.3. *Let $X, Y \in \mathrm{Sm}_S$ and (Z, U, ϕ, g) be an equationally framed correspondence from X to Y [EHK⁺17, Definition 2.1.2]; in other words $Z \subset \mathbb{A}_X^n$, U is an étale neighbourhood of Z , $g : U \rightarrow Y$ and $\phi : U \rightarrow \mathbb{A}^n$ is a framing of Z . Then the following two morphisms are stably homotopic*

$$T^n \wedge X_+ \simeq (\mathbb{P}^1)^{\wedge n} \wedge X_+ \rightarrow (\mathbb{P}^1)_{X_+}^{\times n} / (\mathbb{P}^1)_{X_+}^{\times n} \setminus Z \simeq U/U \setminus Z \xrightarrow{\phi, g} T^n \wedge Y_+$$

$$T^n \wedge X_+ \simeq \mathbb{P}^n/\mathbb{P}^{n-1} \wedge X_+ \rightarrow \mathbb{P}_X^n/\mathbb{P}_X^n \setminus Z \simeq U/U \setminus Z \xrightarrow{\phi, g} T^n \wedge Y_+.$$

Proof. For $E \in \mathcal{S}\mathcal{H}(S)$, precomposition with the first morphism (desuspended by T^n) induces a map $E(Y) \rightarrow E(X)$ known as the *Voevodsky transfer*. Precomposition with the second map induces an “alternative Voevodsky transfer”. It suffices (by the Yoneda lemma) to show that these transfer maps have the same effect (even just on π_0), for every E . In [EHK⁺20a, Theorem 3.2.11], it is shown that the Voevodsky transfer coincides with a further construction known as the *fundamental transfer*. In that proof, all occurrences of $(\mathbb{P}^1)^{\wedge n}$ can be replaced by $\mathbb{P}^n/\mathbb{P}^{n-1}$; one deduces that the alternative Voevodsky transfer also coincides with the fundamental transfer.

The result follows. □

Corollary 7.4. *Let $U \subset \mathbb{A}_S^n$, $F : U \rightarrow \mathbb{A}^n$ have an isolated zero along $Z \subset U$. Then $\mathrm{deg}_Z(F) \in [\mathbb{1}, \mathbb{1}]_S$ is the same as the endomorphism given by the equationally framed correspondence defined by F .*

Proof. By definition, $\mathrm{deg}_Z(F)$ is given by the second morphism of Lemma 7.3, whereas the equationally framed correspondence is given by the first morphism. The result follows. □

Recall also the following.

Proposition 7.5 ([EHK⁺20a], Theorem 3.3.10). *Let $\alpha : S \xleftarrow{\varpi, \tau} Z \rightarrow S$ be a tangentially framed correspondence from S to S over S . Then the trivialization τ of $L_{Z/S}$ induces a transfer map*

$$\varpi_* : [\mathbb{1}, \mathbb{1}]_Z \rightarrow [\mathbb{1}, \mathbb{1}]_S,$$

where $\varpi : Z \rightarrow S$ is the projection. The endomorphism of $\mathbb{1}_S$ corresponding to α is given by $\varpi_*(1)$.

7.2. Main result. Let X/S be smooth, V/X a relatively oriented vector bundle with very non-degenerate section σ and zero scheme Z (which is thus finite over S). Let $Z' \subset Z$ be a clopen component and suppose there are coordinates (ψ, φ, σ') for (V, X, σ, ρ, Z') as in Definition 2.28. Then $\sigma' = (F_1, \dots, F_d)$ determines a function $F : \mathbb{A}_S^d \rightarrow \mathbb{A}_S^d$, and $\varphi(Z')$ is an isolated zero of F .

Theorem 7.6. *Assumptions and notations as above. Let $E \in \mathcal{SH}(S)$ be an SL-oriented ring spectrum with unit map $u : \mathbb{1} \rightarrow E$. Then*

$$\mathrm{ind}_Z(V, \sigma, \rho, E) = u_* \mathrm{deg}_{\varphi(Z')}(F) \in E^0(S).$$

Proof. By Corollary 7.4, Proposition 7.5 and §4.2.2 we have $u_* \mathrm{deg}_{\varphi(Z')}(F) = \mathrm{ind}_Z(\sigma', \sigma', E)$, where σ' is the canonical relative orientation of $\mathcal{O}_{\mathbb{A}^n}^n / \mathbb{A}^n$. The result now follows from Proposition 3.13. \square

Example 7.7. Suppose $S = \mathrm{Spec}(k)$ is a field. Then for well-chosen E (e.g. $E = \mathrm{KO}$ or $E = \mathrm{H}\tilde{\mathbb{Z}}$), the unit map

$$u_* : \mathrm{GW}(k) \simeq [\mathbb{1}, \mathbb{1}]_{\mathcal{SH}(k)} \rightarrow E^0(k)$$

is an isomorphism. We deduce that $\mathrm{ind}_Z(V, \sigma, \rho, E)$ is essentially the same as $\mathrm{deg}_{\varphi(Z)}(F)$.

8. EULER NUMBERS IN KO-THEORY AND APPLICATIONS

As explained in §3.3 and §5.3, we have the motivic ring spectrum KO related to Hermitian K -theory made homotopy invariant, and associated theories of Euler classes and Euler numbers. Using (for example) the construction in §A, we can define KO even if $1/2 \notin S$. There is still a map $\mathrm{GW}(S) \rightarrow \mathrm{KO}(S)$, however we do not know if this is an equivalence, even if S is regular (but we do know this if S is regular and $1/2 \in S$).

Example 8.1. Let $V \rightarrow X$ be a vector bundle and σ a section. Suppose $1/2 \in S$. Then the refined Euler class of V in KO -theory is given by $e(V, \sigma, \mathrm{KO}) = [K(V, \sigma)]$, the class of the Koszul complex. Indeed via Remark 5.15 it suffices to show the analogous result for Thom classes, which is stated on [LR20, p. 34].

Recall that for any lci morphism f we put $\tilde{\omega}_f = \widetilde{\det} L_f$ and $\omega_f = \det L_f$. We show in Proposition B.1 that for $f : X \rightarrow Y$ an lci morphism, we have $f^!(\mathcal{O}) \simeq \tilde{\omega}_f$. Via Proposition B.1, coherent duality (i.e. the adjunction $f_* \dashv f^!$) thus supplies us with a canonical *trace map*

$$\eta_f : f_* \tilde{\omega}_f \simeq f_* f^! \mathcal{O}_Y \rightarrow \mathcal{O}_Y \in D(Y),$$

provided that f is also *proper*. One expects that this can be used to build a map

$$\hat{f}_* : \mathrm{GW}(X, f^! \mathcal{L}) \rightarrow \mathrm{GW}(Y, \mathcal{L}),$$

and moreover that the following diagram commutes

$$\begin{array}{ccc} \mathrm{GW}(X, f^! \mathcal{L}) & \longrightarrow & \mathrm{KO}(X, f^! \mathcal{L}) \\ \hat{f}_* \downarrow & & f_* \downarrow \\ \mathrm{GW}(Y, \mathcal{L}) & \longrightarrow & \mathrm{KO}(Y, \mathcal{L}). \end{array}$$

If we assume that $1/2 \in S$, replace GW by W and KO by KW , then maps \hat{f}_* can be defined and studied using the ideas from §2.2; see also [CH11]. Levine-Raksit [LR20] show that the (modified) diagram commutes provided that X and Y are smooth over a common base with $1/2 \in S$.

If instead we assume that f is finite syntomic, then the analogous result is proved (for $\mathrm{GW} \rightarrow \mathrm{KO}$ and without $1/2 \in S$) in Corollary A.4. This is the only case that we shall use in the rest of this section. Recall the construction of the Scheja-Storch form $\langle - | - \rangle^{\mathrm{SS}}$ from Definition 2.20.

Corollary 8.2. *Let $X \in \mathrm{Sm}_S$, V/X a relatively oriented vector bundle with a very non-degenerate section σ , and Z a clopen component of the zero scheme $Z(\sigma)$. Suppose there exists coordinates (ψ, φ, σ') around Z , as in Definition 2.28.*

Then

$$\mathrm{ind}_Z(\sigma, \rho, \mathrm{KO}) = [\langle - | - \rangle^{\mathrm{SS}}(\varphi(U), \sigma', S)] \in \mathrm{KO}^0(S).$$

Proof. By Proposition 3.13, we may assume that $\psi = \mathrm{id}$ and so on; so in particular $X \subset \mathbb{A}_S^n$. The result now follows from the identification of the transfers in Corollary A.4 (telling us that the index is given by the trace form from coherent duality) and Theorem 2.18 (identifying the coherent duality form with the Scheja-Storch form). \square

Corollary 8.3. *Let S be regular semilocal scheme over a field k of characteristic $\neq 2$.*

- (1) Let $\varpi : S' \rightarrow S$ be a finite syntomic morphism, and τ a trivialization of $L_\varpi \in K(S')$. Then the associated endomorphism of the sphere spectrum over S is given under the isomorphism $[\mathbb{1}, \mathbb{1}]_S \simeq GW(S)$ of [BH17, Theorem 10.12] by the symmetric bilinear form

$$\varpi_*(\mathcal{O}_{S'}) \otimes \varpi_*(\mathcal{O}_{S'}) \rightarrow \varpi_*(\mathcal{O}_{S'}) \xrightarrow{\det(\tau)} \varpi_*(\omega_{S'/S}) \xrightarrow{\eta_\varpi} \mathcal{O}_S.$$

- (2) Let $U \subset \mathbb{A}_S^n$ be open, $F : U \rightarrow \mathbb{A}^n$ have an isolated zero along $Z \subset U$. Then

$$\deg_Z(F) = [\langle - | - \rangle^{\text{SS}}(U, F, S)] \in [\mathbb{1}, \mathbb{1}]_S \simeq GW(S).$$

Proof. The proof of [BH17, Theorem 10.12] shows that the unit map $u : \mathbb{1} \rightarrow \text{KO}$ induces the isomorphism $[\mathbb{1}, \mathbb{1}]_S \simeq GW(S)$. (1) is an immediate consequence of Proposition 7.5, Lemma 4.6(1) and Corollary A.4. Via Corollary 7.4 and Theorem 2.18, (2) is a special case of (1). This concludes the proof. \square

Corollary 8.4. *Let X be essentially smooth over a field k of characteristic $\neq 2$. Let $\varpi : X' \rightarrow X$ be a finite syntomic morphism, and suppose given an orientation $\omega_{X'/X} \xrightarrow{\rho} \mathcal{L}^{\otimes 2}$. Consider the induced transfer*

$$\varpi_* : \text{H}\tilde{\mathbb{Z}}^0(X') \xrightarrow{\rho} \text{H}\tilde{\mathbb{Z}}^0(X', \omega_{X'/X}) \rightarrow \text{H}\tilde{\mathbb{Z}}^0(X) \simeq \underline{GW}(X).$$

Then $\varpi_*(1)$ is given by the image in $\underline{GW}(X)$ of the symmetric bilinear form on

$$\varpi_*(\mathcal{L}) \otimes \varpi_*(\mathcal{L}) \rightarrow \varpi_*(\mathcal{L}^{\otimes 2}) \xrightarrow{\rho} \varpi_*(\omega_{X'/X}) \xrightarrow{\eta_\varpi} \mathcal{O}_X.$$

Proof. Using unramifiedness of \underline{GW} [Mor05, Lemma 6.4.4], we may assume that X is the spectrum of a field. Then X' is semilocal, so $\mathcal{L} \simeq \mathcal{O}$ and we obtain (up to choosing such an isomorphism) $\omega_{X'/X} \xrightarrow{\rho'} \mathcal{O}$. Since $L_{X'/X}$ has constant rank (namely 0), it follows from [BH98, Lemma 1.4.4] that $L_{X'/X} \simeq 0 \in K(X')$. The set of homotopy classes of such trivializations is given by $K_1(X')$, which maps surjectively (via the determinant) onto $\mathcal{O}^\times(X')$. It follows that there exists a trivialization $\tau : 0 \simeq L_{X'/X} \in K(X')$ such that $\det(\tau) = \rho'$. Hence we have a commutative diagram

$$\begin{array}{ccccccc} \varpi_*(\mathcal{L}) \otimes \varpi_*(\mathcal{L}) & \longrightarrow & \varpi_*(\mathcal{L}^{\otimes 2}) & \xrightarrow{\rho} & \varpi_*(\omega_{X'/X}) & \xrightarrow{\eta_\varpi} & \mathcal{O} \\ \parallel & & \parallel & & \parallel & & \parallel \\ \varpi_*(\mathcal{O}) \otimes \varpi_*(\mathcal{O}) & \longrightarrow & \varpi_*(\mathcal{O}) & \xrightarrow{\det \tau} & \varpi_*(\omega_{X'/X}) & \xrightarrow{\eta_\varpi} & \mathcal{O}. \end{array}$$

It follows from Corollary 8.3 that the bottom row is the form $\varpi_*(1)$ arising from the orientation ρ ; by what we just said this is the same as the top row, which is the form we were supposed to obtain.

This concludes the proof. \square

We also point out the following variant.

Corollary 8.5. *Let l/k be a finite extension of fields, $1/2 \in k$. Then Morel's absolute transfer [Mor12, §5.1] $\text{tr}_{l/k} : GW(l, \omega_{l/k}) \rightarrow GW(k)$ is given as follows. Let $\phi : V \otimes_l V \rightarrow l$ be an element of $GW(l)$, $\alpha \in \omega_{l/k}^\times$. Then*

$$\text{tr}_{l/k}(\phi \otimes \alpha) = [V \otimes_k V \rightarrow V \otimes_l V \xrightarrow{\phi} l \xrightarrow{\alpha} \omega_{l/k} \xrightarrow{\eta_{l/k}} k],$$

where $\eta_{l/k} : \omega_{l/k} \rightarrow k$ is the (k -linear) trace map of coherent duality (see §B.1).

Proof. Immediate from [EHK⁺20a, Proposition 4.3.17] (telling us that Morel's transfer coincides with the one from §4.2.4) and Corollary A.4. \square

Remark 8.6. Corollary 8.3(2) generalizes the main result of [KW19], at least for fields of characteristic $\neq 2$.

Remark 8.7. We expect that all of the results in this section extend to fields of characteristic 2 as well. This should be automatic as soon as KO is shown to represent GW in this situation (over regular bases, say).

APPENDIX A. KO VIA FRAMED CORRESPONDENCES

In this section we will construct a strong orientation on KO, and identify some of the transfers. We would like to thank M. Hoyois for communicating these results to us. For another approach to parts of the results in this section see [LÁ17].

We shall make use of the technology of framed correspondences [EHK⁺17]. We write $\text{Corr}^{\text{fr}}(S)$ for the symmetric monoidal ∞ -category of smooth S -schemes and tangentially framed correspondences. Denote by $\text{FSyn}^{\text{or}} \in \text{CAlg}(\mathcal{P}_{\Sigma}(\text{Corr}^{\text{fr}}(S)))$ the stack of finite syntomic schemes Y/X together with a choice of trivialization $\det L_{Y/X} \simeq \mathcal{O}$, with its standard structure of framed transfers. This is constructed in [EHK⁺20b, Example 3.3.4].

Write $\text{Bil} \in \text{CAlg}(\mathcal{P}_{\Sigma}(\text{Sch}_S))$ for the presheaf sending X to the 1-groupoid of pairs (V, ϕ) with $V \rightarrow X$ a vector bundle and $\phi : V \otimes V \rightarrow \mathcal{O}_X$ a non-degenerate, symmetric bilinear form. The commutative monoid structure is given by \otimes . If $f : X \rightarrow Y$ is a finite syntomic morphism then a choice of trivialization $\widetilde{\det} L_f \simeq \omega_f \simeq \mathcal{O}_X$ induces an additive map $\hat{f}_* : \text{Bil}^{\simeq}(X) \rightarrow \text{Bil}^{\simeq}(Y)$; see e.g. §2.2.

Theorem A.1. *There exists a lift $\text{Bil} \in \text{CAlg}(\mathcal{P}_{\Sigma}(\text{Corr}^{\text{fr}}(S)))$ with the transfers given by maps of the form \hat{f}_* , together with a morphism $\text{FSyn}^{\text{or}} \rightarrow \text{Bil} \in \text{CAlg}(\mathcal{P}_{\Sigma}(\text{Corr}^{\text{fr}}(S)))$.*

The morphism $\text{FSyn}^{\text{or}} \rightarrow \text{Bil}$ is informally described as follows: a pair $(f : X \rightarrow Y$ finite syntomic, $\omega_f \simeq \mathcal{O})$ is sent to $f_*(\mathcal{O})$, where $\mathcal{O} \in \text{Bil}(X)$ denotes the vector bundle \mathcal{O}_X with its canonical symmetric bilinear pairing $\mathcal{O}_X \otimes \mathcal{O}_X \rightarrow \mathcal{O}_X$.

Proof. Denote by $K^{\circ} \in \mathcal{P}_{\Sigma}(\text{Sm}_S)$ the rank 0 part of the K -theory presheaf and by $\text{Corr}^{\text{fr}}((\text{Sm}_S)_{/K^{\circ}})$ the subcategory of the category constructed in [EHK⁺20b, §B] on objects (X, ξ) with $X \in \text{Sm}_S$, ξ of rank 0, and morphisms those spans whose left leg is finite syntomic. There are symmetric monoidal functors $\gamma : (\text{Sm}_S)_{/K^{\circ}} \rightarrow \text{Corr}^{\text{fr}}((\text{Sm}_S)_{/K^{\circ}})$ and $\delta : \text{Corr}^{\text{fr}}(S) \rightarrow \text{Corr}^{\text{fr}}((\text{Sm}_S)_{/K^{\circ}})$.

We first lift Bil to $\mathcal{P}_{\Sigma}(\text{Corr}^{\text{fr}}((\text{Sm}_S)_{/K^{\circ}}))$; we let $\text{Bil}(X, \xi)$ be the 1-groupoid of vector bundles with a symmetric bilinear form for the duality $\underline{\text{Hom}}(-, \det \xi)$. Since Bil is 1-truncated, we only need to specify a finite amount of coherence homotopies, so this can be done by hand. Since δ is symmetric monoidal δ^* is lax symmetric monoidal and hence $\delta^*(\text{Bil})$ produces the desired lift.

Let $K' \rightarrow K^{\circ}$ denote the fiber of the determinant map; in other words this is the rank 0 part of K^{SL} . This defines an object of $\text{CAlg}(\mathcal{P}_{\Sigma}(\text{Sm}_S)_{/K^{\circ}}) \simeq \text{CAlg}(\mathcal{P}_{\Sigma}((\text{Sm}_S)_{/K^{\circ}}))$ and $\gamma K' \simeq \text{FSyn}^{\text{or}}$ [EHK⁺20b, Example 3.3.4 and after Example 3.3.6]. To conclude the proof it hence suffices to construct a map $K' \rightarrow \gamma^*(\text{Bil}) \in \text{CAlg}(\mathcal{P}_{\Sigma}(\text{Sm}_S)_{/K^{\circ}})$. Again this only needs a finite amount of coherences; the desired map sends (ξ, ϕ) with $\xi \in K^{\circ}(X)$ and $\phi : \det(\xi) \simeq \mathcal{O}_X$ to (\mathcal{O}, ϕ') where $\phi' : \mathcal{O} \otimes \mathcal{O} \simeq \mathcal{O} \xrightarrow{\cong} \det \xi$. \square

Since group-completion and Zariski localization commute with the forgetful functor $\mathcal{P}_{\Sigma}(\text{Corr}_S^{\text{fr}}) \rightarrow \text{CMon}(\mathcal{P}_{\Sigma}(\text{Sm}_S)) \rightarrow \mathcal{P}_{\Sigma}(\text{Sm}_S)$ [EHK⁺17, Proposition 3.2.14], we deduce that $(L_{\text{Zar}} \text{Bil}^{\text{gp}})(X) \simeq \text{KO}(X)$ [Hor05, Definitions 1.5 and 2.2]; we denote this presheaf by GW . There is thus a canonical Bott element $\beta \in [T^4, \text{GW}]$.

Proposition A.2. *There is a canonical equivalence $(\Sigma_{\text{fr}}^{\infty} \text{GW})[\beta^{-1}] \simeq \text{KO}$, at least if $1/2 \in S$.*

Proof. The spectrum $(\Sigma_{\text{fr}}^{\infty} \text{GW})[\beta^{-1}] \in \mathcal{SH}^{\text{fr}}(S)$ can be modeled by the framed T^4 -prespectrum $(\text{GW}, \text{GW}, \dots)$ with the bonding maps given by multiplication by β . Under the equivalence $\mathcal{SH}^{\text{fr}}(S) \simeq \mathcal{SH}(S)$ [Hoy21], this corresponds to the same prespectrum with transfers forgotten. This is KO by definition. \square

In particular we have constructed an \mathcal{E}_{∞} -structure on KO.

Corollary A.3. *There is a morphism $\text{MSL} \rightarrow \text{KO} \in \text{CAlg}(\mathcal{SH}(S))$, at least if $1/2 \in S$.*

Proof. Take the morphism $\Sigma_{\text{fr}}^{\infty} \text{FSyn}^{\text{or}} \rightarrow \Sigma_{\text{fr}}^{\infty} \text{Bil} \rightarrow \Sigma_{\text{fr}}^{\infty} \text{GW} \rightarrow \text{KO}$ and use that $\Sigma_{\text{fr}}^{\infty} \text{FSyn}^{\text{or}} \simeq \text{MSL}$ [EHK⁺20b, Theorem 3.4.3(i)]. \square

Corollary A.4. *Let $f : Z \rightarrow S$ be a finite syntomic morphism, with $1/2 \in S$, $\tau : L_{Z/S} \simeq 0 \in K(Z)$ a trivialization, $\phi : V \otimes V \rightarrow \mathcal{O}_Z$ be a non-degenerate, symmetric bilinear form (defining an element $[\phi] \in \text{KO}^0(Z)$). Then*

$$[\hat{f}_*(\phi)] = f_*([\phi]) \in \text{KO}^0(S).$$

Here f_* denotes the transfer arising from the six functors formalism, and \hat{f}_* denotes the transfer constructed above using coherent duality.

Proof. We first give a simplified proof assuming that S is affine. Denote by $\mathrm{KO}^{\mathrm{fr}} \in \mathcal{SH}^{\mathrm{fr}}(S)$ a lift of KO . By [EHK⁺20a, Theorem 3.3.10], for any morphism $p : Z \rightarrow Y$ with $Y \in \mathrm{Sm}_S$ and a form ψ on Y , the six functors transfer of $p^*([\psi])$ along f coincides with the framed transfer of $[\psi]$ along the correspondence $S \xleftarrow{\tau} Z \rightarrow Y$. By Proposition A.2 we can take $\mathrm{KO}^{\mathrm{fr}} = \Sigma_{\mathrm{fr}}^{\infty} \mathrm{GW}[\beta^{-1}]$; it follows that there is a map $\mathrm{Bil} \rightarrow \Omega_{\mathrm{fr}}^{\infty} \mathrm{KO}^{\mathrm{fr}} \in \mathcal{P}_{\Sigma}(\mathrm{Corr}^{\mathrm{fr}})$. We thus deduce that

$$\hat{f}_*(p^*\psi) = f_*(p^*[\psi]) \in \mathrm{KO}^0(S).$$

The result would follow if there exist p, ψ with $p^*\psi = \phi$. Under our simplifying assumption that S is affine, this is always the case; see [EHK⁺20b, Proposition A.0.4 and Example A.0.6(5)].

To deal with the general case, we begin with some constructions. Given $F \in \mathcal{P}(\mathrm{Sm}_S)$, denote by $f^*F \in \mathcal{P}(\mathrm{Sm}_Z)$ the left Kan extension. If F comes from a presheaf with framed transfers, then the transfers along (base changes of) f assemble to a map $f_*f^*F \rightarrow F$. Given $E \in \mathcal{SH}(S)$, we obtain $\Omega^{\infty}E \in \mathcal{P}(\mathrm{Sm}_S)$ and can apply this construction. On the other hand there is a map $f_*\Omega^{\infty}f^*E \rightarrow \Omega^{\infty}E$ coming from the six functors transfer as well as a map $f^*\Omega^{\infty}E \rightarrow \Omega^{\infty}f^*E$, and [EHK⁺20a, Theorem 3.3.10] implies that the following diagram commutes

$$\begin{array}{ccc} f_*f^*\Omega^{\infty}E & \longrightarrow & f_*\Omega^{\infty}f^*E \\ \downarrow & \swarrow & \\ \Omega^{\infty}E & & \end{array}$$

Now we continue with the proof. We have a commutative diagram

$$\begin{array}{ccc} f_*f^*\mathrm{Bil}_S & \longrightarrow & f_*\mathrm{Bil}_Z \\ \downarrow & \swarrow & \\ \mathrm{Bil}_S & & \end{array}$$

here $\mathrm{Bil}_T \in \mathcal{P}(\mathrm{Sm}_T)$ denotes the stack of symmetric bilinear forms, the map $f_*f^*\mathrm{Bil}_S \rightarrow \mathrm{Bil}_S$ comes from the above construction with $F = \mathrm{Bil}_S$, the map $f_*\mathrm{Bil}_Z \rightarrow \mathrm{Bil}_S$ is the transfer \hat{f}_* , and commutativity holds by construction. Applying the above construction with $E = \mathrm{KO}$ and using its naturality in F , we obtain all in all the following commutative diagram

$$\begin{array}{ccccccc} f_*\mathrm{Bil}_Z & \xleftarrow{w} & f_*f^*\mathrm{Bil}_S & \longrightarrow & f_*f^*\Omega^{\infty}\mathrm{KO} & \longrightarrow & f_*\Omega^{\infty}f^*\mathrm{KO} \\ & \searrow & \downarrow & & \downarrow & \swarrow & \\ & & \mathrm{Bil}_S & \longrightarrow & \Omega^{\infty}\mathrm{KO} & & \end{array}$$

It follows from [EHK⁺20b, Proposition A.0.4 and Example A.0.6(5)] that $f^*\mathrm{Bil}_S \rightarrow \mathrm{Bil}_Z$ is a Zariski equivalence, and hence the map labelled w is a Nisnevich equivalence (f being finite). Since $f_*\Omega^{\infty}f^*\mathrm{KO}$ and $\Omega^{\infty}\mathrm{KO}$ are Nisnevich local, we may invert w above to obtain the following commutative diagram

$$\begin{array}{ccc} f_*\mathrm{Bil}_Z & \longrightarrow & f_*f^*\Omega^{\infty}\mathrm{KO} \\ \hat{f}_* \downarrow & & f_* \downarrow \\ \mathrm{Bil}_S & \longrightarrow & \Omega^{\infty}\mathrm{KO}. \end{array}$$

The result follows. □

Remark A.5. If $1/2 \notin S$, then we could define KO as $(\Sigma_{\mathrm{fr}}^{\infty} \mathrm{GW})[\beta^{-1}]$. Then Corollaries A.3 and A.4 (as well as Proposition A.2) remain true. The problem is that we no longer know what theory KO represents.

APPENDIX B. MISCELLANEOUS

We collect some results which we believe are well-known, but for which we could not find convenient references.

B.1. Cotangent complexes and dualizing complexes. For any morphism of schemes $f : X \rightarrow Y$, there is the cotangent complex $L_f \in D(X)$ (see e.g. [Sta18, Tag 08P5]). If f is lci, then L_f is perfect [Sta18, Tag 08SH] and hence defines a point $L_f \in K(X)$. Consequently in this case we can make sense of the graded determinant $\widetilde{\det}L_f \in \mathrm{Pic}(D(X))$; this is (locally) a shift of a line bundle.

Proposition B.1. *Let $f : X \rightarrow Y$ be lci. Then there is a canonical isomorphism $f^!\mathcal{O}_Y \simeq \widetilde{\det}L_f$.*

Recall that by our conventions, X and Y are separated and of finite type over some noetherian scheme S ; in particular they are themselves noetherian. We strongly believe that these assumptions are immaterial.

Proof. Both sides are compatible with passage to open subschemes of X . Locally on X , f factors as a regular immersion followed by a smooth morphism, say $f = pi$. For either a regular immersion or a smooth morphism g , we have isomorphisms

$$(18) \quad g^! \mathcal{O}_Y \simeq \widetilde{\det} L_g,$$

as desired [Har66, Definition III.2, Proposition 7.2]. For composable lci morphisms p, i , we have

$$(19) \quad (pi)^!(\mathcal{O}) \simeq i^* p^!(\mathcal{O}) \otimes i^!(\mathcal{O}).$$

Similarly we have a canonical cofiber sequence $i^* L_p \rightarrow L_{pi} \rightarrow L_i$ and hence

$$(20) \quad \widetilde{\det} L_{pi} \simeq i^* \widetilde{\det} L_p \otimes \widetilde{\det} L_i.$$

Combining (18), (19) and (20), we thus obtain an isomorphism

$$\alpha_{p,i} : f^! \mathcal{O}_X \simeq i^! p^! \mathcal{O}_X \simeq \widetilde{\det} L_f.$$

We have thus shown that $f^! \mathcal{O}_Y$ is locally isomorphic to $\widetilde{\det} L_f$ (via $\alpha_{p,i}$), and hence that $A := f^! \mathcal{O}_Y \otimes (\widetilde{\det} L_f)^{-1}$ is an \mathcal{O}_X -module concentrated in degree 0. Exhibiting an isomorphism as claimed is the same as exhibiting $A \simeq \mathcal{O}_X$, or equivalently a section $a \in \Gamma(X, A)$ which locally on X corresponds to an isomorphism. Since A is 0-truncated, we may construct a locally. In other words, we need to exhibit a cover $\{U_n\}_n$ of X and isomorphisms $\alpha_n : f^! \mathcal{O}_Y|_{U_n} \simeq \widetilde{\det} L_f|_{U_n}$ such that on $U_n \cap U_m$ we have $\alpha_n \simeq \alpha_m$. Hence, since we are claiming to exhibit a *canonical* isomorphism, we may do so locally on X . We may thus assume that f factors as pi , for a smooth morphism $p : V \rightarrow Y$ and a regular immersion $i : X \rightarrow V$. We have already found an isomorphism in this situation, namely $\alpha_{p,i}$. What remains to do is to show that this isomorphism is independent of the choice of factorization $f = pi$.

Thus let $i' : X \rightarrow V'$ and $p' : V' \rightarrow Y$ be another such factorization. We need to show that $\alpha_{p,i} = \alpha_{p',i'}$. By considering $V' \times_Y V$, we may assume given a commutative diagram

$$\begin{array}{ccccc} X & \xrightarrow{i'} & V' & \xrightarrow{p'} & Y \\ \parallel & & q \downarrow & & \parallel \\ X & \xrightarrow{i} & V & \xrightarrow{p} & Y, \end{array}$$

where q is smooth. If in (20) both f and g are smooth, then the isomorphism arises from the first fundamental exact sequence of Kähler differentials [Har66, Proposition III.2.2], and hence is the same as the isomorphism (19). It follows that we may assume that $p = \text{id}$. The isomorphism $i'^! q^! \simeq i^!$ is explained in [Har66, Proposition III.8.2] and reduces via formal considerations (that apply in the same way to $\widetilde{\det} L_-$) to the case of a smooth morphism with a section.

We are thus reduced to the following problem. Let $p : V \rightarrow X$ be smooth and $i : X \rightarrow V$ a regular immersion which is a section of p . The coherent duality formalism provides us with an isomorphism $\omega_{X/V} \otimes i^* \omega_{V/X} \simeq \mathcal{O}_X$; we need to check that this is the same as the isomorphism $\det L_i \otimes i^* \det L_p \simeq \det L_{\text{id}} = \mathcal{O}_X$ coming from (20). By [Har66, Lemma III.8.1, Definition III.1.5], the first isomorphism arises from the second fundamental exact sequence of Kähler differentials. This is the same as the second isomorphism.

This concludes the proof. \square

B.2. Grothendieck–Witt rings and Witt rings. Let R be a commutative ring. A *symmetric space over R* means a finitely generated projective R -module M together with a non-degenerate, symmetric bilinear form $\varphi : M \times M \rightarrow R$. The Grothendieck group on the semiring of isomorphism classes of symmetric spaces over R (with operations given by direct sum and tensor product) is denoted $\text{GW}(R)$ and called the *Grothendieck–Witt ring of R* . A symmetric space (M, φ) is called *metabolic* if there exists a summand $N \subset M$ with $N = N^\perp$. The quotient of $\text{GW}(R)$ by the subgroup (which is an ideal) generated by metabolic spaces is denoted $\text{W}(R)$ and called the *Witt ring of R* .

Lemma B.2. *Let R be a commutative ring and (M, φ) a symmetric space. The following are equivalent:*

- (1) M is metabolic
- (2) M contains an isotropic subspace of half rank: there exists a summand $N \subset M$ such that $\dim M = 2 \dim N$ and $\varphi|_N = 0$.

Now suppose that all finitely generated projective R -modules are free of constant rank, e.g. R a local ring. In this case the image of M in $\mathrm{GW}(R)$ is given by nh , where $\dim M = 2n$ and h denotes the hyperbolic plane, corresponding to the matrix $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, which also satisfies $h = 1 + \langle -1 \rangle \in \mathrm{GW}(R)$.

Moreover we have a pullback square

$$\begin{array}{ccc} \mathrm{GW}(R) & \longrightarrow & \mathbb{Z} \\ \downarrow & & \downarrow \\ \mathrm{W}(R) & \longrightarrow & \mathbb{Z}/2, \end{array}$$

where the horizontal maps are given by rank. In particular $\mathrm{GW}(R) \rightarrow \mathrm{W}(R) \times \mathbb{Z}$ is injective.

Proof. The equivalence of (1) and (2) is [Kne77, Corollary I.3.2]. The fact that the image in $\mathrm{GW}(R)$ is given by nh follows from [Kne77, Proposition I.3.2 and Corollary I.3.1].

Consider the symmetric space $M = \langle 1 \rangle \oplus h$. Thus $M \simeq R^3$ has basis e_1, e_2, e_3 with $\langle e_1, e_1 \rangle = 1$, $\langle e_1, e_2 \rangle = 0$, and so on. Direct computation shows that $f_1 = e_1 + e_2$, $f_2 = e_1 - e_3$, $f_3 = e_3 - e_1 - e_2$ is an orthogonal basis exhibiting $M \simeq \langle 1 \rangle \oplus \langle 1 \rangle \oplus \langle -1 \rangle$. Thus $h = 1 + \langle -1 \rangle \in \mathrm{GW}(R)$, as claimed.

We have $\mathrm{W}(R) = \mathrm{GW}(R)/J$, where the ideal J is generated by metabolic spaces. By the previous assertion, $J = \mathbb{Z} \cdot h$, and so the rank homomorphism maps J isomorphically onto $2\mathbb{Z}$. The pullback square follows formally. \square

Lemma B.3. *Let R be a local ring.*

- (1) *Let (M, φ) be a symmetric space over R . Then M admits an orthogonal basis if and only if there exists $m \in M$ with $\varphi(m, m) \in R^\times$.*
- (2) *$\mathrm{GW}(R)$ is generated by elements of the form $\langle a \rangle$, with $a \in R^\times$.*

Proof. (1) If e_1, \dots, e_n is an orthogonal basis then $\varphi(e_i, e_i) = a_i$ whereas $\varphi(e_i, e_j) = 0$ for $i \neq j$; it follows now from non-degeneracy that $a_1 \in R^\times$, and so the condition is necessary. Now suppose that $m \in M$ with $\varphi(m, m) \in R^\times$. By [MH73, Theorem I.3.2] we get $M \simeq Rm \oplus (Rm)^\perp$. Consider an isomorphism $M \simeq Re_1 \oplus \dots \oplus Re_n \oplus N$, where n is maximal. We wish to show that $N = 0$. We know that $n > 0$ (by existence of m) and if $x \in N$ then $(*) \varphi(x, x) \notin R^\times$ (because else we could split off Rx as before, contradicting maximality). Replacing M by $e_1R \oplus N$, we may assume that $n = 1$. Since $\varphi|_N$ is non-degenerate and R is local, if $N \neq 0$ there exist $y, z \in N$ with $\varphi(y, z) = 1$. Set $e'_1 = e_1 + y$ and $f = e_1 + \lambda z$ (with $\lambda \in R$). Then $\varphi(e'_1, e'_1) = \varphi(e_1, e_1) + \varphi(y, y) \in R^\times$ by $(*)$ (and using that R is local), and similarly $\varphi(f, f) \in R^\times$. On the other hand $\varphi(e'_1, f) = \varphi(e_1, e_1) + \lambda$, and hence there exists a (unique) value of λ such that $f \in (Re'_1)^\perp$. It follows that $M \simeq Re'_1 \oplus Rf \oplus N'$, in contradiction of maximality of n .

(2) If M is an inner product space then $M \oplus \langle 1 \rangle$ admits an orthogonal basis by (1), and hence $[M] = [M \oplus \langle 1 \rangle] - \langle 1 \rangle \in \mathrm{GW}(R)$ can be expressed in terms of elements of the form $\langle a \rangle$. \square

Corollary B.4. *We have $\mathrm{GW}(\mathbb{F}_2) = \mathbb{Z}$.*

Proof. It is immediate from Lemma B.3 that $\mathbb{Z} \rightarrow \mathrm{GW}(\mathbb{F}_2)$ is surjective. Since the rank provides a retraction, the map is also injective, hence an isomorphism. \square

B.3. Regular sequences.

Lemma B.5. *Let S be a scheme, $X \rightarrow S$ a smooth morphism, $f_1, \dots, f_n \in \mathcal{O}_X(X)$ and put $Z = Z(f_1, \dots, f_n)$. Assume that for all $s \in S$, either Z_s is empty or else $\dim Z_s \leq \dim X_s - n$. Then $Z \rightarrow S$ is flat and for each $z \in Z$, $f_1, \dots, f_n \in \mathcal{O}_{Z,z}$ is a strongly regular sequence. In particular f_1, \dots, f_n is a regular sequence and $Z \rightarrow X$ is a regular immersion.*

Proof. For $x \in X$ there exist affine open subschemes $x \in U \subset X$, $S' \subset S$ and a factorization $U \rightarrow \mathbb{A}_{S'}^d \rightarrow S' \rightarrow S$ of $U \rightarrow S$ with $U \rightarrow \mathbb{A}_{S'}^d$ étale [Sta18, Tag 01V4]. Then $Z' := Z \cap U \rightarrow S'$ still has fibers of dimension $\leq d - n$ and hence equal to $d - n$ by Krull's principal ideal theorem [Sta18, Tag 0BBZ]. This implies that $Z' \rightarrow S'$ is a relative global complete intersection (in the sense of [Sta18, Tag 00SP]) [EHK⁺17, Lemma 2.1.15], whence flat [Sta18, 00SW], and (f_1, \dots, f_n) is a strongly regular sequence in $\mathcal{O}_{Z',z'}$ for any $z' \in Z'$ [Sta18, Tag 00SV(1)]. Since x was arbitrary, it follows that $Z \rightarrow S$ is flat and (f_1, \dots, f_n) form a strongly regular sequence in $\mathcal{O}_{Z,z}$ for any $z \in Z$. For the last statement, we note that if $x \in X \setminus Z$ then $f_i \in \mathcal{O}_{X,x}$ is a unit for some i , and hence $K(f_\bullet)_x \simeq 0 \simeq (\mathcal{O}_Z)_x$, whereas if $x \in Z$ then $K(f_\bullet)_x \simeq \mathcal{O}_{Z,z}$ by strong regularity [Sta18, Tag 062F]. \square

Competing interests: The authors declare none.

REFERENCES

- [AB84] M. F. Atiyah and R. Bott. The moment map and equivariant cohomology. *Topology*, 23(1):1–28, 1984.
- [AF16] A. Ask and J. Fasel. Comparing Euler classes. *Q. J. Math.*, 67(4):603–635, 2016.
- [Ana15] Alexey Ananyevskiy. The special linear version of the projective bundle theorem. *Compos. Math.*, 151(3):461–501, 2015.
- [Ana20] Alexey Ananyevskiy. SL-oriented cohomology theories. In *Motivic homotopy theory and refined enumerative geometry*, volume 745 of *Contemp. Math.*, pages 1–19. Amer. Math. Soc., Providence, RI, [2020] ©2020.
- [Bac17] Tom Bachmann. The generalized slices of hermitian k-theory. *Journal of Topology*, 10(4):1124–1144, 2017. arXiv:1610.01346.
- [Bac18] Tom Bachmann. Some remarks on units in grothendieck–witt rings. *Journal of Algebra*, 499:229 – 271, 2018. arXiv:1707.08087.
- [Bac20] Tom Bachmann. η -periodic motivic stable homotopy theory over dedekind domains. arXiv:2006.02086, 2020.
- [BBM⁺21] Thomas Brazelton, Robert Burklund, Stephen McKean, Michael Montoro, and Morgan Opie. The trace of the local \mathbb{A}^1 -degree. *Homology Homotopy Appl.*, 23(1):243–255, 2021.
- [BF17] Tom Bachmann and Jean Fasel. On the effectivity of spectra representing motivic cohomology theories. 2017. arXiv:1710.00594.
- [BG1⁺71] Pierre Berthelot, Alexandre Grothendieck, Luc Illusie, et al. *Théorie des intersections et théorème de Riemann-Roch*, volume 225. Springer, 1971.
- [BH98] Winfried Bruns and H Jürgen Herzog. *Cohen-macaulay rings*. Cambridge university press, 1998.
- [BH17] Tom Bachmann and Marc Hoyois. Norms in motivic homotopy theory. 2017. arXiv:1711.03061.
- [BH20] Tom Bachmann and Michael J. Hopkins. η -periodic motivic stable homotopy theory over fields. arXiv:2005.06778, 2020.
- [BKW20] Candace Bethea, Jesse Leo Kass, and Kirsten Wickelgren. Examples of wild ramification in an enriched Riemann–Hurwitz formula. In *Motivic homotopy theory and refined enumerative geometry*, volume 745 of *Contemp. Math.*, pages 69–82. Amer. Math. Soc., Providence, RI, [2020] ©2020.
- [BM00] Jean Barge and Fabien Morel. Groupe de Chow des cycles orientés et classe d’Euler des fibrés vectoriels. *C. R. Acad. Sci. Paris Sér. I Math.*, 330(4):287–290, 2000.
- [BØ21] Tom Bachmann and Paul Arne Østvær. Topological models for stable motivic invariants of regular number rings. arXiv:2102.01618, 2021.
- [BW20] Olivier Benoist and Olivier Wittenberg. On the integral Hodge conjecture for real varieties, I. *Invent. Math.*, 222(1):1–77, 2020.
- [CD19] Denis-Charles Cisinski and Frédéric Déglise. *Triangulated categories of mixed motives*. Springer Monographs in Mathematics. Springer, Cham, [2019] ©2019.
- [CDH⁺20] Baptiste Calmès, Emanuele Dotto, Yonatan Harpaz, Fabien Hebestreit, Markus Land, Kristian Moi, Denis Nardin, Thomas Nikolaus, and Wolfgang Steimle. Hermitian k-theory for stable ∞ -categories iii. ArXiv 2009.07225, 2020.
- [CF17a] Baptiste Calmès and Jean Fasel. A comparison theorem for mw-motivic cohomology. *arXiv preprint arXiv:1708.06100*, 2017.
- [CF17b] Baptiste Calmès and Jean Fasel. Finite chow–witt correspondences. *arXiv preprint*, 2017. arXiv:1412.2989.
- [CH09] Baptiste Calmès and Jens Hornbostel. Tensor-triangulated categories and dualities. *Theory Appl. Categ.*, 22(6):136–200, 2009.
- [CH11] Baptiste Calmès and Jens Hornbostel. Push-forwards for Witt groups of schemes. *Comment. Math. Helv.*, 86(2):437–468, 2011.
- [Con00] Brian Conrad. *Grothendieck duality and base change*. Number 1750. Springer Science & Business Media, 2000.
- [DJK18] Frédéric Déglise, Fangzhou Jin, and Adeel A. Khan. Fundamental classes in motivic homotopy theory. *Preprint*, available at <https://arxiv.org/abs/1805.05920>, 2018.
- [DM98] Olivier Debarre and Laurent Manivel. Sur la variété des espaces linéaires contenus dans une intersection complète. *Math. Ann.*, 312(3):549–574, 1998.
- [EH16] David Eisenbud and Joe Harris. *3264 and all that—a second course in algebraic geometry*. Cambridge University Press, Cambridge, 2016.
- [EHK⁺17] Elden Elmanto, Marc Hoyois, Adeel A. Khan, Vladimir Sosnilo, and Maria Yakerson. Motivic infinite loop spaces. arXiv preprint 1711.05248, 2017.
- [EHK⁺20a] Elden Elmanto, Marc Hoyois, Adeel A. Khan, Vladimir Sosnilo, and Maria Yakerson. Framed transfers and motivic fundamental classes. *J. Topol.*, 13(2):460–500, 2020.
- [EHK⁺20b] Elden Elmanto, Marc Hoyois, Adeel A. Khan, Vladimir Sosnilo, and Maria Yakerson. Modules over algebraic cobordism. *Forum Math. Pi*, 8:e14, 44, 2020.
- [Eis95] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.
- [EL77] David Eisenbud and Harold I. Levine. An algebraic formula for the degree of a C^∞ map germ. *Ann. of Math. (2)*, 106(1):19–44, 1977. With an appendix by Bernard Teissier, “Sur une inégalité à la Minkowski pour les multiplicités”.
- [Fas13] Jean Fasel. The projective bundle theorem for \mathbb{P}^j -cohomology. *J. K-Theory*, 11(2):413–464, 2013.
- [FK13] Sergey Finashin and Viatcheslav Kharlamov. Abundance of real lines on real projective hypersurfaces. *Int. Math. Res. Not. IMRN*, (16):3639–3646, 2013.
- [FK15] S. Finashin and V. Kharlamov. Abundance of 3-planes on real projective hypersurfaces. *Arnold Math. J.*, 1(2):171–199, 2015.
- [FK21] Sergey Finashin and Viatcheslav Kharlamov. Segre Indices and Welschinger Weights as Options for Invariant Count of Real Lines. *Int. Math. Res. Not. IMRN*, (6):4051–4078, 2021.
- [Ful84] W. Fulton. *Intersection theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer-Verlag, 1984.

- [GP99] T. Graber and R. Pandharipande. Localization of virtual classes. *Invent. Math.*, 135(2):487–518, 1999.
- [GP18] Owen Gwilliam and Dmitri Pavlov. Enhancing the filtered derived category. *Journal of Pure and Applied Algebra*, 222(11):3621–3674, 2018.
- [Har66] Robin Hartshorne. *Residues and duality*, volume 20. Springer, 1966.
- [Har13] Robin Hartshorne. *Algebraic geometry*, volume 52. Springer Science & Business Media, 2013.
- [Hor05] Jens Hornbostel. \mathbb{A}^1 -representability of hermitian k -theory and witt groups. *Topology*, 44(3):661–687, 2005.
- [Hoy17] Marc Hoyois. The six operations in equivariant motivic homotopy theory. *Advances in Mathematics*, 305:197–279, 2017.
- [Hoy21] Marc Hoyois. The localization theorem for framed motivic spaces. *Compos. Math.*, 157(1):1–11, 2021.
- [HWXZ19] Jens Hornbostel, Matthias Wendt, Heng Xie, and Marcus Zibrowius. The real cycle class map. *arXiv preprint arXiv:1911.04150*, 2019.
- [Ill71] L. Illusie. *Complexe cotangent et déformations*. Number v. 1 in Lecture notes in mathematics. Springer-Verlag, 1971.
- [Jac17] Jeremy Jacobson. Real cohomology and the powers of the fundamental ideal in the witt ring. *Annals of K-Theory*, 2(3):357–385, 2017.
- [Khi77] G. N. Khimshiashvili. The local degree of a smooth mapping. *Sakharth. SSR Mecn. Akad. Moambe*, 85(2):309–312, 1977.
- [Kne77] Manfred Knebusch. Symmetric bilinear forms over algebraic varieties. In G. Orzech, editor, *Conference on quadratic forms*, volume 46 of *Queen’s papers in pure and applied mathematics*, pages 103–283. Queens University, Kingston, Ontario, 1977.
- [Knu91] Max-Albert Knus. *Quadratic and Hermitian forms over rings*, volume 294 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1991. With a foreword by I. Bertuccioni.
- [KW19] Jesse Leo Kass and Kirsten Wickelgren. The class of Eisenbud–Khimshiashvili–Levine is the local \mathbb{A}^1 -brouwer degree. *Duke Mathematical Journal*, 168(3):429–469, 2019.
- [KW21] Jesse Leo Kass and Kirsten Wickelgren. An arithmetic count of the lines on a smooth cubic surface. *Compos. Math.*, 157(4):677–709, 2021.
- [LÁ17] Alejo López-Ávila. *E_∞ -ring Structures in Motivic Hermitian K -theory*. PhD thesis, Universität Osnabrück, 2017.
- [Lam05] T. Y. Lam. *Introduction to quadratic forms over fields*, volume 67 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2005.
- [Lev19] Marc Levine. Motivic Euler characteristics and Witt-valued characteristic classes. *Nagoya Mathematical Journal*, 236:251 – 310, 2019. Special Issue: Celebrating the 60th birthday of Shuji Saito.
- [Lev20] Marc Levine. Aspects of enumerative geometry with quadratic forms. *Doc. Math.*, 25:2179–2239, 2020.
- [LR20] Marc Levine and Arpon Raksit. Motivic Gauss-Bonnet formulas. *Algebra Number Theory*, 14(7):1801–1851, 2020.
- [McK21] Stephen McKean. An arithmetic enrichment of Bézout’s Theorem. *Math. Ann.*, 379(1-2):633–660, 2021.
- [MH73] John Willard Milnor and Dale Husemoller. *Symmetric bilinear forms*, volume 60. Springer, 1973.
- [Mor05] Fabien Morel. The stable \mathbb{A}^1 -connectivity theorems. *K-theory*, 35(1):1–68, 2005.
- [Mor12] Fabien Morel. *\mathbb{A}^1 -Algebraic Topology over a Field*. Lecture Notes in Mathematics. Springer Berlin Heidelberg, 2012.
- [OT14] Christian Okonek and Andrei Teleman. Intrinsic signs and lower bounds in real algebraic geometry. *J. Reine Angew. Math.*, 688:219–241, 2014.
- [Pau20] Sabrina Pauli. Quadratic types and the dynamic euler number of of lines on a quintic 3-fold. ArXiv preprint 2006.12089, 2020.
- [PW10a] Ivan Panin and Charles Walter. On the algebraic cobordism spectra $m\mathbb{S}l$ and $m\mathbb{S}p$. *arXiv preprint arXiv:1011.0651*, 2010.
- [PW10b] Ivan Panin and Charles Walter. Quaternionic grassmannians and pontryagin classes in algebraic geometry. *arXiv preprint arXiv:1011.0649*, 2010.
- [PW18] I. Panin and C. Walter. On the motivic commutative ring spectrum **BO**. *Algebra i Analiz*, 30(6):43–96, 2018.
- [Sch10] Marco Schlichting. The mayer-vietoris principle for grothendieck-witt groups of schemes. *Inventiones mathematicae*, 179(2):349–433, 2010.
- [Sol06] Jake P. Solomon. Intersection theory on the moduli space of holomorphic curves with Lagrangian boundary conditions. *PhD Thesis*, available at <https://arxiv.org/abs/math/0606429>, 2006.
- [Spi18] Markus Spitzweck. A commutative \mathbb{P}^1 -spectrum representing motivic cohomology over Dedekind domains. *Mém. Soc. Math. Fr. (N.S.)*, (157):110, 2018.
- [SS75] Günter Scheja and Uwe Storch. Über Spurfunktionen bei vollständigen Durchschnitten. *J. Reine Angew. Math.*, 278/279:174–190, 1975.
- [ST15] Marco Schlichting and Girja S Tripathi. Geometric models for higher grothendieck–witt groups in \mathbb{A}^1 -homotopy theory. *Mathematische Annalen*, 362(3-4):1143–1167, 2015.
- [Sta18] The Stacks Project Authors. *Stacks Project*. <http://stacks.math.columbia.edu>, 2018.
- [SW21] Padmavathi Srinivasan and Kirsten Wickelgren. An arithmetic count of the lines meeting four lines in P^3 . *Trans. Amer. Math. Soc.*, 374(5):3427–3451, 2021.
- [Wen20] Matthias Wendt. Oriented Schubert calculus in Chow–Witt rings of Grassmannians. In *Motivic homotopy theory and refined enumerative geometry*, volume 745 of *Contemp. Math.*, pages 217–267. Amer. Math. Soc., Providence, RI, [2020] ©2020.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA, USA
Email address: `tom.bachmann@zoho.com`

CURRENT: K. WICKELGREN, DEPARTMENT OF MATHEMATICS, DUKE UNIVERSITY, DURHAM, NC, USA
Email address: `kirsten.wickelgren@duke.edu`